

8-2009

# Review: Transport Losses in Market Weight Pigs: I. A Review of Definitions, Incidence, and Economic Impact

M. J. Ritter

*University of Illinois at Urbana-Champaign*

M. Ellis

*University of Illinois at Urbana-Champaign*

N. L. Berry

*Cargill Inc.*

S E. Curtis

*University of Illinois at Urbana-Champaign*

L. Anil

*University of Minnesota - Twin Cities*

Follow this and additional works at: [http://lib.dr.iastate.edu/ans\\_pubs](http://lib.dr.iastate.edu/ans_pubs)

 *next page for additional authors*  
Part of the [Agriculture Commons](#), and the [Meat Science Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/ans\\_pubs/96](http://lib.dr.iastate.edu/ans_pubs/96). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Review: Transport Losses in Market Weight Pigs: I. A Review of Definitions, Incidence, and Economic Impact

## Abstract

Transport losses (dead and nonambulatory pigs) present animal welfare, legal, and economic challenges to the US swine industry. The objectives of this review are to explore 1) the historical perspective of transport losses; 2) the incidence and economic implications of transport losses; and 3) the symptoms and metabolic characteristics of fatigued pigs. In 1933 and 1934, the incidence of dead and nonambulatory pigs was reported to be 0.08 and 0.16%, respectively. More recently, 23 commercial field trials (n = 6,660,569 pigs) were summarized and the frequency of dead pigs, nonambulatory pigs, and total transport losses at the processing plant were 0.25, 0.44, and 0.69% respectively. In 2006, total economic losses associated with these transport losses were estimated to cost the US pork industry approximately \$46 million. Furthermore, 0.37 and 0.05% of the nonambulatory pigs were classified as either fatigued (nonambulatory, noninjured) or injured, respectively, in 18 of these trials (n = 4,966,419 pigs). Fatigued pigs display signs of acute stress (open-mouth breathing, skin discoloration, muscle tremors) and are in a metabolic state of acidosis, characterized by low blood pH and high blood lactate concentrations; however, the majority of fatigued pigs will recover with rest. Transport losses are a multifactorial problem consisting of people, pig, facility design, management, transportation, processing plant, and environmental factors, and, because of these multiple factors, continued research efforts are needed to understand how each of the factors and the relationships among factors affect the well-being of the pig during the marketing process.

## Keywords

Dead, fatigued, nonambulatory, pig, transport

## Disciplines

Agriculture | Animal Sciences | Meat Science

## Comments

This article is from *Professional Animal Scientist* 25 (2009): 404.

## Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

## Authors

M. J. Ritter, M. Ellis, N. L. Berry, S E. Curtis, L. Anil, E. Berg, M. Benjamin, D. Butler, C. Dewey, B Driessen, Paul R. DuBois, J. D. Hill, J. N. Marchant-Forde, P. Matzat, John J. McGlone, P. Mormede, T. Moyer, K. Pfalzgraf, J. Salak-Johnson, M. Siemens, J. Sterle, C. Stull, T. Whiting, B. Wolter, S. R. Niekamp, and Anna K. Johnson

## REVIEWS

# REVIEW: Transport Losses in Market Weight Pigs: I. A Review of Definitions, Incidence, and Economic Impact

M. J. Ritter,<sup>\*1,2</sup> M. Ellis,<sup>\*</sup> N. L. Berry,<sup>†</sup> S. E. Curtis,<sup>\*</sup> L. Anil,<sup>‡</sup> E. Berg,<sup>§</sup> M. Benjamin,<sup>#</sup> D. Butler, C. Dewey,<sup>¶</sup> B. Driessen,<sup>\*\*</sup> P. DuBois,<sup>††</sup> J. D. Hill,<sup>‡‡</sup> J. N. Marchant-Forde,<sup>§§</sup> P. Matzat,<sup>###</sup> J. McGlone, P. Mormede,<sup>¶¶</sup> T. Moyer,<sup>\*\*\*</sup> K. Pfalzgraf,<sup>†††</sup> J. Salak-Johnson,<sup>\*</sup> M. Siemens,<sup>††</sup> J. Sterle,<sup>‡‡‡</sup> C. Stull,<sup>§§§</sup> T. Whiting,<sup>####</sup> B. Wolter, S. R. Niekamp,<sup>¶¶¶</sup> and A. K. Johnson<sup>\*\*\*\*</sup>

<sup>\*</sup>Department of Animal Sciences, University of Illinois, IL 61801; <sup>†</sup>Cargill Inc., Coralville, IA 52241; <sup>‡</sup>Department of Veterinary Population Medicine, University of Minnesota, MN 55108; <sup>§</sup>Department of Animal Sciences, North Dakota State University, ND 58108; <sup>#</sup>Veterinary Services Consulting, Millarville, Alberta, Canada T0L 1R0; <sup>¶</sup>Murphy-Brown LLC, NC 28398; <sup>¶¶</sup>Department of Population Medicine, University of Guelph, Ontario, Canada N1G2W1; <sup>\*\*</sup>K. H. Kempen and Zootechnical Centre-K. U. Leuven, Belgium B-3360; <sup>††</sup>Cargill Meat Solutions, Wichita, KS 67202; <sup>‡‡</sup>Innovative Livestock Solutions, Blackie, Alberta, Canada T0L0J0; <sup>§§</sup>Livestock Behavior Research Unit, USDA-ARS, West Lafayette, IN 47907; <sup>###</sup>Elanco Animal Health, Greenfield, IN 46140; <sup>||</sup> || Pork Industry Institute, Department of Animal and Food Sciences, Texas Tech University, Lubbock 79409; <sup>¶¶¶</sup>National Institute for Agricultural Research, University of Bordeaux, Bordeaux, France; <sup>\*\*\*</sup>Hatfield Quality Meats, Hatfield, PA 19440; <sup>†††</sup>Tyson Fresh Meats, Springdale, AR 72762; <sup>‡‡‡</sup>Department of Animal Science, Texas AgriLife Extension, Texas A&M University, College Station 77843; <sup>§§§</sup>School of Veterinary Medicine, University of California, Davis 95616; <sup>####</sup>Veterinary Services Branch, Manitoba Agriculture and Food, Winnipeg, Manitoba, Canada R3T 5S6; <sup>||</sup> || || The Maschhoffs, Carlyle, IL 62231; <sup>¶¶¶</sup>National Pork Board, Clive, IA 50325; <sup>\*\*\*\*</sup>Department of Animal Science, Iowa State University, Ames 50011

## ABSTRACT

*Transport losses (dead and nonambulatory pigs) present animal welfare, legal, and economic challenges to the US swine industry. The objectives of this review are to explore 1) the historical perspective of transport losses; 2) the incidence*

*and economic implications of transport losses; and 3) the symptoms and metabolic characteristics of fatigued pigs. In 1933 and 1934, the incidence of dead and nonambulatory pigs was reported to be 0.08 and 0.16%, respectively. More recently, 23 commercial field trials (n = 6,660,569 pigs) were summarized and the frequency of dead pigs, nonambulatory pigs, and total transport losses at the processing plant were 0.25, 0.44, and 0.69% respectively. In 2006, total economic losses associated with these trans-*

*port losses were estimated to cost the US pork industry approximately \$46 million. Furthermore, 0.37 and 0.05% of the non-ambulatory pigs were classified as either fatigued (nonambulatory, noninjured) or injured, respectively, in 18 of these trials (n = 4,966,419 pigs). Fatigued pigs display signs of acute stress (open-mouth breathing, skin discoloration, muscle tremors) and are in a metabolic state of acidosis, characterized by low blood pH and high blood lactate concentrations; however, the majority of fatigued pigs*

<sup>1</sup>Corresponding author: [ritterma@lilly.com](mailto:ritterma@lilly.com)

<sup>2</sup>Current address: Elanco Animal Health, 56776 241st Street, Suite 200, Ames, IA 50010.

will recover with rest. Transport losses are a multifactorial problem consisting of people, pig, facility design, management, transportation, processing plant, and environmental factors, and, because of these multiple factors, continued research efforts are needed to understand how each of the factors and the relationships among factors affect the well-being of the pig during the marketing process.

**Key words:** dead, fatigued, nonambulatory, pig, transport

## INTRODUCTION

Dead and nonambulatory pigs at the processing plant are a multifactorial problem that can be influenced by pig, facility design, people, management, transportation, processing plant, and environmental factors (Anderson et al., 2002; Ellis et al., 2003; Ellis and Ritter, 2005a), and these losses represent multiple challenges for the entire US food chain. First, improving the welfare of finished pigs during transport and reducing the incidence of dead and nonambulatory pigs has become an animal welfare priority (National Pork Board, 2007). Second, increasingly strict rules, regulations, and enforcement are being considered for nonambulatory livestock [e.g., the Downed Animal and Food Safety Protection Act—Bill H. R. 661 (US House of Representatives, 2007) and Bill S. 394 (US Senate, 2007)]. Third, transport losses represent direct financial losses to pig producers and pork processors. Dead

and nonambulatory pigs have been estimated to cost the US pork industry \$50 to \$100 million annually (Ellis et al., 2003).

In 2004, the Animal Welfare Committee of the National Pork Board sponsored a workshop that involved scientists, government officials, and industry representatives from North America and Europe to review the scientific literature pertaining to transport losses in market weight pigs. The goals of this workshop were to identify key gaps in current knowledge and to identify future research needs. Therefore, the specific objectives of this paper are 1) to discuss pig transport losses from a historical perspective; 2) to determine the incidence and economic impact of transport losses in the United States; and 3) to describe the symptoms and metabolic characteristics of fatigued pigs.

## REVIEW AND DISCUSSION

### Defining Transport Losses

The term “transport losses” refers to pigs that die or become nonambulatory at any stage of the marketing process, defined as movement from the grower-finisher environment to stunning at the abattoir. Pigs that die during transport are referred to as “dead on arrival,” whereas pigs that die after having been unloaded are termed “dead in yard” or “dead in pen.” A “nonambulatory pig” is a pig unable to move or keep up with its contemporaries at the processing

plant (Anderson et al., 2002). Several terms are used throughout the industry for nonambulatory pigs, including cripples, slows, stressors, subjects, and suspects.

Two types of nonambulatory pigs commonly observed are fatigued and injured pigs (Ellis and Ritter, 2005a,b). “Fatigued pigs” are pigs that, without obvious injury, trauma, or disease, refuse to walk or keep up with their contemporaries at any stage of the marketing process (Ritter et al., 2005). In addition, “injured pigs” are those that become nonambulatory or that have a compromised ability to ambulate because of structural unsoundness or an injury sustained before or during the marketing process (Ellis and Ritter, 2005b).

### Historical Perspective

Transport losses in pigs during marketing from the farm to the processing plant are not a new issue in the swine industry. In fact, the National Livestock Loss Prevention Board was developed in 1934 to monitor losses occurring in livestock during transportation. In 1933 and 1934, the National Livestock Loss Prevention Board monitored the incidence of dead and nonambulatory pigs for shipment by rail and by truck at 17 livestock markets, and the incidence of dead and nonambulatory pigs was reported as 0.08 and 0.16%, respectively (Smith, 1937; Table 1).

The National Livestock Loss Prevention Board also continued to monitor the incidence of dead and nonambulatory pigs for deliveries by truck (Table 2) and by rail (Table 3) at the Omaha Livestock Market from 1934 to 1951. During that period, the incidence of dead pigs ranged from 0.07 to 0.19%, whereas the incidence of nonambulatory pigs ranged from 0.14 to 0.29%, with total losses ranging from 0.27 to 0.46% for deliveries by truck (Table 2). For deliveries by rail, the incidence of dead pigs ranged from 0.01 to 0.09%, that of nonambulatory pigs ranged from 0.11 to 0.21%, and that of total losses ranged from 0.14 to 0.29% (Table 3).

**Table 1. National Livestock Loss Prevention Board survey of losses at 17 hog markets in 1933 and 1934<sup>1</sup>**

Delivery method	Year	Pigs received, no.	Dead, %	Nonambulatory, %	Total losses, <sup>2</sup> %
By rail	1933	4,883,427	0.078	0.140	0.218
	1934	2,836,588	0.079	0.147	0.226
By truck	1933	12,192,828	0.080	0.166	0.246
	1934	9,294,602	0.101	0.195	0.296

<sup>1</sup>Adapted from Smith (1937).

<sup>2</sup>Total losses = dead + nonambulatory pigs.

**Table 2. A summary of transport losses collected by the National Livestock Loss Prevention Board on deliveries by truck at the Omaha Livestock Market from 1934 to 1951<sup>1,2,3</sup>**

Year	Pigs received, no.	Dead, %	Nonambulatory, %	Total losses, <sup>4</sup> %
1934	2,485,148	0.12	0.20	0.32
1935	1,061,945	0.10	0.24	0.33
1936	1,574,254	0.10	0.24	0.33
1937	842,056	0.07	0.20	0.27
1938	824,519	0.07	0.28	0.35
1939	1,128,145	0.08	0.23	0.31
1940	1,412,936	0.09	0.22	0.31
1941	1,022,623	0.10	0.29	0.39
1942	1,283,330	0.12	0.26	0.38
1943	1,992,810	0.16	0.22	0.39
1944	2,615,133	0.14	0.18	0.32
1945	1,121,004	0.18	0.24	0.42
1946	1,471,496	0.19	0.27	0.46
1947	1,799,590	0.16	0.21	0.37
1950	2,475,122	0.17	0.24	0.41
1951	2,980,217	0.16	0.18	0.34

<sup>1</sup>Spencer (1942): data from 1934 to 1942.

<sup>2</sup>Spencer (1947): data from 1943 to 1947.

<sup>3</sup>Spencer (1951): data from 1950 and 1951.

<sup>4</sup>Total losses = dead + nonambulatory pigs.

Smith (1937) provided recommendations for reducing transport losses in pigs during marketing. Interestingly, the following recommendations generally hold true today:

1. The railcar or truck should be cleaned and properly bedded before loading.
2. In hot weather, use sand for bedding, soak the sand before loading, and shower the pigs during transport. Do not apply a heavy stream of cold water to the back of a hot pig as this can result in death.
3. In the winter, use straw or hay for bedding and line the inside of the truck or car with heavy building paper to protect the pigs from cold air.
4. Withdraw feed before loading and do not feed pigs during transportation.
5. Handle pigs quietly and calmly.
6. Do not use sticks, clubs, whips, or prods to move pigs as these handling tools may cause carcass bruising. Instead, use

canvas “flappers” to minimize carcass bruising.

7. If possible, do not mix unfamiliar pigs during transport.
8. Do not overcrowd pigs during transport, especially in hot weather.

In the 1960s, producers, researchers, and veterinarians reported unexplained deaths in heavy-muscled pigs during routine handling and transportation (Topel et al., 1968). From a survey conducted by the Livestock Conservation Institute, the percentage of pigs that died during transport reportedly increased by 61% from 1964 to 1972, and the incidence of nonambulatory pigs increased by 17% during that same period (Rosse, 1972). It was later determined that most of these unexplained deaths were attributed to porcine stress syndrome (Topel et al., 1968; Topel and Christian, 1981).

During the 1980s and 1990s, the research focus was more on the general well-being of pigs in relation to journey times and stocking densities. The

information gathered during these decades has largely shaped legislation on the transportation of livestock in the European Union (European Economic Community, 1991; European Commission, 1995). More recently, the focus has shifted back toward transport losses, and several research projects from 2000 to 2007 have focused on the problem areas identified by Smith (1937). These efforts included evaluating the effects of handling tools (McGlone et al., 2004), handling intensity (Hamilton et al., 2004), feed withdrawal before loading (Ritter, 2007), mixing unfamiliar pigs during transport (Ritter, 2007), and transport floor space (Ritter, 2007) on the stress responses and transport losses of pigs during the marketing process. Furthermore, priorities of the swine industry included the development of water-application recommendations during hot weather and boarding and bedding recommendations for cold weather to provide optimal care and comfort for pigs during transportation (National Pork Board, 2007).

### *Incidence of Transport Losses*

**Dead Pigs.** The percentages of dead pigs at USDA-inspected plants are reported by the Food Safety and Inspection Service as “swine condemned ante-mortem for deads,” and these national statistics are available to the public via the Freedom of Information Act. The yearly incidence of dead market pigs at USDA-inspected plants for calendar years 1991 through 2006 is presented in Figure 1 (Food Safety and Inspection Service, 2007b). The incidence of dead market swine at US plants was very low in 1991 (0.08%) and 1992 (0.07%); however, the percentage of dead pigs at US plants increased 3-fold between 1993 and 1998 (0.10 and 0.30%, respectively). It is unclear why this value increased greatly over this period, but some potential explanations include changes in genetics, increased slaughter weights, and increased size of production operations (Ellis et al., 2003). For example, the national average for slaughter weights in the



United States increased from 114.8 kg in 1993 to 117.5 kg in 1999 (US-DA-National Agricultural Statistics Service, 2008).

From 1998 to 2001, the percentage of dead pigs peaked and remained relatively constant (at 0.28 to 0.30%), whereas from 2001 to 2002, the percentage of dead pigs at US plants decreased from 0.29 to 0.22%, where it remains today. This decrease can be attributed to greater industry awareness of losses during the marketing process. In 2002, the Transport Quality Assurance program of the National Pork Board was made available, and there was a concerted focus on research that yielded important knowledge (National Pork Board, 2008a,b).

Beginning in about 2005, several pork processors began euthanizing nonambulatory pigs on the trailer and in the plant that had a low likelihood of recovering (e.g., pigs with broken legs; pigs with a rectal temperature greater than 41.1°C) to minimize stress and suffering (M. Siemens, Cargill Meat Solutions, Wichita, KS,

personal communication). In most cases, nonambulatory pigs that are killed are recorded as dead pigs; therefore, the definition of a dead pig at US processing plants has now been altered to include pigs that are killed at the plant.

A summary of the incidence of dead pigs at the processing plant for 1960 through 2004 is presented by country in Table 4. The current US national average for percentage of pigs dead at the processing plant (0.22%; Figure 1) is similar to values reported for Canada from 1996 to 2004 (range: 0.07 to 0.17%) and Spain (range: 0.15 to 0.33%), but is higher than values reported for Denmark from 1996 to 2002 (0.02%), the Czech Republic from 1997 through 2004 (0.11%), for France during 1995 and 1997 (0.08 and 0.07%, respectively), for Germany from 1999 through 2003 (range: 0.10 to 0.17%), and for the United Kingdom from 1990 through 1994 (range: 0.05 to 0.11%).

**Nonambulatory Pigs.** Currently, national statistics are not available for the incidence of nonambulatory pigs

at US processing plants. However, 23 commercial field trials have reported incidences of dead pigs, nonambulatory pigs, and total losses (dead and nonambulatory pigs) at US processing plants that have occurred before pigs reached the weigh scale (Table 5). The weighted averages across the 23 field studies (n = 6,660,569 pigs) were 0.25% for dead pigs (range: 0.00 to 0.77%), 0.44% for nonambulatory pigs (range: 0.11 to 2.34%), and 0.69% for total losses (range: 0.14 to 2.39%). Furthermore, nonambulatory pigs at the processing plant were classified as fatigued or injured in 18 of these 23 field studies (Table 5). The weighted averages across the 18 studies (n = 4,966,419 pigs) were 0.37% fatigued (range: 0.05 to 1.98%) and 0.05% injured (range: 0.04 to 0.36%), suggesting a 7:1 ratio of fatigued to injured pigs at the processing plant. Additionally, the percentage of nonambulatory pigs at the farm was reported in only 12 of the 23 studies (n = 101,417 pigs), and the weighted average for nonambulatory pigs at the farm across those studies was 0.11% (range: 0.00 to 0.47%).

The incidence of nonambulatory pigs in other countries has been poorly documented as well. Sunstrum et al. (2006) monitored transport losses in Ontario in 2003 and reported that the percentage of subject pigs at the plant was 0.27%, which included 0.13% fatigued pigs, 0.12% injured pigs, and 0.02% cull pigs (e.g., prolapses, intact males, severe bruises, tail bites, etc.). The literature is apparently devoid of any reports of nonambulatory pigs in European countries. However, Averós et al. (2008) recently monitored 739 journeys to 37 processing plants in 5 different European Union countries and reported that the percentages of dead and injured pigs during unloading were 0.11 and 0.36%, respectively. Interestingly, these authors included pigs with skin blemishes and lameness in their definition of injured pigs. Therefore, it is unclear if any of these injured pigs became nonambulatory at the processing plant.

**Table 3. A summary of transport losses collected by the National Livestock Loss Prevention Board on deliveries by rail at the Omaha Livestock Market from 1934 to 1951<sup>1,2,3</sup>**

Year	Pigs received, no.	Dead, %	Nonambulatory, %	Total losses, <sup>4</sup> %
1934	322,584	0.04	0.13	0.18
1935	78,263	0.06	0.11	0.17
1936	188,534	0.02	0.12	0.14
1937	267,626	0.02	0.15	0.18
1938	389,852	0.04	0.21	0.25
1939	501,131	0.01	0.14	0.16
1940	648,573	0.03	0.16	0.19
1941	677,410	0.02	0.18	0.20
1942	934,490	0.03	0.17	0.20
1943	1,031,280	0.04	0.13	0.17
1944	880,118	0.06	0.14	0.20
1945	681,825	0.05	0.17	0.22
1946	670,364	0.09	0.20	0.29
1947	600,034	0.06	0.20	0.26
1950	258,789	0.04	0.17	0.21
1951	183,717	0.05	0.18	0.23

<sup>1</sup>Spencer (1942): data from 1934 to 1942.

<sup>2</sup>Spencer (1947): data from 1943 to 1947.

<sup>3</sup>Spencer (1951): data from 1950 and 1951.

<sup>4</sup>Total losses = dead + nonambulatory pigs.

**Table 4. Summary of total preslaughter deaths (transport and lairage) by country**

Country	Year	Total deaths, %	Reference
Belgium	1993	0.30	Christensen et al. (1994)
Canada	2004	0.12	Haley et al. (2008b)
	2004	0.10	Haley et al. (2008b)
	2003	0.07	Sunstrum et al. (2006)
	2003	0.09	Benjamin (2005)
	2001	0.17	Haley et al. (2008a)
	1996	0.14	Murray (2000)
Czech Republic	1997–2004	0.11	Vecerek et al. (2006)
Denmark	1998–2002	0.02	Barton Gade et al. (2007)
	1996	0.02	Barton Gade (1997)
	1993	0.03	Christensen et al. (1994)
	1984	0.12	Barton Gade et al. (2007)
	1978	0.12	Barton Gade et al. (2007)
France	1997	0.07	Colleu and Chevillon (1999)
	1995	0.08	Colleu and Chevillon (1999)
Germany	2003	0.10	Werner et al. (2007)
	2002	0.10	Werner et al. (2007)
	2001	0.13	Werner et al. (2007)
	2000	0.15	Werner et al. (2007)
	1999	0.17	Werner et al. (2007)
	1993	0.50	Christensen et al. (1994)
Italy	1993	0.10	Christensen et al. (1994)
The Netherlands	1993	0.16	Christensen et al. (1994)
	—	0.15	Lambooy and Engel (1991)
	1980	0.21	van Logtestijn et al. (1982)
	1976	0.30	Corstiaensen et al. (1977)
	1976	0.38	van Logtestijn et al. (1982)
	1972	0.52	van Logtestijn et al. (1982)
	1968	0.47	van Logtestijn et al. (1982)
	1964	0.28	van Logtestijn et al. (1982)
	1960	0.15	van Logtestijn et al. (1982)
Portugal	1993	0.16	Christensen et al. (1994)
Spain	—	0.33	Gosálvez et al. (2006)
	—	0.15	Palacio et al. (1996)
	1992–1994	0.22	Guàrdia et al. (1996)
Sweden	1976–1977	0.13	Fabiansson et al. (1979)
	1969	0.13	Fabiansson et al. (1979)
United Kingdom	1994	0.05	Riches et al. (1996)
	1993–1994	0.07	Riches et al. (1996)
	1993	0.09	Christensen et al. (1994)
	1990–1992	0.11	Abbott et al. (1995)
	1991–1992	0.07	Warriss and Brown (1994)
	1985	0.27	Robertson (1987)

*Continued*

### *Economic Implications*

**Swine Producers.** Economic losses experienced by pork producers include a complete loss of value on dead pigs and substantial price discounts on nonambulatory pigs. Discounts for nonambulatory pigs vary by region and by processing plant in the United States, and these discounts have been estimated to be as high as 30% of the total value of the pig (M. Ritter, Elanco Animal Health, Greenfield, IN, personal communication). There also may be economic losses associated with carcass trim loss on nonambulatory pigs because of carcass bruising.

Economic losses experienced by US swine producers in 2006 caused by transport losses in market weight pigs are summarized in Tables 6 and 7. The average US slaughter weight in 2006 was 122.15 kg. The estimated farrow-to-finish cost of production, average pig value, and net profit on a per-pig basis were \$105.03, \$125.50, and \$20.47, respectively (Table 6). The percentages of dead and nonambulatory pigs used in this economic analysis were 0.22% (Figure 1) and 0.44% (Table 5), respectively.

Economic losses were divided into “direct financial losses” and “indirect financial losses” (Table 7). Direct financial losses were defined as the price paid for dead and nonambulatory pigs minus the average farrow-to-finish cost of production. For the purposes of this analysis, it was assumed that dead pigs had zero value, whereas nonambulatory pigs were discounted by 30% (\$37.65) of the total value of the pig (\$125.50; Table 6). Thus, direct financial losses for dead pigs and nonambulatory pigs were calculated to be \$105.03 and \$17.18 per pig, respectively (Table 7).

Indirect financial losses refer to lost profit opportunities, which averaged \$20.47 per pig in 2006 (Table 6). When applied to the 103,688,100 pigs that were slaughtered in the United States in 2006, direct financial losses and indirect financial losses were determined to be \$31,796,810 and \$14,008,481, respectively (Table 7). Therefore, total economic losses asso-

**Table 4 (continued). Summary of total preslaughter deaths (transport and lairage) by country**

Country	Year	Total deaths, %	Reference
	1977–1978	0.07	Sains (1980)
	1970–1972	0.07	Allen et al. (1974)
	1969–1972	0.08	Smith and Allen (1976)
	1961–1973	0.10	Allen (1979)

food programs (e.g., school lunch programs and military commissaries), thereby requiring additional time and labor to segregate carcasses of nonambulatory pigs from those of normal pigs to ensure that meat from these carcasses is not used in these programs (M. Siemens, 2009, Cargill Meat Solutions, Wichita, KS, personal communication).

**Further Processors and Retailers.**


Economic losses associated with nonambulatory pigs may also be incurred by further processors and retailers. Carr et al. (2005) evaluated fresh pork quality traits of 246 fatigued pigs, and reported that the vast majority of fatigued pigs produced pork with high ultimate pH (6.0), low Minolta L\* values (45.72), and low drip losses (1.91%). These quality attributes are indicative of dark, firm, and dry pork; however, a small percentage of the fatigued pigs yielded pale, soft, and exudative pork. It is well-documented that short-term stress immediately before slaughter increases the rate of postmortem metabolism, resulting in pale, soft, and exudative pork, whereas long-term stress before slaughter depletes muscle glycogen stores, resulting in dark, firm, and

ciated with dead and nonambulatory pigs were estimated to be approximately \$46 million in 2006 (Table 7). This value is similar to the estimate of Ellis et al. (2003), who reported that transport losses cost US swine producers between \$50 million and \$100 million dollars annually.

**Pork Processors.** Pork processing plants have disposal costs for dead pigs, increased labor costs to move nonambulatory pigs quickly and humanely to the USDA suspect pen, and increased attention to regulations and personnel training on how to handle nonambulatory pigs. The USDA evaluates how nonambulatory pigs are handled at each plant and inspects all nonambulatory pigs for health before slaughter. Improper handling of non-

ambulatory pigs at a processing plant can result in a USDA noncompliance report. Furthermore, inhumane acts that are considered by the USDA inspector to be of an “egregious nature” will result in suspension of inspection at the processing plant (Food Safety and Inspection Service, 2003, 2007a, 2008a,b).

Two of the 5 “willful acts of abuse” monitored in the American Meat Institute’s animal handling audit of processing plants pertain to the handling of nonambulatory pigs. These willful acts of abuse include dragging conscious, nonambulatory pigs and driving normal pigs over the top of a nonambulatory pig (Grandin, 2007). Furthermore, meat from nonambulatory animals cannot enter government



**Figure 1.** Yearly incidence of dead market swine at USDA-inspected plants for the calendar years 1991 to 2006 (Food Safety and Inspection Service, 2007b).



**Table 5. A 23-trial summary on the incidence of dead and nonambulatory (NA) pigs in the United States<sup>1</sup>**

Reference	Year	Truckloads, no.	Pigs, no.	Farm NA, <sup>2</sup> %	Losses at the processing plant				
					Dead, %	Fatigued, <sup>3</sup> %	Injured, %	Total NA, <sup>4</sup> %	Total losses, <sup>5</sup> %
Berry et al. (2007)									
Study 1	2006– 2007	186	31,264	—	0.51	0.72	0.05	0.77	1.27
Study 2	2006– 2007	270	38,771	—	0.37	0.74	0.04	0.78	1.15
Ellis et al. (2003)									
Study 1	2000– 2001	2,199	367,453	—	0.28	—	—	0.81	1.09
Study 2	2001– 2002	93	15,736	—	0.19	—	—	0.62	0.81
Study 3	2000	24	4,052	—	0.25	—	—	0.75	1.00
Study 4	2000	23	3,761	—	0.70	—	—	1.55	2.25
Fitzgerald et al. (2009)	2005– 2006	12,333	2,053,945	—	0.25	0.55	0.05	0.60	0.85
Johnson et al. (2008)	2007	33	5,091	0.05	0.12	0.34	0.14	0.48	0.60
Lewis (2006)									
Study 1	2004	31	5,006	—	0.08	1.98	0.36	2.34	2.39
Study 2	2004	41	5,262	—	0.12	1.45	0.04	1.49	1.61
Murphy (2007)	2006– 2007	24	2,634	0.12	0.46	0.21	0.09	0.29	0.75
Rademacher and Davies (2005)	2002– 2005	7,396	1,303,148	—	0.37	—	—	0.40	0.77
Ritter (2007)									
Study 1	2003– 2004	74	12,511	0.26	0.23	0.55	0.24	0.85	1.08
Study 2	2004– 2005	42	6,953	0.20	0.52	0.79	0.16	0.95	1.47
Study 3	2005	35	3,827	0.00	0.05	0.16	0.08	0.24	0.29
Study 4	2005	37	4,027	0.00	0.08	0.10	0.12	0.22	0.30
Study 5	2005– 2006	109	17,256	0.11	0.35	0.42	0.08	0.51	0.90
Ritter (unpublished data)									
Study 1	2007	39	2,839	0.00	0.77	0.39	0.11	0.51	1.28
Study 2	2007	22	3,740	0.47	0.13	0.40	0.13	0.53	0.67
Stewart et al. (2008)									
Study 1	2007	32	5,884	0.09	0.04	0.05	0.05	0.11	0.14
Study 2	2007	14	2,591	0.04	0.00	0.12	0.04	0.16	0.16
Sutherland et al. (2006)	2004– 2005	16,323	2,730,754	—	0.18	0.21	0.05	0.26	0.44
Swan et al. (2007)	2003– 2004	192	34,064	0.06	0.26	0.65	0.34	0.99	1.25
Totals and weighted averages <sup>6</sup>		39,572	6,660,569	0.11	0.25	0.37	0.05	0.44	0.69

<sup>1</sup>All of the studies except Berry et al. (2007), Stewart et al. (2008), and Fitzgerald et al. (2009) stopped monitoring dead and nonambulatory pigs after pigs crossed the weigh scale. Rademacher and Davies (2005) reported dead pigs during unloading and after the pigs crossed the weigh scale.

<sup>2</sup>Farm NA = all nonambulatory pigs at the farm during loading.

<sup>3</sup>Fatigued = all nonambulatory, noninjured pigs at the processing plant.

<sup>4</sup>Total NA = all nonambulatory (fatigued and injured) pigs at the processing plant.

<sup>5</sup>Total losses = all dead and nonambulatory pigs at the processing plant.

<sup>6</sup>Weighted averages were adjusted for the total number of pigs in each study.

**Table 6. Assumptions used for calculating the economic impact of transport losses in market weight pigs on the US pork industry in 2006**

2006 economic assumption	Amount
Market hog statistic	
Number of pigs slaughtered <sup>1</sup>	103,688,100
Average slaughter wt, <sup>1</sup> kg	122.15
Market hog production cost	
Average farrow-to-finish break-even cost, <sup>2</sup> \$/kg	\$0.86
Average farrow-to-finish cost of production, <sup>3</sup> \$/pig	\$105.03
Market hog price	
Average live price paid, <sup>4</sup> \$/kg	\$1.03
Average pig value, <sup>5</sup> \$/pig	\$125.50
Market hog profit	
Average net profit, <sup>6</sup> \$/pig	\$20.47
Price paid for dead and nonambulatory pigs	
Dead pigs, <sup>7</sup> \$/pig	\$0.00
Nonambulatory pigs, <sup>8</sup> \$/pig	\$87.85

<sup>1</sup>Value obtained from USDA-National Agricultural Statistics Service (2007b).  
<sup>2</sup>Value obtained from Meyer (2008).  
<sup>3</sup>Average farrow-to-finish cost of production = average farrow-to-finish break-even cost × average slaughter weight.  
<sup>4</sup>Value obtained from USDA-National Agricultural Statistics Service (2007a).  
<sup>5</sup>Average pig value = average live price paid × average slaughter weight.  
<sup>6</sup>Average net profit = average pig value – average farrow-to-finish cost of production.  
<sup>7</sup>Assumes complete loss of value on dead pigs.  
<sup>8</sup>Assumes nonambulatory pigs are discounted 30%.

*Symptoms and Metabolic Characteristics of Fatigued Pigs*

**Symptoms.** Anderson et al. (2002) reported that fatigued pigs at processing plants displayed the acute-stress symptoms of open-mouth breathing (44%), skin discoloration (77%), muscle tremors (83%), and abnormal vocalizations (30%). Additionally, a strong positive relationship ( $r = 0.81$ ) between dead and nonambulatory pigs has been reported (Hamilton et al., 2003), which suggests a common link or cause for dead and nonambulatory pigs. On this basis, a continuum of stress was recently proposed for the development of fatigued and dead pigs (Ritter et al., 2005). As the pig begins to experience stress, it will display open-mouth breathing, skin discoloration, or both. If the stress is not removed or if additional stressors are introduced, the pig will become reluctant to move, make abnormal vocalizations, develop muscle tremors, or some combination of these signs. At this stage, the pig may become overwhelmed by the accumulation of stress, in which case the pig will collapse and become nonambulatory, and, in extreme cases, death may ensue.

dry pork (reviewed by Gregory, 1994). On this basis, Carr et al. (2005) hypothesized that the fresh

quality carcass traits of fatigued pigs depend on when the pig becomes fatigued during the marketing process.

**Table 7. Economic impact of transport losses in market weight pigs on the US pork industry in 2006<sup>1</sup>**

Variable	Transport losses		Direct financial losses		Indirect financial losses	
	Average losses, <sup>2,3</sup> %	Total losses, <sup>4</sup> no.	Average losses, <sup>5</sup> \$/pig	Total losses, <sup>6</sup> \$	Average losses, <sup>7</sup> \$/pig	Total losses, <sup>8</sup> \$
Dead pigs	0.22	228,114	(\$105.03)	(\$23,958,813)	(\$20.47)	(\$4,669,494)
Nonambulatory pigs	0.44	456,228	(\$17.18)	(\$7,837,997)	(\$20.47)	(\$9,338,987)
Total		684,341		(\$31,796,810)		(\$14,008,481)

<sup>1</sup>Values are based on the economic assumptions described in Table 6.  
<sup>2</sup>The value for percentage of dead pigs in 2006 was obtained from Figure 1.  
<sup>3</sup>The value for percentage of nonambulatory pigs in 2006 was obtained from Table 5.  
<sup>4</sup>Calculated by multiplying the percentage of dead or nonambulatory pigs by the total number of pigs slaughtered in 2006.  
<sup>5</sup>Average direct financial losses = (price paid for dead or nonambulatory pigs) – (average farrow-to-finish cost of production).  
<sup>6</sup>Total direct financial losses = (average direct financial losses) × (number of dead or nonambulatory pigs).  
<sup>7</sup>Average in-direct financial losses = average net profit per pig (Table 6).  
<sup>8</sup>Total in-direct financial losses = (average net profit per pig) × (number of dead or nonambulatory pigs).

**Metabolic Characteristics.** Ivers et al. (2002) measured several metabolic parameters in normal and fatigued pigs during unloading at the processing plant. Compared with normal pigs from the same trailer load, fatigued pigs had higher blood lactate, ammonia, sodium, potassium, cortisol, epinephrine, and norepinephrine concentrations while having lower blood pH, bicarbonate, base excess, calcium, partial pressure of carbon dioxide, and insulin values. Additionally, fatigued pigs had lower liver glycogen concentrations and lower glycolytic potential values in the LM and semitendinosus muscles than normal pigs. Thus, the researchers concluded that fatigued pigs are in a metabolic state of acidosis characterized by high blood lactate and low blood pH values (Anderson et al., 2002; Ivers et al., 2002).

Additionally, it is plausible that pigs could develop fatigue from chronic stress. Chronic stress depletes muscle glycogen stores and may result in physical exhaustion and fatigue (reviewed by Gregory, 1994, 1996). In support of this theory, Carr et al. (2005) reported that the majority of fatigued pigs evaluated had high LM ultimate pH (5.90 to 7.00), suggesting that muscle glycogen stores were substantially reduced before slaughter. However, the relationship between muscle glycogen stores and the incidence of fatigued pigs is currently unknown and warrants additional research.

#### ***Do Fatigued Pigs Recover?***

Blood pH, lactate, bicarbonate, and base-excess values of aggressively handled pigs return to baseline resting values by 2 h posthandling (Anderson et al., 2002; Bertol et al., 2002). Furthermore, Ritter et al. (2006) monitored 25 pigs that had become fatigued on the trailer during loading, and reported that 72% of these pigs were normal during unloading at the processing plant after a 3-h journey. Results of these studies suggest that fatigued pigs may recover if allowed to rest for at least 2 to 3 h.

## IMPLICATIONS

Transport losses on average total 0.7%, meaning that more than 99% of pigs that are transported walk off the truck, walk through the plant, and are processed without delay. Despite the large percentage of unaffected pigs, transport losses have been estimated to cost the US swine industry approximately \$46 million annually. More than 80% of pigs that become nonambulatory are in a state of metabolic acidosis and are classified as fatigued, yet the majority of these pigs will recover fully if given time to rest. Because transport losses are a multifactorial problem consisting of pig, facility design, people, management, transportation, processing plant, and environmental factors, continued research efforts are needed to understand how each of these factors and the relationships among factors affect the well-being of the pig during the marketing process.

## ACKNOWLEDGMENTS

The authors acknowledge the National Pork Board Animal Welfare Committee for organizing and sponsoring this project.

## LITERATURE CITED

Abbott, T. A., H. J. Guise, E. J. Hunter, R. H. C. Penny, P. J. Barnes, and C. Easby. 1995. Factors influencing pig deaths during transit: An analysis of drivers' reports. *Anim. Welfare* 4:29.

Allen, W. M. 1979. Losses of pigs due to the "acute stress syndrome" in the United Kingdom. *Acta Agric. Scand.* 21:495.

Allen, W. M., C. N. Hebert, and L. P. Smith. 1974. Deaths during and after transportation in pigs in Great Britain. *Vet. Rec.* 94:212.

Anderson, D. B., D. J. Ivers, M. E. Benjamin, H. W. Gonyou, D. J. Jones, K. D. Miller, R. K. McGuffey, T. A. Armstrong, D. H. Mowrey, L. F. Richardson, R. Seneriz, J. R. Wagner, L. E. Watkins, and A. G. Zimmermann. 2002. Physiological responses of market hogs to different handling practices. p. 399 in Proc. Am. Assoc. Swine Vet., Kansas City, MO.

Averós, X., T. G. Knowles, S. N. Brown, P. D. Warriss, and L. F. Gosálvez. 2008. Factors affecting the mortality of pigs being transported to slaughter. *Vet. Rec.* 163:386.

Barton Gade, P. 1997. The effect of pre-slaughter handling on meat quality in pigs. p. 100 in *Manipulating Pig Production*. P. D. Cranwell, ed. Australas. Pig Sci. Assoc., Victoria, Australia.

Barton Gade, P., L. Christensen, M. Baltzer, and J. Valentin Petersen. 2007. Causes of pre-slaughter mortality in Danish slaughter pigs with special emphasis on transport. *Anim. Welfare* 16:459.

Benjamin, M. E. 2005. Pig handling: Physiological responses and meat quality discussion. In Proc. 2005 Manitoba Swine Seminar. <http://www.gov.mb.ca/agriculture/livestock/pork/pdf/bab19s01.pdf> Accessed May 14, 2009.

Berry, N., A. Johnson, J. Hill, T. Baas, L. Karriker, and K. Stalder. 2007. Loading gantry versus traditional chute for the finisher pig: Effect on welfare parameters at time of marketing. p. 137 in Proc. Allen D. Leman Swine Conf., St. Paul, MN.

Bertol, T. M., M. Ellis, D. N. Hamilton, and F. McKeith. 2002. Effect of handling intensity on blood acid-base balance in slaughter pigs. *J. Anim. Sci.* 80(Suppl. 2):86. (Abstr.)

Carr, S. N., J. P. Gooding, P. J. Rincker, D. N. Hamilton, M. Ellis, J. Killefer, and F. K. McKeith. 2005. A survey of pork quality of downer pigs. *J. Muscle Foods* 16:298.

Christensen, L., P. Barton Gade, and L. Blaabjerg. 1994. Investigation of transport conditions in participating countries in the EC project: PL92063. p. W-201 in Proc. 40th Int. Congr. Meat Sci. Technol., The Hague, the Netherlands.

Colleu, T., and P. Chevillon. 1999. Effect of climatic parameters and distance on pig mortality during transport. *Techni-Porc.* 22:31.

Corstiaensen, G. P., J. J. M. De Bruin, L. H. H. M. Lendfers, J. G. van Logtestijn, and A. Th. Verdijk. 1977. Pig losses during transport and in the pens of slaughterhouses in the Netherlands in 1976. *Tijdschr. Diergeneeskd.* 102:811.

Ellis, M., F. McKeith, D. Hamilton, T. Bertol, and M. Ritter. 2003. Analysis of the current situation: What do downers cost the industry and what can we do about it? p. 1 in Proc. 4th Am. Meat Sci. Assoc. Pork Quality Symp., Columbia, MO.

Ellis, M., and M. Ritter. 2005a. Impact of animal handling and transportation conditions on losses of slaughter weight swine during transport. p. 199 in Proc. Am. Assoc. Swine Vet., Toronto, Ontario, Canada.

Ellis, M., and M. Ritter. 2005b. Transport losses: Causes and solutions. p. 176 in Proc. Allen D. Leman Swine Conf., St. Paul, MN.

European Commission. 1995. Council Directive 95/29/EC, On the protection of animals during transport and amending Directive 91/628/EEC. *Off. J. Eur. Communities L* 148:52.

- European Economic Community. 1991. Council Directive 91/628/EEC, On the protection of animals during transport and amending Directives 90/425/EEC and 91/496/EEC. *Off. J. Eur. Communities L* 340:17.
- Fabiansson, S., K. Lundström, and I. Hansson. 1979. Mortality among pigs during transport and waiting time before slaughter in Sweden. *Swed. J. Agric. Res.* 9:25.
- Fitzgerald, R. F., K. J. Stalder, J. O. Matthews, C. M. Schultz Kaster, and A. K. Johnson. 2009. Factors associated with fatigued, injured, and dead pig frequency during transport and lairage at a commercial abattoir. *J. Anim. Sci.* 87:1156.
- Food Safety and Inspection Service. 2003. FSIS Directive 6900.2 Revision 1: Humane handling and slaughter of livestock. <http://www.fsis.usda.gov/OPPDE/RDAD/FSISDirectives/6900.2Rev1.pdf> Accessed Nov. 15, 2008.
- Food Safety and Inspection Service. 2007a. FSIS Directive 6100.1: Ante-mortem livestock inspection. Food Safety Inspection Service, Washington, DC.
- Food Safety and Inspection Service. 2007b. Market swine condemned ante-mortem for deads in USDA inspected plants for the calendar years of 1991 to 2006. FOIA Case No. 07-148. Food Safety Inspection Service, Washington, DC.
- Food Safety and Inspection Service. 2008a. Humane slaughter of livestock. Code of Federal Regulations, Part 313. Food Safety Inspection Service, Washington, DC.
- Food Safety and Inspection Service. 2008b. Humane handling activities and documentation in livestock slaughter establishments. FSIS Notice 16-08. Food Safety Inspection Service, Washington, DC.
- Gosálvez, L. F., X. Averós, J. J. Valdelvira, and A. Herranz. 2006. Influence of season, distance and mixed loads on the physical and carcass integrity of pigs transported to slaughter. *Meat Sci.* 73:553.
- Grandin, T. 2007. Recommended Animal Handling Guidelines and Audit Guide. 2007 ed. Am. Meat Inst. Found., Washington, DC. <http://www.meatami.com/ht/a/GetDocumentAction/i/1489> Accessed Dec. 4, 2008.
- Gregory, N. G. 1994. Preslaughter handling, stunning and slaughter. *Meat Sci.* 36:45.
- Gregory, N. G. 1996. Welfare and hygiene during preslaughter handling. *Meat Sci.* 43:S35.
- Guàrdia, M., M. Gispert, and A. Diestre. 1996. Mortality rates during transport and lairage in pigs for slaughter. *Meat Focus Int.* 10:362.
- Haley, C., C. E. Dewey, T. Widowski, and R. Friendship. 2008a. Association between in-transit loss, internal trailer temperature, and distance traveled by Ontario market hogs. *Can. J. Vet. Res.* 72:385.
- Haley, C., C. E. Dewey, T. Widowski, Z. Poljak, and R. Friendship. 2008b. Factors associated with in-transit losses of market hogs in Ontario in 2001. *Can. J. Vet. Res.* 72:377.
- Hamilton, D. N., M. Ellis, T. M. Bertol, and K. D. Miller. 2004. Effects of handling intensity and live weight on blood acid-base status in finishing pigs. *J. Anim. Sci.* 82:2405.
- Hamilton, D. N., M. Ellis, G. E. Bressner, B. F. Wolter, D. J. Jones, and L. E. Watkins. 2003. Relationships between environmental conditions on trucks and losses during transport to slaughter in finishing pigs. *J. Anim. Sci.* 81(Suppl. 2):53. (Abstr.)
- Ivers, D. J., L. F. Richardson, D. J. Jones, L. E. Watkins, K. D. Miller, J. R. Wagner, R. Seneriz, A. G. Zimmermann, K. A. Bowers, and D. B. Anderson. 2002. Physiological comparison of downer and non-downer pigs following transportation and unloading at a processing plant. *J. Anim. Sci.* 80(Suppl. 2):39. (Abstr.)
- Johnson, A., L. Sadler, M. Faga, C. Feuerbach, H. Hill, R. Bailey, and M. Ritter. 2008. Effects of facility design on the welfare of market weight pigs during loading and unloading. *J. Anim. Sci.* 86(E Suppl. 2):297. (Abstr.)
- Lambooy, E., and B. Engel. 1991. Transport of slaughter pigs by truck over a long distance: some aspects of loading density and ventilation. *Livest. Prod. Sci.* 28:163.
- Lewis, C. R. 2006. Transportation and handling of swine in commercial operations. MS Thesis. Texas Tech University, Lubbock.
- McGlone, J. J., R. L. McPherson, and D. L. Anderson. 2004. Case study: Moving devices for finishing pigs—Efficacy of electric prod, board, paddle, or flag. *Prof. Anim. Sci.* 20:518.
- Meyer, S. 2008. Trends in U.S. pork production. In Proc. 2008 Banff Pork Seminar. <http://www.banffpork.ca/breakout/breakout2008.shtml#8> Accessed March 5, 2008.
- Murphy, C. M. 2007. The effect of floor space and external environmental conditions on the internal environment of a swine livestock trailer and transport losses. MS Thesis. University of Illinois, Urbana-Champaign.
- Murray, A. C. 2000. Reducing losses from gate to packer. *Adv. Pork Prod.* 11:175.
- National Pork Board. 2007. National Pork Board request for proposals: Information for spring 2007 funding. <http://www.pork.org/Documents/RFP/2007SpringCall.doc> Accessed Nov. 15, 2008.
- National Pork Board. 2008a. Animal welfare—Research. <http://www.pork.org/PorkScience/Research/Research.aspx?c=4> Accessed Dec. 2, 2008.
- National Pork Board. 2008b. Transport Quality Assurance. <http://www.pork.org/Producers/TQA/TQA.aspx> Accessed Nov. 15, 2008.
- Palacio, J., S. Garcia-Belenguer, F. M. Gascon, F. Liste, C. Orega, B. Lobera, I. Martin-Maestro, J. A. Angel, J. C. Lles, and F. Bayo. 1996. Mortality in slaughter pigs during transport. *Invest. Agric., Prod. Sanid. Anim.* 11:159.
- Rademacher, C., and P. Davies. 2005. Factors associated with the incidence of mortality during transport of market hogs. p. 186 in Proc. Allen D. Leman Swine Conf., St. Paul, MN.
- Riches, H. L., H. J. Guise, R. H. C. Penny, T. A. Jones, and A. Cuthbertson. 1996. A national survey of transport conditions for pigs. *Pig J.* 38:8.
- Ritter, M., M. Ellis, M. Benjamin, E. Berg, P. DuBois, J. Marchant-Forde, A. Green, P. Matzat, P. Mormede, T. Moyer, K. Pfalzgraf, M. Siemens, J. Sterle, T. Whiting, B. Wolter, and A. Johnson. 2005. The fatigued pig syndrome. *J. Anim. Sci.* 83(Suppl. 1):258. (Abstr.)
- Ritter, M. J. 2007. Effects of animal handling and transportation factors on the welfare, stress responses, and incidences of transport losses in market weight pigs at the packing plant. PhD Diss. University of Illinois, Urbana-Champaign.
- Ritter, M. J., M. Ellis, J. Brinkmann, J. M. DeDecker, M. E. Kocher, K. K. Keffaber, B. A. Peterson, J. M. Schlipf, and B. F. Wolter. 2006. Effect of floor space during transport of market weight pigs on incidence of transport losses (dead and non-ambulatory pigs) at the packing plant and relationships between transport conditions and losses. *J. Anim. Sci.* 84:2856.
- Robertson, J. F. 1987. Bacon pigs: A practical approach to the reduction of deaths in transit. New perspectives in pig production. p. 23 in Proc. North of Scotland College of Agriculture Pig Conference, North of Scotland College of Agriculture, Aberdeen, UK.
- Rosse, J. C. 1972. Trend in hog transit deaths. Livestock Conservation Inc., Omaha, NE.
- Sains, A. 1980. Deaths in transit: What British surveys show. *Pig Farming* 28:40.
- Smith, L. P., and W. M. Allen. 1976. A study of the weather conditions related to the death of pigs during and after their transportation in England. *Agric. Meteorol.* 16:115.
- Smith, W. W. 1937. Marketing: Market classes and grades. p. 450 in *Pork Production*. The MacMillan Company, New York, NY.
- Spencer, W. T. 1942. 1942 annual report to the live stock sanitary and loss prevention board of the Omaha livestock market interests on the eradication and control of live stock diseases and prevention of losses inci-



- dent to marketing live stock. The Live Stock and Loss Prevention Board, Omaha, NE.
- Spencer, W. T. 1947. 26th annual report to the live stock sanitary and loss prevention board of the Omaha livestock market interests on the eradication and control of live stock diseases and prevention of losses incident to marketing. The Live Stock and Loss Prevention Board, Omaha, NE.
- Spencer, W. T. 1951. 30th annual report to the live stock sanitary and loss prevention board of the Omaha livestock market interests on the eradication and control of live stock diseases and prevention of losses incident to marketing. The Live Stock and Loss Prevention Board, Omaha, NE.
- Stewart, G., M. Ritter, M. Culbertson, G. Mann, and R. Wofford. 2008. Effects of previous handling and feed withdrawal prior to loading on transport losses in market weight pigs. p. 359 in Proc. Am. Assoc. Swine Vet. Mtg., San Diego, CA.
- Sunstrum, J., C. Dewey, and C. Haley. 2006. Clinical signs of stress in finisher pigs transported to market in the summer. p. 45 in Proc. Am. Assoc. Swine Vet., Orlando, FL.
- Sutherland, M. A., A. McDonald, and J. J. McGlone. 2006. Factors influencing the percentage of dead and fatigued pigs during transport. Page 609 in Proc. 19th Int. Pig Vet. Soc. Congr., Copenhagen, Denmark.
- Swan, J. E., M. H. Gillis, K. D. Miller, J. D. Muegge, D. H. Mowrey, T. A. Armstrong, W. C. Weldon, and M. J. Ritter. 2007. Effects of ractopamine on transport losses in market weight pigs. *J. Anim. Sci.* 85(Suppl. 1):127. (Abstr.)
- Topel D. G. Bicknell E. J. Preston K. S. Christian L. L. Matsushima C. Y.. 1968. Porcine stress syndrome. *Mod. Vet. Pract.* 49:40, 59.
- Topel, D. G., and L. L. Christian. 1981. Porcine stress syndrome. p. 647 in *Diseases of Swine*. 5th ed. A. D. Leman, R. D. Glock, W. L. Mengeling, R. H. C. Penny, E. Scholl, and B. Straw, ed. The Iowa State University Press, Ames.
- USDA-National Agricultural Statistics Service. 2007a. Agricultural prices—2006 summary. <http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1003> Accessed Nov. 2, 2007.
- USDA-National Agricultural Statistics Service. 2007b. Quick stats—US and state slaughter statistics on barrows and gilts for the year of 2006. <http://www.nass.usda.gov/index.asp> Accessed Nov. 1, 2007.
- USDA-National Agricultural Statistics Service. 2008. Quick stats—US and state slaughter statistics on barrows and gilts for the years of 1993 to 1999. <http://www.nass.usda.gov/QuickStats/index.asp> Accessed March 5, 2008.
- US House of Representatives. 2007. H. R. 661: Downed Animal and Food Safety Protection Act. <http://thomas.loc.gov/cgi-bin/query/z?c110:h661> Accessed May 20, 2009.
- US Senate. 2007. S. 394: Downed Animal and Food Safety Protection Act. <http://thomas.loc.gov/cgi-bin/query/z?c100:s394> Accessed May 20, 2009.
- van Logtestijn, J. G., A. M. T. C. Romme, and G. Eikelenboom. 1982. Losses caused by transport of slaughter pigs in the Netherlands. p. 105 in *Transport of Animals Intended for Breeding, Production and Slaughter* Vol. 18 of *Current Topics in Veterinary Medicine and Animal Science*. R. Moss, ed. Martinus Nijhoff Publishers, London, UK.
- Vecerek, V., M. Malena, M. Malean Jr., E. Voslarova, and P. Chloupek. 2006. The impact of the transport distance and season on losses of fattened pigs during transport to the slaughterhouse in the Czech Republic in the period from 1997 to 2004. *Vet. Med.-Czech.* 51:21.
- Warriss, P. D., and S. N. Brown. 1994. A survey of mortality in slaughter pigs during transport and lairage. *Vet. Rec.* 134:513.
- Werner, C., K. Reiners, and M. Wicke. 2007. Short as well as long transport duration can affect the welfare of slaughter pigs. *Anim. Welfare* 16:385.