

3-2001

## Seed physiological performance of soybeans with altered saturated fatty acid contents

Tong Wang

*Iowa State University*, [tongwang@iastate.edu](mailto:tongwang@iastate.edu)

Teresa Harp

*Iowa State University*

Earl G. Hammond

*Iowa State University*


Joseph S. Burris

*Iowa State University*

Walter R. Fehr

*Iowa State University*

Follow this and additional works at: [http://lib.dr.iastate.edu/fshn\\_ag\\_pubs](http://lib.dr.iastate.edu/fshn_ag_pubs)

 Part of the [Food Science Commons](#), and the [Human and Clinical Nutrition Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/fshn\\_ag\\_pubs/1](http://lib.dr.iastate.edu/fshn_ag_pubs/1). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Food Science and Human Nutrition at Iowa State University Digital Repository. It has been accepted for inclusion in Food Science and Human Nutrition Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Seed physiological performance of soybeans with altered saturated fatty acid contents

## **Abstract**

Soybean (*Glycine max* (L.) Merr.) seeds with elevated or reduced percentages of palmitate and elevated percentages of stearate were compared with seeds of typical composition in tests for germination, seedling growth rate and leachate conductivity. In general, seeds with altered compositions did well in these physiological tests, but their vigour tended to be negatively correlated with the percentages of stearate and palmitate in various lipid classes.

## **Keywords**

germination, *Glycine max* (L.) Merr., membrane permeability, neutral lipids, phospholipids, saturated fatty acids, seedling growth rate

## **Disciplines**

Food Science | Human and Clinical Nutrition

## **Comments**

This article is from *Seed Science Research*, 11 (2001): 93–97, doi: [10.1079/SSR200063](https://doi.org/10.1079/SSR200063).

## Seed Science Research

<http://journals.cambridge.org/SSR>


Additional services for **Seed Science Research**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



---

## Seed physiological performance of soybeans with altered saturated fatty acid contents

Tong Wang, Teresa Harp, Earl G. Hammond, Joseph S. Burrisa and Walter R. Fehr

Seed Science Research / Volume 11 / Issue 01 / March 2011, pp 93 - 97  
DOI: 10.1079/SSR200063, Published online: 22 February 2011

**Link to this article:** [http://journals.cambridge.org/abstract\\_S0960258501000101](http://journals.cambridge.org/abstract_S0960258501000101)

### How to cite this article:

Tong Wang, Teresa Harp, Earl G. Hammond, Joseph S. Burrisa and Walter R. Fehr (2011). Seed physiological performance of soybeans with altered saturated fatty acid contents. Seed Science Research, 11, pp 93-97 doi:10.1079/SSR200063

**Request Permissions :** [Click here](#)

## SHORT COMMUNICATION

## Seed physiological performance of soybeans with altered saturated fatty acid contents

Tong Wang<sup>1\*</sup>, Teresa Harp<sup>1</sup>, Earl G. Hammond<sup>1</sup>, Joseph S. Burris<sup>2</sup> and Walter R. Fehr<sup>3</sup>

<sup>1</sup>Department of Food Science and Human Nutrition; <sup>2</sup>Seed Science Center; <sup>3</sup>Department of Agronomy, Iowa State University, Ames, IA 50011, USA

### Abstract

Soybean (*Glycine max* (L.) Merr.) seeds with elevated or reduced percentages of palmitate and elevated percentages of stearate were compared with seeds of typical composition in tests for germination, seedling growth rate and leachate conductivity. In general, seeds with altered compositions did well in these physiological tests, but their vigour tended to be negatively correlated with the percentages of stearate and palmitate in various lipid classes.

**Keywords:** germination, *Glycine max* (L.) Merr., membrane permeability, neutral lipids, phospholipids, saturated fatty acids, seedling growth rate

### Introduction

Modification of the fatty acid composition of soybean oil to make it more competitive in various segments of the food and industrial oil markets (Hammond, 1992) and to make it more nutritional (Hu *et al.*, 1997) has been an important objective of plant breeding and molecular genetics in recent years. Altered fatty acid composition has been developed through traditional plant breeding (Wilson *et al.*, 1981) and application of chemical mutagens (Hammond and Fehr, 1975; Wilcox *et al.*, 1984; Fehr *et al.*, 1991) that have extended the range of the five major fatty acids normally found in soybean oil. One alteration of interest has been elevation of the saturated fatty acids, palmitate and stearate (Hartmann *et al.*, 1996; Rebetzke *et al.*, 1996, 1998; Stoltzfus *et al.*, 2000). Elevated palmitate may be used to achieve a desirable crystal structure in plastic

fats, and soybeans with elevated saturated fatty acids may be used to make margarine and shortening stocks without hydrogenation, thereby reducing processing cost and avoiding the formation of *trans* fatty acids (Kok, 1998). Soybean oil with elevated palmitate also exhibits improved oxidative stability when used as frying oil (Shen *et al.*, 1997).

Among the lipid components of soybean seeds, 0.3–0.6% phospholipids (PL) are present, with phosphatidylcholine (PC), phosphatidylethanolamine (PE) and phosphatidylinositol (PI) as the major classes (Hui, 1996; Wang *et al.*, 1997). These PLs are structural components of the plasma membrane. Membrane PLs must have the correct balance of saturated and unsaturated fatty acids (Chapman, 1973) to be in the proper physical state for cells to perform their metabolic tasks. In soybean mutants whose triacylglycerol (TAG) saturate percentages are elevated, PL saturate percentages also are increased (Mounts *et al.*, 1996; Wang *et al.*, 1997). Soybean seeds with elevated stearate frequently have poor field germination and reduced yield (observed by W.R. Fehr, Agronomy Department, Iowa State University, data not published). When canolas were genetically modified to produce oils with greater than 30% stearate, their germination rates were lower, and seedlings were less vigorous. Accumulation of stearate in membrane lipids may have made membranes less adaptable to changes in temperature and moisture content (Thompson and Li, 1997).

The standard germination test is the most widely accepted procedure for estimating seed viability, but it is an inadequate predictor of field emergence (Dornbos, 1988). Tests of seedling vigour were developed to complement viability tests. The seedling growth rate (SGR) test gives a reproducible and objective evaluation of differences in growth rate. The conductivity test is a measurement of electrolyte leakage from plant tissues, and the leachate

---

\*Correspondence  
Fax: 515-294-8181  
Email: tongwang@iastate.edu

conductivity has been negatively correlated with seed viability and vigour (McKersie and Senaratna, 1983).

In this study, we report the effect of altered saturated fatty acid content in soybean seed lipids on germination, seedling growth rate, and seed electrolyte retention.

## Materials and methods

### *Soybean sample selection for physiological tests*

Commercial soybean cultivars and experimental soybean lines were provided by W.R. Fehr at Iowa State University and Pioneer Hi-Bred International Inc. of Des Moines, IA, USA. One hundred individual seeds from each soybean line with altered palmitate or stearate content were analysed by gas chromatography (GC) for fatty acid composition (Hammond, 1991), and 12 soybean lines were selected for the widest possible ranges of palmitate and stearate percentages and for the least variation in seed-to-seed composition.

### *Lipid compositional analysis*

PL isolation, class separation and fatty acid composition analysis were performed as described by Wang *et al.* (1997).

### *Germination evaluation*

The rolled-towel germination method (AOSA, 1986) was used. Each roll consisted of two standard-weight paper towels (35.5 × 63 cm). Each towel was moistened with 30 g of water. On the lower towel, 50 seeds were oriented into two rows with 25 seeds per row. The seeds were covered with an upper towel, the two paper towels were loosely rolled, and the rolls were placed upright in plastic containers that were loosely covered with plastic bags and placed in unlighted germinators at 15, 25 and 35°C. After 7 days, germination counts were made. Three replications for each line/temperature combination were performed.

### *Seedling growth rate*

The seedling growth rate (SGR) vigour test, which evaluates the growth of seedlings from a standard germination test, was performed according to the procedures described in the Rules for Testing Seeds (AOSA, 1986). After germination counts, the normal seedlings were freed of cotyledons, dried at 84°C for 63 h, and weighed. SGR was expressed as mg/seedling.

### *Conductivity test*

A G-2000 seed analyser (Wavefront Inc., Ann Arbor, MI, USA) was used to measure the electrical conductivities of the leachates for each of 50 individual seeds at ambient temperature. Each seed was placed in a cell filled with 3.5 ml distilled water on a 100-cell tray. Conductivity readings in micro-Siemens were taken after leaching for 1, 3, 6, 12 and 24 h. Three 50-seed replicates were tested for each line.

### *Statistical analysis*

A general linear model procedure of SAS was used for analysis of variance, and the least significant difference (LSD) was determined for mean comparison (SAS, 1984).

## Results and discussion

### *Fatty acid composition*

The average fatty acid composition of the soybean lines tested and the standard deviations among 100 individual seeds are shown in Table 1. Two commercial cultivars, Sturdy and Kenwood, were included in the study. For the soybean lines with elevated stearate, there was considerably greater variation among individual seeds. The unsaturated fatty acid percentages varied to compensate for the changes in the saturated fatty acid percentages.

The fatty acid composition of the major lipid classes in the experimental lines, except for 'Sturdy', is presented in Table 2. In lines with lipids elevated in saturates, there were corresponding increases in the PLs. PC and PE contained higher palmitate percentages than TAG except for two of the lines with elevated palmitate (PAL53 and PAL82). PE generally contained more palmitate but less stearate than PC and TAG except in the reduced palmitate lines, while PI contained the highest palmitate and stearate of the PL. PI palmitate was also higher than that of TAG in all instances, but PI stearate was frequently less than that of TAG in elevated saturate lines. When one of the saturates was elevated or reduced in PL, the other saturate tended to change in the opposite direction. These observations suggest that the saturate content of soybean PL is regulated in some way.

### *Seed physiological behaviour*

Data for germination percentages, SGR and leachate conductivity of the 12 soybean lines are presented in Table 3. Germination percentages at 15 and 25°C were generally high except for PAL79, which had reduced

**Table 1.** Average soybean fatty acid percentages (mole) and compositional variation (SD) among 100 seeds

Classification	Sample ID	16:0 <sup>a</sup>	SD	18:0	SD	18:1	SD	18:2	SD	18:3	SD
Typical composition	Sturdy	10.1	0.3	4.3	1.5	24.3	2.2	52.9	1.8	8.4	0.7
	STE07	10.4	0.6	4.3	0.4	31.9	4.5	46.8	3.6	6.6	0.6
	STE15	11.1	0.5	4.0	0.3	28.3	3.8	49.3	2.7	7.3	1.1
	Kenwood	11.6	0.5	4.4	0.3	22.6	2.7	53.3	2.2	8.1	0.7
Reduced 16:0	PAL79	3.0	0.9	2.9	0.3	16.1	2.5	65.1	2.0	12.9	1.9
	PAL22	3.3	0.5	4.1	0.5	26.2	4.0	56.8	3.6	9.7	1.3
Elevated 16:0	PAL53	26.5	0.9	4.8	0.6	15.3	1.2	44.5	1.0	8.9	0.8
	PAL82	32.5	1.9	5.3	0.8	12.9	1.6	37.3	1.5	12.1	1.2
Elevated 18:0	STE39	10.1	0.5	15.6	2.6	16.7	2.0	48.3	1.9	9.4	0.9
	STE62	9.4	0.6	21.8	4.0	17.5	1.7	43.4	3.7	8.0	0.8
	STE73	8.7	0.8	25.1	5.9	17.5	1.9	40.6	4.7	8.2	0.7
	STE71	9.6	0.5	26.6	3.6	16.9	1.0	39.2	3.5	7.8	0.7

<sup>a</sup>The number before the colon is the chain length of the fatty acid; the number after the colon is the number of double bonds in each fatty acid molecule.

**Table 2.** Saturated fatty acid percentages of neutral lipid (TAG) and phospholipids (PLs) for various lines of soybean seeds

Sample classification	Sample ID	TAG		PC		PE		PI	
		16:0	18:0	16:0	18:0	16:0	18:0	16:0	18:0
Typical composition	STE07	10.4	4.3	14.4	3.7	20.5	2.9	29.9	7.5
	STE15	11.1	4.0	15.0	3.7	20.1	2.8	32.3	6.9
	Kenwood	11.6	4.4	15.1	4.5	22.1	3.4	33.0	9.1
Reduced 16:0 <sup>a</sup>	PAL79	3.0	2.9	8.6	6.8	11.9	7.3	17.3	15.9
	PAL22	3.3	4.1	9.2	5.6	14.9	6.5	23.2	14.1
Elevated 16:0	PAL53	26.5	4.8	21.4	2.8	26.6	1.6	40.2	4.6
	PAL82	32.5	5.3	25.4	3.4	26.4	1.8	36.7	6.8
Elevated 18:0	STE39	10.1	15.6	11.7	8.6	18.2	5.9	24.8	15.2
	STE62	9.4	21.8	11.5	11.2	15.2	7.6	21.6	20.6
	STE73	8.7	25.1	11.2	11.9	16.0	8.3	22.2	19.3
	STE71	9.6	26.6	12.3	11.6	17.5	8.3	21.1	22.0

<sup>a</sup>See Table 1 for explanation of the fatty acid symbols.

saturate content and shrivelled seeds. Severe seed fungal decay occurred at 35°C, making evaluation of germination at this temperature less accurate. SGR was much greater at 25 and 35°C than at 15°C, and in most instances SGR was slightly better at 25 than 35°C.

The electrical conductivity of the 24-h leachate was generally lower for typical-composition lines (300–330  $\mu$ Siemens) than for those with altered compositions (360–600  $\mu$ Siemens), except for STE07, which had significantly higher conductivity, and STE39, which had a relatively low conductivity compared to the lines with similar fatty acid compositions. PAL79 had an unusually high conductivity (594  $\mu$ Siemens). The conductivity was also measured at 1, 3, 6 and 12 h. The values at 1 and 3 h were low (below 100  $\mu$ Siemens) and similar for all lines (data not shown). At 6 and 12 h, the relative

conductivity differences among lines were similar to those at 24 h.

Correlation coefficients obtained from linear correlation analysis between neutral and PL saturates and germination percentage, SGR, and conductivity are shown in Table 4. The reduced saturate line, PAL79, was not included in the analysis because of its shrivelled seeds and poor germination. Generally, germination percentages were negatively correlated with saturated fatty acid content for all classes of soybean lipids, although many of the correlation coefficients were not statistically significant at the 5% level. Elevated saturate lines had significantly lower germination at 25°C, especially for elevated stearate lines, which showed significant negative correlation for all lipid classes. At 35°C the stearate percentages of PC and PE were significantly negatively correlated with germination.

**Table 3.** Soybean seed germination, seedling growth rate (SGR) and conductivity tests of various types of soybean seeds

Sample classification	Sample ID	Germination (%)			SGR (mg/seedling)			Conductivity ( $\mu$ Siemens, 24 h)
		15°C	25°C	35°C	15°C	25°C	35°C	
Typical	Sturdy	99.3	98.0	80.0	21.0	48.7	40.7	314.5
	STE07	96.7	95.3	76.0	22.3	48.1	43.4	431.7
	STE15	97.3	97.3	59.3	23.7	47.6	44.5	328.8
	Kenwood	98.7	96.7	46.7	23.4	50.1	43.9	305.6
Reduced 16:0 <sup>a</sup>	PAL79	82.7	42.0	17.3	14.3	41.4	37.9	594.4
	PAL22	98.7	94.7	72.7	22.7	43.1	40.3	364.6
Elevated 16:0	PAL53	99.3	91.3	75.3	21.9	42.1	41.6	434.8
	PAL82	95.3	82.7	26.0	19.4	47.5	52.9	519.7
Elevated 18:0	STE39	96.0	93.3	64.0	24.4	46.5	37.2	329.2
	STE62	97.3	92.7	33.3	23.2	46.3	43.1	467.7
	STE73	98.7	86.7	6.0	20.9	45.9	38.1	504.8
	STE71	88.7	88.7	38.0	17.8	40.5	42.0	363.5
	LSD <sub>0.05</sub>	3.7	8.2	19.4	2.2	4.2	8.0	37.8

<sup>a</sup>See Table 1 for explanation of the fatty acid symbols.

**Table 4.** Correlation ( $R^2$ ) of neutral and polar lipid saturated fatty acid percentages with germination, SGR and conductivity tests

	Germination (%)			SGR (mg/seedling)			Conductivity ( $\mu$ Siemens)
	15°C	25°C	35°C	15°C	25°C	35°C	
TG 16:0 <sup>b</sup>	n.s. <sup>a</sup>	-0.85	n.s.	-0.80	n.s.	n.s.	n.s.
TG 18:0	n.s.	-0.92	n.s.	n.s.	-0.79	n.s.	n.s.
TG 16:0 + 18:0	n.s.	-0.84	-0.65	-0.63	n.s.	n.s.	n.s.
PC 16:0	n.s.	-0.82	n.s.	-0.81	n.s.	n.s.	n.s.
PC 18:0	n.s.	-0.90	-0.76	n.s.	n.s.	n.s.	n.s.
PC 16:0 + 18:0	n.s.	-0.85	n.s.	n.s.	n.s.	n.s.	0.63
PE 16:0	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
PE 18:0	n.s.	-0.91	-0.76	n.s.	n.s.	n.s.	n.s.
PE 16:0 + 18:0	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
PI 16:0	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
PI 18:0	n.s.	-0.85	n.s.	n.s.	-0.76	n.s.	n.s.
PI 16:0 + 18:0	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.

<sup>a</sup>n.s. denotes not statistically significant at 5% level.

<sup>b</sup>See Table 1 for explanation of the fatty acid symbols.

SGRs were negatively correlated with saturate content at 15 and 25°C for all lipid classes, although only a few were statistically significant. At 15°C, palmitate in TAG and PC was significantly negatively correlated with SGR, whereas at 25°C, stearate in TAG and PI was significantly negatively correlated. More soybean lines should be included in future experiments in order to demonstrate a more definite effect of saturate percentage on seed performance.

Conductivity of seed leachate was generally not significantly correlated with saturate content, except

for PC total saturates. Nevertheless, the correlations were all positive. Possibly, increased saturate in PC makes membranes more permeable.

In general, the lines with altered fatty acid compositions did well in the tests of seed viability and vigour, but under some conditions there was a significant negative correlation between elevated lipid saturates and germination, SGR and electrolyte retention. Some of these data suggest that increased stearate percentage may be more detrimental to the viability and vigour of soybean seeds than increased

palmitate. These effects may be related to the changes elevated saturates had caused in the physical properties of the TAG and PL classes (Wang, 1998). Hypothetically, soybeans with high saturate percentages should perform better at a relatively high temperature because their membranes would be more fluid, and they might be too rigid at low temperatures.

### Acknowledgement

This work was supported by a research grant from Pioneer Hi-Bred International, Inc., Des Moines, IA, USA.

### References

- AOSA (Association of Official Seed Analysts) (1986) Rules for testing seeds. *Proceedings of the Association of Official Seed Analysts* 6, 1–126.
- Chapman, D. (1973) Some recent studies of lipids, lipid-cholesterol and membrane systems. pp. 91–144 in Chapman, D.; Wallach, D.F.H. (Eds) *Biological membranes*, volume 2. New York, Academic Press.
- Dornbos, D.L. (1988) Soybean seed yield viability and vigor, and chemical composition resulting from drought and high temperature stress during seed fill. PhD dissertation, Iowa State University, Ames, IA.
- Fehr, W.R., Welke, G.A., Hammond, E.G., Duvick, D.N. and Cianzio, S.R. (1991) Inheritance of elevated palmitic acid content in soybean seed oil. *Crop Science* 31, 1522–1524.
- Hammond, E.G. (1991) Organization of rapid analysis of lipids in many individual plants. *Modern Methods of Plant Analysis* 12, 321–329.
- Hammond, E.G. (1992) Genetic alteration of food fats and oils. pp. 313–327 in Chow, C.K. (Ed.) *Fatty acids in foods and their health implications*. New York, Marcel Dekker.
- Hammond, E.G. and Fehr, W.R. (1975) Oil quality improvement in soybeans – *Glycine max* (L.) Merr. *Fette Seifen Anstrichmittel Verbunden Mit Der Zeitschrift die Ernährungsindustrie* 77, 97–101.
- Hartmann, R.B., Fehr, W.R., Welke, G.A., Hammond, E.G., Duvick, D.N. and Cianzio, S.R. (1996) Association of elevated palmitate content with agronomic and seed traits of soybean. *Crop Science* 36, 1466–1470.
- Hu, F.B., Stampfer, M.J., Manson, J.E., Rimm, E., Colditz, G.A., Rosner, B.A., Hennekens, C.H. and Willett, W.C. (1997) Dietary fat intake and the risk of coronary heart disease in women. *New England Journal of Medicine* 337, 1491–1499.
- Hui, Y.H. (1996) Lecithins. pp. 316–317 in Hui, Y.H. (Ed.) *Bailey's industrial oil and fat products*, volume 1. Edible oil and fat: General application. New York, John Wiley & Sons, Inc.
- Kok, L. (1998) *Trans-free margarine from highly saturated soybean oil*. MSc thesis, Iowa State University, Ames, IA.
- McKersie, B.D. and Senaratna, T. (1983) Membrane structure in germinating seed. *Recent Advances in Phytochemistry* 17, 29–52.
- Mounts, T.L., Abidi, S.L. and Rennick, K.A. (1996) Effect of genetic modification on the content and composition of bioactive constituents in soybean oil. *Journal of the American Oil Chemists' Society* 73, 581–586.
- Rebetzke, G.J., Pantalone, V.R., Burton, J.W., Carver, B.F. and Wilson, R.F. (1996) Phenotypic variation for saturated fatty acid content in soybean. *Euphytica* 91, 289–295.
- Rebetzke, G.J., Burton, J.W., Carter, T.E. and Wilson, R.F. (1998) Genetic variation for modifiers controlling reduced saturated fatty acid content in soybean. *Crop Science* 38, 303–308.
- SAS (1984) *SAS user's guide*. Cary, NC, Statistics Institute Inc.
- Shen, N., Fehr, W., Johnson, L. and White, P. (1997) Oxidative stabilities of soybean oils with elevated palmitate and reduced linolenate contents. *Journal of the American Oil Chemists' Society* 74, 299–302.
- Stoltzfus, D.L., Fehr, W.R. and Welke, G.A. (2000) Relationship of elevated palmitate to soybean seed traits. *Crop Science* 40, 52–54.
- Thompson, G.A. and Li, C. (1997) Altered fatty acid composition of membrane lipids in seeds and seedling tissue of high-saturate canolas. pp. 313–315 in Williams, J.P.; Khan, M.U.; Lem, N.W. (Eds) *Physiology, biochemistry and molecular biology of plant lipids*. Boston, Kluwer Academic Publishers.
- Wang, T. (1998) Phospholipid fatty acid composition of modified soybeans and the effect of saturated fatty acid contents on seed performance. PhD dissertation, Iowa State University, Ames, IA.
- Wang, T., Hammond, E.G. and Fehr, W.R. (1997) Phospholipid fatty acid composition and stereospecific distribution of soybeans with wide range of fatty acid composition. *Journal of the American Oil Chemists' Society* 74, 1587–1594.
- Wilcox, J.R., Cavins, J.F. and Nielsen, N.C. (1984) Genetic alteration of soybean oil composition by a chemical mutagen. *Journal of the American Oil Chemists' Society* 61, 97–100.
- Wilson, R.F., Burton, J.W. and Brim, C.A. (1981) Progress in the selection for altered fatty acid composition in soybeans. *Crop Science* 21, 788–791.

Received 3 September 1998  
 accepted after revision 25 September 2000  
 © CAB International, 2001



# Grassland Ecophysiology and Grazing Ecology

*Edited by G Lemaire, INRA, France, J Hodgson, Massey University, New Zealand,  
A Moraes, Universidade Federal do Parana, P C de F Cavalho and C Nabinger, Universidade Federal  
do Rio Grande do Sul, Brazil*

October 2000                      432 pages                      Hardback  
ISBN 0 85199 452 0  
£65.00 (US\$120.00)

**Readership:** Plant ecology/physiology, grassland science, animal science.

The sustainability and ecophysiology of grazing systems are of growing importance. This research level book presents edited, key papers from the International Symposium on Grassland Ecophysiology and Grazing Ecology held in Curitiba, Brazil in August 1999. It considers how plants within grasslands respond to and are adapted to grazing animals.

**This title:**

- Is of worldwide interest, covering both temperate and tropical pastures
- Includes contributors and leading authorities from North and South America, Europe and Australasia

**Contents:**

- Sustainability of grazing systems: goals, concepts and methods

**Part I: Environmental constraints and plant responses to defoliation**

- Effects of nitrogen and water supply on N and C fluxes and partitioning in defoliated swards
- An integrated view of C and N uses in leaf growth zones of defoliated grasses
- Effects of grazing on the roots and rhizosphere of grasses
- Reserve formation and recycling of carbon and nitrogen during regrowth of defoliated plants

**Part II: Morphogenesis of pasture species and adaptation to defoliation**

- Shoot morphological plasticity of grasses: leaf growth *vs.* tillering
- Tiller dynamics of grazed swards
- Effect of nitrogen on some morphogenetical traits of temperate and tropical perennial forage grasses
- Modelling the dynamics of temperate grasses and legumes in cut mixtures

**Part III: Plant-Animal interactions**

- Plant-animal interactions in complex plant communities: from mechanism to modelling
- Modeling spatial aspects of plant-animal interactions
- Defoliation patterns and herbage intake on pastures
- Selective grazing on grass-legume mixtures in tropical pastures

**Part IV: Sustainable grazing management of natural pastures**

- Leaf tissue turnover and efficiency of herbage utilization
- Dynamics of heterogeneity in a grazed sward
- Soil-plant-animal interactions and impact on nitrogen and phosphorus cycling and re-cycling in grazed pasture
- Sustainable management of pasture and rangelands

**Part V: Problems of Animal Production Related to Pastures in Subtropical and Temperate Regions of South America**

- *Campos* in Southern Brazil
- *Campos* in Uruguay
- Argentina's Humid Pampa

For further information or to order please contact CABI *Publishing*, UK or an exclusive CABI *Publishing* distributor in your area.

Please add £2.50 per book postage and packing (excluding UK).

CABI *Publishing*, CAB International, Wallingford, Oxon, OX10 8DE, UK

Tel: +44(0)1491 832111 Fax: +44(0)1491 829292 Email: [orders@cabi.org](mailto:orders@cabi.org)

CABI *Publishing*, CAB International, 10 East 40th Street, Suite 3203, New York, NY 10016, USA

Tel: +1 212 481 7018 Fax: +1 212 686 7993 Email: [cabi-nao@cabi.org](mailto:cabi-nao@cabi.org)