

2012

Swine Feed Efficiency: Influence of Market Weight

David R. Stender

Iowa State University, dstender@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/ipic_factsheets



Part of the [Agriculture Commons](#), and the [Animal Sciences Commons](#)

Recommended Citation

Stender, David R., "Swine Feed Efficiency: Influence of Market Weight" (2012). *Iowa Pork Industry Center Fact Sheets*. 9.
http://lib.dr.iastate.edu/ipic_factsheets/9

This Article is brought to you for free and open access by the Iowa Pork Industry Center at Iowa State University Digital Repository. It has been accepted for inclusion in Iowa Pork Industry Center Fact Sheets by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Swine Feed Efficiency: Influence of Market Weight

Abstract

To understand the reason that F/G becomes worse as pigs mature, it is important to understand the physiology of growth. During the growth cycle of a pig, the pounds of feed required per pound of gain increases, or said another way, the feed efficiency becomes worse as the pig's weight increases. Because it is more efficient to build muscle than fat, the efficiency of converting feed to live weight gain is best for young pigs and declines as pigs grow larger and older. Early growing pigs convert dietary feed at less than 2:1 F/G while finishing pigs convert feed at over 3:1 F/G. The poorer feed efficiency is caused by two major factors. First, the feed required for maintenance relative to lean growth increases as the pig gets larger. Second, the composition of gain shifts from primarily lean growth to a larger segment of lipid accretion as a pig approaches mature weight. Therefore, the mature frame size of the genetics is an important factor in feed efficiency near market weight. Early maturing pigs enter the less efficient fattening phase at a lighter final weight. Modern high lean genetics typically stay in the lean growth phase longer.

Keywords

IPIC 25h, Swine Feed Efficiency

Disciplines

Agriculture | Animal Sciences

Swine Feed Efficiency: Influence of Market Weight

Author
David R. Stender, Iowa State University

Reviewers
Mike D. Tokach, Kansas State University
Joel M. DeRouchey, Kansas State University

Introduction

To understand the reason that F/G becomes worse as pigs mature, it is important to understand the physiology of growth. During the growth cycle of a pig, the pounds of feed required per pound of gain increases, or said another way, the feed efficiency becomes worse as the pig's weight increases. Because it is more efficient to build muscle than fat, the efficiency of converting feed to live weight gain is best for young pigs and declines as pigs grow larger and older. Early growing pigs convert dietary feed at less than 2:1 F/G while finishing pigs convert feed at over 3:1 F/G. The poorer feed efficiency is caused by two major factors. First, the feed required for maintenance relative to lean growth increases as the pig gets larger. Second, the composition of gain shifts from primarily lean growth to a larger segment of lipid accretion as a pig approaches mature weight. Therefore, the mature frame size of the genetics is an important factor in feed efficiency near market weight. Early maturing pigs enter the less efficient fattening phase at a lighter final weight. Modern high lean genetics typically stay in the lean growth phase longer (Figure 1).

Understanding the differences in lean and fat deposition curves is important for feeding programs and marketing decisions. Genetics that have higher lean potential are more efficient in converting feed than those of lower lean gains. Therefore, the influence of market weight on feed efficiency is less for high lean pigs, because they enter the fattening phase of production at higher market weights. The decision for optimal market weight is influenced by the feed efficiency curve, the packer grid/matrices and the feed cost per pound of gain.

Feed Efficiency at Market Weight

Table 1 provides an example of pig performance expectations for high lean pigs in a commercial production system from 240 to 280 lb. Although feed efficiency increases as pigs get heavier, it is not at the same rate as other genetics that mature earlier.

Figure 1. Genetic Potential Impacts Backfat Depth




Table 1. Feed to Gain Ratios for Contemporary High Lean Genetics (data collected by Dr. Matt Swantek)

Item	Weight Range, lb			
	240-250	250-260	260-270	270-280
Average daily gain, lb	1.97	1.95	1.94	1.92
Average daily feed intake, lb	6.82	7.12	7.48	7.84
Feed/gain	3.46	3.65	3.87	4.09

The weight range in Table 1 covers the weight range of most packer grids with F/G increasing from 3.46 to 4.09; an 18.2% loss in feed efficiency comparing a 245 lb pig to a 275 lb pig. The impact of F/G at heavier weight will also impact the cumulative feed efficiency on the final closeout. Using the same data as in Table 1, cumulative wean-to-finish feed efficiency increases as market weight increases (Table 2).

Table 2. Cumulative Performance at Different Final Weights.

Item	Weight Range, lb			
	250	260	270	280
Average daily gain, lb	1.49	1.50	1.51	1.53
Average daily feed intake, lb	3.85	3.95	4.06	4.17
Feed/gain	2.59	2.63	2.68	2.73
Days on feed	160	165	170	176

In contrast to the data from modern, high lean pigs in Table 2, data from Headley et al. (1961) shows the extreme difference in genetic improvement over the last 50 years. Their data showed that F/G was 3.5 for a 160 lb pig, 4.1 for a 200 lb pig, and 4.4 for a 220 lb pig.

Because there are genetic maturity differences and different feeding strategies between lines of pigs, F/G from 250 to 280 will vary. There are only two methods to determine the final stage F/G ratios: incrementally weigh feed disappearance and pigs over time, or use accurate closeout records. It is important to calculate the incremental F/G at different end weights for your genetics and feed management. This can be done by mathematically converting cumulative F/G to incremental ratios. The incremental ratios can then be used as part of the target weight marketing decision.

Removing Heavy Pigs From Pens Improves F/G

Another consideration related to market end weight F/G is marketing strategy. The first marketing of the barn is important, as the heaviest pigs must be properly identified and a relatively constant number of pigs should be removed from every pen. Once removed, growth rate and F/G will improve for pigs remaining in the pen (DeDecker et al., 2005). Removing 25% of the pigs at 19 days before marketing resulted in more total weight marketed (0.3%) and a 17 lb feed savings per pig in the study. Higher removal rates had more feed savings (52.6 lb), but less pen weight sold (3.3%). Initial stocking density (space per pig) will have a major influence on the impact of sorting strategy on F/G. As stocking density increases, the benefit of removing pigs on growth of the remaining pigs in the pen increases.

Conclusion

Good decision making regarding target market weight is multi-faceted. For maximizing profit, consideration of all the factors involved is critical, including: feed efficiency, return over variable cost, packer grids and their premium and discount schedules, ability of stockman to sort heavy pigs, seasonal market cycles and pig removal strategies.

Understanding on-farm growing and finishing pig feed efficiency and growth curves are important to determine the optimal marketing strategy. High lean genetics may be taken to heavier weights with less effect on feed efficiency. Every operation should independently evaluate the F/G near market weight from records as an important piece of determining target end weight. Other considerations when making a target weight decision include: knowing and utilizing pigs removal strategies, selection strategies to market the correct pigs, packer weight grids (including discounts and premiums), and seasonal market cycles.

References

- DeDecker, J.M., M. Ellis, B.F. Wolter, B.P. Corrigan, S.E. Curtis, E.N. Parr, and D.M. Webel. 2005. Effects of proportion of pigs removed from a group and subsequent floor space on growth performance of finishing pigs. *J. Anim. Sci.* 83:449-454.
- Headley, V.E., E.R. Miller, D.E. Ullrey and J.A. Hofer. 1961. Application of the equation of the curve of diminishing increment to swine nutrition. *J. Anim. Sci.* 20:311.
- Latorre, M.A., R. Lázaro, D.G. Valencia, P. Medel, and G.G. Mateos. 2004. Meat Science and Muscle Biology The effects of gender and slaughter weight on the growth performance, carcass traits, and meat quality characteristics of heavy pigs. *J. Anim. Sci.* 82:526-533.

... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410, or call 800-795-3272 (voice) or 202-720-6382 (TDD). USDA is an equal opportunity provider and employer. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa