

2000

Replacement and deletion mutations in the catalytic domain and belt region of *Aspergillus awamori* glucoamylase to enhance thermostability

Hsuan-Liang Liu
Iowa State University

Yann Doleyres
Iowa State University

Pedro M. Coutinho
Iowa State University

See next page for additional authors

Follow this and additional works at: http://lib.dr.iastate.edu/cbe_pubs

 Part of the [Biochemical and Biomolecular Engineering Commons](#), [Biological Engineering Commons](#), [Food Chemistry Commons](#), and the [Human and Clinical Nutrition Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/cbe_pubs/30. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Chemical and Biological Engineering at Iowa State University Digital Repository. It has been accepted for inclusion in Chemical and Biological Engineering Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Replacement and deletion mutations in the catalytic domain and belt region of *Aspergillus awamori* glucoamylase to enhance thermostability

Abstract

Three single-residue mutations, Asp71→Asn, Gln409→Pro and Gly447→Ser, two long-to-short loop replacement mutations, Gly23-Ala24-Asp25-Gly26-Ala27-Trp28-Val29-Ser30→Asn-Pro-Pro (23–30 replacement) and Asp297-Ser298-Glu299-Ala300-Val301→Ala-Gly-Ala (297–301 replacement) and one deletion mutation removing Glu439, Thr440 and Ser441 (Δ 439–441), all based on amino acid sequence alignments, were made to improve *Aspergillus awamori* glucoamylase thermostability. The first and second single-residue mutations were designed to introduce a potential *N*-glycosylation site and to restrict backbone bond rotation, respectively, and therefore to decrease entropy during protein unfolding. The third single-residue mutation was made to decrease flexibility and increase *O*-glycosylation in the already highly *O*-glycosylated belt region that extends around the globular catalytic domain. The 23–30 replacement mutation was designed to eliminate a very thermolabile extended loop on the catalytic domain surface and to bring the remainder of this region closer to the rest of the catalytic domain, therefore preventing it from unfolding. The 297–301 replacement mutant GA was made to understand the function of the random coil region between α -helices 9 and 10. Δ 439–441 was constructed to decrease belt flexibility. All six mutations increased glucoamylase thermostability without significantly changing enzyme kinetic properties, with the 23–30 replacement mutation increasing the activation free energy for thermoinactivation by about 4 kJ/mol, which leads to a 4°C increase in operating temperature at constant thermostability.

Keywords

Aspergillus awamori, belt region, catalytic domain, glucoamylase, mutations, thermostability

Disciplines

Biochemical and Biomolecular Engineering | Biological Engineering | Chemical Engineering | Food Chemistry | Food Science | Human and Clinical Nutrition

Comments

This is a post-print of an article from *Protein Engineering*, 13, no. 9 (2000): 655–659, doi: [10.1093/protein/13.9.655](https://doi.org/10.1093/protein/13.9.655).

Authors

Hsuan-Liang Liu, Yann Doleyres, Pedro M. Coutinho, Clark Ford, and Peter J. Reilly

Replacement and Deletion Mutations in the Catalytic Domain and Belt Region of *Aspergillus awamori* Glucoamylase to Enhance Thermostability

Hsuan-Liang Liu, Yann Doleyes, Pedro M. Coutinho, Clark Ford¹, and Peter J. Reilly²

Departments of Chemical Engineering and ¹Food Science and Human Nutrition, Iowa State University, Ames, IA 50011, USA

²To whom correspondence should be addressed (Phone: 1-515-294-5968; fax: 1-515-294-2689; e-mail: reilly@iastate.edu)

Running head: **Mutations enhancing glucoamylase thermostability**

Three single-residue mutations, Asp71→Asn, Gln409→Pro, and Gly447→Ser, two long-to-short loop replacement mutations, Gly23-Ala24-Asp25-Gly26-Ala27-Trp28-Val29-Ser30→Asn-Pro-Pro (23-30 replacement) and Asp297-Ser298-Glu299-Ala300-Val301→Ala-Gly-Ala (297-301 replacement), and one deletion mutation removing Glu439, Thr440, and Ser441 (Δ 439-441), all based on amino acid sequence alignments, were made to improve *Aspergillus awamori* glucoamylase thermostability. The first and second single-residue mutations were designed to introduce a potential *N*-glycosylation site and to restrict backbone bond rotation, respectively, and therefore to decrease entropy during protein unfolding. The third single-residue mutation was made to decrease flexibility and increase *O*-glycosylation in the already highly *O*-glycosylated belt region that extends around the globular catalytic domain. The 23-30 replacement mutation was designed to eliminate a very thermolabile extended loop on the catalytic domain surface and to bring the remainder of this region closer to the rest of the catalytic domain, therefore preventing it from unfolding. The 297-301 replacement mutant GA was made to understand the function of the random coil region between α -helices 9 and 10. Δ 439-441 was constructed to decrease belt flexibility. All six mutations increased glucoamylase thermostability without significantly changing enzyme kinetic properties, with the 23-30 replacement mutation increasing the activation free energy for thermoinactivation by about 4 kJ/mol, which leads to a 4°C increase of operating temperature at constant thermostability.

Introduction

Glucoamylase [α -(1,4)-D-glucan glucohydrolase, EC 3.2.1.3, GA] is an industrially important, multidomain, *N*- and *O*-glycosylated enzyme that produces glucose from starch dextrin. A mutant GA with higher thermostability would be desirable because it could be used at higher temperatures, decreasing the required reaction time because of its higher activity.

GA undergoes irreversible thermoinactivation at moderately acidic pHs and 70°C (Munch and Tritsch, 1990). Neither deamidation (Chen *et al.*, 1994a,b) nor Asp-X peptide bond hydrolysis (Chen *et al.*, 1995a) determines the rate during GA thermoinactivation at 70°C and above, confirming earlier work (Munch and Tritsch, 1990). Aggregation and precipitation occur after the rate-determining step, because GA thermoinactivation follows first-order kinetics (Munch and Tritsch, 1990). Mismatched disulfide bonds are not found at 70°C and pH 3.5 and 4.5, which means that some other mechanism must control thermoinactivation at high temperatures. This mechanism appears to be a change in the protein structure such as unfolding that destroys the integrity of the active site (Munch and Tritsch, 1990; Williamson *et al.*, 1992b; Chen *et al.*, 1994a,b).

Knowledge of amino acid sequence homology and three-dimensional structures allows the prediction of important roles of specific amino acid residues on enzyme catalytic mechanism, thermostability, substrate selectivity, and structure-function relationships. The primary sequences of archaeal, eubacterial, and fungal GAs were aligned by Coutinho and Reilly (1994a,b, 1997) and have served along with the GA catalytic domain three-dimensional structure (Aleshin *et al.*, 1992) as important guides for various protein engineering studies of GA (Ford, 1999; Reilly, 1999).

Mutant GAs have been constructed to increase thermostability by eliminating thermoinactivation sites, by increasing α -helix stability, by introducing additional disulfide linkages, and by substituting Pro residues. Specifically, the *Aspergillus awamori*/*Aspergillus niger* (GAs from the two species are identical) Asn182→Ala and Asn182→Asp mutations increased GA thermostability up to 70°C, with specific activities similar to wild-type GA and the second with about 25% lower specific activity (Chen *et al.*, 1994a,b). Mutations Asp257→Gln, Asp293→Gln, and

Asp293→Glu gave slightly increased thermostability below 70°C at pH 4.5 with retention of about three-quarters of wild-type specific activity (Chen *et al.*, 1995a). Mutations Gly137→Ala, Gly139→Ala, and Gly137→Ala/Gly139→Ala, designed to stiffen α -helix 4, significantly increased thermostability at pH 4.5 well above 70°C without loss of specific activity, although performance at other pHs varied (Chen *et al.*, 1996). Mutation Ala246→Cys, proposed to create a disulfide linkage with Cys320, increased thermostability with retention of 27% and 85% of wild-type activity with maltose and maltoheptaose substrates, respectively (Fierobe *et al.*, 1996). Double mutation Asn20→Cys/Ala27→Cys was designed to create a disulfide bond on the catalytic domain surface, and successfully stabilized GA against unfolding (Li *et al.*, 1998). Mutations Ser436→Pro (Li *et al.*, 1997) and Ser30→Pro (Allen *et al.*, 1998) stabilized GA by reducing the conformational entropy of unfolding. In general, incorporating several beneficial mutations in one GA further increased thermostability and specific activity (Li *et al.*, 1998; Allen *et al.*, 1998).

This is one article of a series (Chen *et al.*, 1994a,b, 1995a, 1996; Li *et al.*, 1997, 1998; Allen *et al.*, 1998) on site-directed mutagenesis of *A. awamori* GA to improve GA thermostability. It reports construction and performance of three single-residue mutations, Asp71→Asn, Gln409→Pro, and Gly447→Ser, two loop-shortening replacement mutations, Gly23-Ala24-Asp25-Gly26-Ala27-Trp28-Val29-Ser30→Asn-Pro-Pro (23-30 replacement) and Asp297-Ser298-Glu299-Ala300-Val301→Ala-Gly-Ala (297-301 replacement), and one deletion mutation removing Glu439, Thr440, and Ser441 (Δ 439-441) designed by comparison with other GAs (Coutinho and Reilly, 1997). Figure 1 shows where each mutation occurs on the enzyme.

Materials and Methods

Materials

YEPM18, a yeast expression vector carrying the wild-type GA cDNA and the *Saccharomyces cerevisiae* strain C468 (α *leu2-3 leu2-112 his3-11 his3-15 mal-*) (Innis *et al.*, 1985) was a generous gift from Cetus Corporation. All restriction enzymes were purchased from Promega. Acar-

bose was a gift from Miles Laboratories. Peptide-*N*-glycosidase F was purchased from Oxford GlycoSystems.

Site-directed mutagenesis

Site-directed mutagenesis was performed with the Promega Altered Sites II *in vitro* mutagenesis system. An *Xba*I-*Hind*III fragment of pGEM-GA containing the wild-type GA cDNA (Fang and Ford, 1998) was inserted into the Promega pALTER-1 vector, making a GA cDNA-containing vector to be used as the double-stranded DNA template in the Altered Sites II system (Fang *et al.*, 1998b). The following mutagenic oligonucleotide primers were synthesized at the Iowa State University Nucleic Acid Facility: 5'-GCT CGT ACT ATC CTG AAT AAC ATC **AAT CCG CCC** GGC GCG GAC TCT GGC ATT GTC GTT GCT-3' (23-30 replacement), 5'-CTC TTC CGA AAT GGA **AAT** ACC AGT CTC CTC T-3' (Asp71→Asn), 5'-TAT ACC CTC AAC GAT GGT CTC AGT **GCT GGA GCT** GCG GTG GGT CGG TAC CCT GAG GAC-3' (297-301 replacement), 5'-AG TCT GAT GGC GAG CCG CTT TCC GCT CGC GA-3' (Gln409→Pro), 5'-GTC GTG CCT GCT TCT TGG GGC --- --- --- GCC AGC AGC GTG CCC GGC ACC TGT-3' (439-441 deletion), and 5'-GCC AGC AGC GTG CCC **AGC** ACC TGT GCG GCC A-3' (Gly447→Ser), the bold letters and the dashed lines indicating the changed and deleted nucleotides, respectively. All mutations were verified by DNA sequencing.

Expression and purification of GA

GA genes containing mutations were subcloned into the yeast expression vector YEpPM18 and transformed into *S. cerevisiae* strain C468 by electroporation, as described previously (Chen *et al.*, 1994b). Then mutant and wild-type GAs were produced in shaker flasks at 30°C and 170 rpm for 5 days without pH control. The cultures were centrifuged to remove the yeast cells, and then the culture supernatants were concentrated, dialyzed against 0.5 M NaCl/0.1 M NaOAc buffer (pH 4.4) with a 10-kDa cutoff Amicon S1 spiral ultrafiltration cartridge, and applied to acarbose-Sepharose affinity chromatography for purification (Chen *et al.*, 1994b). SDS-PAGE that demonstrated

enzyme homogeneity was carried out using 0.75 mm-thick polyacrylamide gels (Garfin, 1990).

Specific activities and kinetic parameters

GA specific activity at 50°C and pH 4.4 was determined with 4% (w/v) maltose (Fang *et al.*, 1998b). One unit (IU) was defined as the amount of enzyme required to produce 1 $\mu\text{mol}/\text{min}$ glucose under the conditions of the assay. Values of k_{cat} and K_{M} on maltose were obtained at 45°C and pH 4.4 in 0.05 M NaOAc buffer (Fang *et al.*, 1998a). Enzyme concentrations were determined by the Pierce bicinchoninic acid protein assay, using bovine serum albumin as the standard.

Irreversible thermoinactivation

Wild-type and mutant GAs (0.475 μM) were incubated for 12 min in 0.05 M NaOAc buffer, pH 4.4, at six temperatures between 65 and 77.5°C. Samples were taken at 2-min intervals, quickly chilled on ice, and then stored at 4°C for 24 h before being subjected to residual activity assay at 50°C (Liu *et al.*, 1998). The inactivation rate coefficients (k_d) for mutant and wild-type GAs were obtained from a semilogarithmic plot of residual activity versus inactivation time. The activation free energies (ΔG^\ddagger) for thermoinactivation of mutant and wild-type GAs were obtained from a semilogarithmic plot of k_d/T versus $1/T$.

De-N-glycosylation of Asp71→Asn mutant and wild-type GAs

Portions of 20 μg of Asp71→Asn mutant and wild-type GAs were added to 10 μl of 5X incubation buffer supplemented with 0.5% SDS and 5% β -mercaptoethanol, followed by denaturation of these two glycoproteins by heating at 100°C for 2 min. After the mixtures cooled to room temperature, 15 μl of peptide-*N*-glycosidase F was added to the two mixtures and each was incubated for 18 h at 37°C.

MALDI mass spectra

Asp71→Asn mutant and wild-type GAs, with and without de-*N*-glycosylation, were submitted to

the Protein Facility at Iowa State University to obtain MALDI mass spectra. Samples were dispersed in a large excess of a matrix material containing a chromophore for the laser light.

Results

GA specific activity and kinetics

The enzymatic properties of mutated and wild-type GAs are listed in Table I. Specific activities and catalytic efficiencies (k_{cat}/K_M) for all the mutated GAs are slightly higher than those of wild-type GAs, except for the Asp71→Asn mutated GA, which has the lowest catalytic efficiency, and the GA with mutation $\Delta 439-443$, which has the lowest specific activity. There is a positive correlation between specific activity and catalytic efficiency.

GA thermostability

The irreversible thermoinactivation of wild-type and mutated GAs follows first-order kinetics. Figure 2 shows the effect of temperature on k_d for wild-type and mutated GAs, while Table II shows the changes of the activation free energies for thermoinactivation ($\Delta\Delta G^\ddagger$) for mutated GAs relative to wild-type GA at 65 and 75°C. All the mutated GAs have positive $\Delta\Delta G^\ddagger$ values, meaning that they all are more thermostable than wild-type GA at both temperatures. The 23-30 replacement mutation gave the highest positive $\Delta\Delta G^\ddagger$ values of all the mutated GAs at both 65 and 75°C and therefore yielded the most thermostable mutated GA. All the mutated GAs except that with mutation Gly447→Ser have lower $\Delta\Delta G^\ddagger$ values at 75°C than at 65°C, indicating that their thermostabilities are less increased over that of wild-type GA by mutation at the higher temperature.

De-N-glycosylation and MALDI mass spectra

MALDI spectra of untreated and de-*N*-glycosylated Asp71→Asn mutant and wild-type GAs showed that the molecular weights of the two forms decreased by 2.2 and 2.3 kDa, respectively, upon *N*-deglycosylation. This suggests that Asp71→Asn mutant GA is as *N*-glycosylated as wild-

type GA. Therefore this mutation does not cause *N*-glycosylation of Asn71 as was possible. These results also confirm that the two *N*-glycosylation sites of wild-type GA have approximately thirteen sugar residues, as found by X-ray crystallography (Aleshin *et al.*, 1994).

Discussion

Of the three potential *N*-glycosylation sites in *Aspergillus* GAs, Asn171 and Asn395 but not Asn182 are *N*-glycosylated in *A. niger* (Svensson *et al.*, 1983), *A. awamori* var. *X100* (Aleshin *et al.*, 1992), and *A. awamori* GA, the latter expressed in *S. cerevisiae* (Chen *et al.*, 1994b). GAs from the first two sources have β -C1-linked *N*-acetyl-D-glucosamine residues initiating five- and eight-residue carbohydrate chains on Asn171 and Asn395, respectively (Aleshin *et al.*, 1994). Mutation Asn171→Ser GA has very similar catalytic efficiencies for malto- and isomaltooligosaccharide hydrolysis and very similar thermostability as wild-type *A. niger* GA (Stoffer *et al.*, 1997), while mutation Asn395→Gln dramatically decreases secretion as well as thermostability without changing specific activity (Chen *et al.*, 1994b), meaning that different *N*-glycosylation sites in the catalytic domain may play different roles in GA expression and unfolding.

In the present study, Asp71 between α -helices 2 and 3 in the *A. awamori* GA catalytic domain was replaced by Asn because GAs from *Neurospora crassa*, *Hormoconis resiniae*, *Humicola grisea*, *Rhizopus oryzae*, and *Saccharomycopsis fibuligera* all have Asn residues potentially involved in *N*-glycosylation at this position (Coutinho and Reilly, 1994b, 1997). This mutation did not yield *N*-glycosylation, but did slightly increase thermostability compared to wild-type GA while slightly decreasing specific activity.

A second single mutation was the replacement of Gln409 in *A. awamori* GA by Pro because GAs from *N. crassa*, *H. resiniae*, *H. grisea*, *Arxula adenivorans*, *Methanococcus jannaschii*, and *Clostridium* sp. have Pro at this position (Coutinho and Reilly, 1997). Pro residues restrict backbone bond rotation because of their pyrrolidine rings, and therefore they are believed to decrease entropy during protein unfolding by reducing the numbers of unfolded conformations that can be sampled by the protein (Matthews *et al.*, 1987). Suzuki *et al.* (1989, 1991) proposed that

increased numbers of Pro residues often correlate with increased protein thermostability. In general, the criteria used for the successful replacement of amino acid residues by Pro are that their values of ϕ , the dihedral angle of the C α -N bond, be $-80^\circ < \phi < -60^\circ$ (Balaji *et al.*, 1989) and that they be located in turns or random coils, the favored locations of Pro residues (Nicholson *et al.*, 1992; Ueda *et al.*, 1993).

The effect of the Gln409→Pro mutation on GA thermoinactivation shows that the backbone rigidity, greater space filling, and more hydrophobic interactions introduced by substituting Pro can help to prevent GA from unfolding under elevated temperatures.

Mutation Gly447→Ser GA was constructed to decrease flexibility and increase *O*-glycosylation in the already highly *O*-glycosylated belt region that extends around the catalytic domain between residues 441 and 470. The absence of a side chain in Gly gives the peptide backbone much greater conformational flexibility; therefore our purpose here was to investigate whether replacing Gly447 tended to limit possible conformations and therefore to stabilize GA. The replacement with Ser was also based on homology because GAs from *Aspergillus terreus*, *Aspergillus oryzae*, and *H. grisea* all have a Ser residue in position 447 (Coutinho and Reilly, 1997).

The increased thermostability caused by this mutation supports previous findings (Evans *et al.*, 1990; Libby *et al.*, 1994) that the belt region plays a role in GA thermostability by stabilizing a critical hydrophobic folding subunit (Coutinho and Reilly, 1997) and that *O*-linked carbohydrates appear to increase the rigidity and decrease the thermal vibration of the GA polypeptide chain (Williamson *et al.*, 1992a,b). In addition, this mutated GA has slightly higher specific activity and catalytic efficiency than wild-type GA, suggesting that this mutation does not significantly alter the structure of the enzyme around its active site or its interaction with the substrate. This result was expected because the belt region is located on the surface of the catalytic domain and does not play an important role in the GA catalytic mechanism.

Deletion mutations have been made to analyze the role of the belt, linker, and starch-binding domain of *A. awamori* GA (Evans *et al.*, 1990; Libby *et al.*, 1994; Chen *et al.*, 1995b). In this study, we constructed two long-to-short loop replacement mutations and one deletion mutation in

the same GA to improve its thermostability. A short-to-long loop replacement mutation had already been performed (Liu *et al.*, 1998), leading to a more selective but less active GA.

Previously made mutations suggest that the extended loop on the catalytic domain surface between α -helices 1 and 2 is critical for irreversible thermoinactivation and substrate selectivity of GA (Li *et al.*, 1998; Allen *et al.*, 1998; Liu *et al.*, 1998, 1999). It belongs to the most hydrophobic folding unit, which appears to strongly influence GA thermostability (Coutinho and Reilly, 1997). To further investigate the role of the loop and the region around it, a replacement mutation involving residues 23-30 was constructed to mimic the short amino acid sequence from *R. oryzae* GA. Previous studies led to two thermostable mutations, Asn20→Cys/Ala27→Cys (Li *et al.*, 1998) and Ser30→Pro (Allen *et al.*, 1998), and one thermosensitive mutation, Ala27→Pro (Li *et al.*, 1997), in this region. This extended loop exhibits a larger flexibility than its adjacent regions, shown by B-factors in *A. awamori* X100 GA crystallographic studies (Aleshin *et al.*, 1992, 1994), and takes a different position in the *S. fibuligera* GA structure (Sevčik *et al.*, 1997). The 23-30 replacement mutation was designed to eliminate the long loop, tightening this region together with the rest of the catalytic domain and therefore preventing it from unfolding.

The 23-30 replacement mutation increased specific activity slightly compared to wild-type GA and led to the most thermostable mutant GA constructed in this study, giving an approximately 4°C increase in operating temperature at constant thermostability. The current study further highlights the importance of this loop region on GA thermostability.

Residues Asp297 to Val301 occur in the random coil region of unknown function between α -helices 9 and 10. To understand the function of this region, we replaced residues 297-301 with three residues, Ala, Gly, and Ala, that mimic the amino acid sequence of *S. fibuligera* GA (Coutinho and Reilly, 1994a). The specific activity of this GA is only 20% that of *A. niger* GA on soluble starch (Solovicová *et al.*, 1996). Thus this region is perhaps important for GA specific activity and thermostability.

The 297-301 replacement mutation slightly increased specific activity and thermostability compared to wild-type GA, meaning that GA thermostability was increased by bringing two α -helices

closer together to increase unfolding entropy. The GA catalytic domain is composed of thirteen α -helices (52% of all its amino acid residues), with twelve of the helices folding into an unusual α/α barrel. Many of the remaining amino acids form β -sheets to link these α -helices (Aleshin *et al.*, 1992). The random coils between different α -helices may also play an important role in GA thermostability. For instance, mutation Ser436→Pro, changing a residue located in a random coil of unknown function, strongly decreases the entropy of unfolding by occupying more space and therefore enhancing hydrophobic interactions in this void (Li *et al.*, 1997).

Deletion mutation Δ 439-441 was made to decrease belt flexibility and *O*-glycosylation. This particular deletion was made because *Corticium rolfsii* GA does not have these three amino acids. A change in stability of this mutant GA should tell us more about the role played by the belt region in GA thermostability. Mutations in the belt region are interesting because they should not affect the center of the catalytic domain and thus should not modify specific activity. The belt region is important because this highly *O*-glycosylated domain may be involved in GA secretion, since the catalytic domain cannot be expressed extracellularly without this belt region (Libby *et al.*, 1994). In addition, its presence may prevent the catalytic domain from unfolding at higher temperatures.

Mutation Δ 439-441 increases GA thermostability more at lower than at higher temperatures. Three different effects may be involved. Reducing belt flexibility should decrease enzyme unfolding at all temperatures. However, the deletion of the three amino acids brings Gly438 adjacent to Ala442, possibly causing Gly-Ala bond hydrolysis as previously seen (Chen *et al.*, 1995a). Furthermore, the loss of two *O*-glycosylation sites caused by the deletion may partially counteract increased belt rigidity. Overall, both the Δ 439-441 and Gly447→Ser mutations strengthen the structure of the belt region of *A. awamori* GA and decrease unfolding of the catalytic domain.

In summary, we have successfully increased GA thermostability by three single-point, two replacement, and one deletion mutation. The 23-30 replacement mutation is the most thermostable mutation yet obtained in GA, along with the Ser30→Pro/Gly137→Ala and Asn20→Cys/Ala27→Cys/Ser30→Pro/Gly137→Ala mutations (Allen *et al.*, 1998), which also contain mutated residues in this part of GA. Therefore, the extended loop region on the catalytic domain surface be-

tween α -helices 1 and 2 plays a very important role in determining GA thermostability as well as substrate selectivity. The highly *O*-glycosylated belt region on the catalytic domain surface between residues 440 and 471 also has an important role in the mechanism governing the irreversible thermoinactivation of GA; this result strongly agrees with previously suggested functions of the *O*-glycosylation region, which include protecting GA from thermal denaturation. Amino acid sequence homology provides direct and important information in predicting the functions of some important amino acid residues.

Acknowledgements

This project was supported by the U. S. Department of Agriculture through the National Research Initiative Competitive Grants Program and by Genencor International, Inc. The authors thank James Meade for his gift of wild-type GA gene and plasmid, Jay Shetty for the gift of acarbose, and Alain Laederach for preparing Figure 1. Y. D. visited Iowa State University through an undergraduate exchange program between it and Ecole Polytechnique Fédérale de Lausanne, Switzerland.

References

- Aleshin, A., Golubev, A., Firsov, L. M. and Honzatko, R. B. (1992) *J. Biol. Chem.*, **267**, 19291-19298.
- Aleshin, A. E., Hoffman, C., Firsov, L. M. and Honzatko, R. B. (1994) *J. Mol. Biol.*, **238**, 575-591.
- Allen, M. J., Coutinho, P. M. and Ford, C. F. (1998) *Protein Eng.*, **11**, 783-788.
- Balaji, V. N., Mobasser, A. and Rao, S. N. (1989) *Biochem. Biophys. Res Commun.*, **160**, 109-114.
- Chen, H.-M., Bakir, U., Reilly, P. J. and Ford, C. (1994a) *Biotechnol. Bioeng.*, **43**, 101-105.
- Chen, H.-M., Ford, C. and Reilly, P. J. (1994b) *Biochem J.*, **301**, 275-281.
- Chen, H.-M., Ford, C. and Reilly, P. J. (1995a) *Protein Eng.*, **8**, 575-582.

- Chen, H.-M., Li, Y., Panda, T., Buehler, F. U., Ford, C. and Reilly, P. J. (1996) *Protein Eng.*, **9**, 499-505.
- Chen, L., Coutinho, P. M., Nikolov, Z. and Ford, C. (1995b) *Protein Eng.*, **8**, 1049-1055.
- Coutinho, P. M. and Reilly, P. J. (1994a) *Protein Eng.*, **7**, 393-400.
- Coutinho, P. M. and Reilly, P. J. (1994b) *Protein Eng.*, **7**, 749-760.
- Coutinho, P. M. and Reilly, P. J. (1997) *Proteins: Struct. Funct. Genet.*, **29**, 334-347.
- Evans, R., Ford, C., Sierks, M. R., Nikolov, Z. and Svensson, B. (1990) *Gene*, **91**, 131-134.
- Fang, T.-Y. and Ford, C. (1998) *Protein Eng.*, **11**, 383-388.
- Fang, T.-Y., Coutinho, P. M., Reilly, P. J. and Ford, C. (1998a), *Protein Eng.*, **11**, 119-126.
- Fang, T.-Y., Honzatko, R. B., Reilly, P. J. and Ford, C. (1998b) *Protein Eng.*, **11**, 127-133.
- Fierobe, H.-P., Stoffer, B. B., Frandsen, T. P. and Svensson, B. (1996) *Biochemistry*, **35**, 8696-8704.
- Ford, C. (1999) *Curr. Opin. Biotechnol.*, **10**, 353-357.
- Garfin, D. E. (1990) *Meth. Enzymol.*, **182**, 425-441.
- Innis, M. A., Holland, M. J., McCabe, P. C., Cole, G. E., Wittman, V. P., Tal, R., Watt, K., Gelfand, D. H., Holland, J. P. and Meade, J. H. (1985) *Science*, **228**, 21-26.
- Li, Y., Reilly, P. J. and Ford, C. (1997) *Protein Eng.*, **10**, 1199-1204.
- Li, Y., Coutinho, P. M. and Ford, C. (1998) *Protein Eng.*, **11**, 661-667.
- Libby, C. B., Cornett, C. A. G., Reilly, P. J. and Ford, C. (1994) *Protein Eng.*, **7**, 1109-1114.
- Liu, H.-L., Coutinho, P. M., Ford, C. and Reilly, P. J. (1998) *Protein Eng.*, **11**, 389-398.
- Liu, H.-L., Ford, C. and Reilly, P. J. (1999) *Protein Eng.*, **12**, 163-172.
- Matthews, B. W., Nicholson, H. and Becktel, W. J. (1987) *Proc. Natl. Acad. Sci. U.S.A.*, **84**, 6663-6667.
- Munch, O. and Tritsch, D. (1990) *Biochim. Biophys. Acta*, **1041**, 111-116.
- Nicholson, H., Tronrud, D. E., Becktel, W. J. and Matthews, B. W. (1992) *Biopolymers*, **32**, 1431-1441.
- Reilly, P. J. (1999) *Starch/Stärke*, **51**, 269-274.

- Sevčík, J., Hostinová, E., Gašperík, J., Solovicová, A., Wilson, K. S. and Dauter, Z. (1997) Structure 1AYX.
- Solovicová, A., Gašperík, J. and Hostinová, E. (1996) *Biochem Biophys. Res. Commun.*, **224**, 790-795.
- Stoffer, B., Dupont, C., Frandsen, T. P., Lehmbeck, J. and Svensson, B. (1997) *Protein Eng.*, **10**, 81-87.
- Suzuki, Y. (1989) *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.*, **65**, 146-148.
- Suzuki, Y., Hatagaki, K. and Oda, H. (1991) *Appl. Microbiol. Biotechnol.*, **34**, 707-714.
- Svensson, B., Larsen, K., Svendsen, I. and Boel, E. (1983) *Carlsberg Res. Commun.*, **48**, 529-544.
- Ueda, T., Tamura, T., Maeda, Y., Hashimoto, Y., Miki, T., Yamada, H. and Imoto, T. (1993) *Protein Eng.*, **6**, 183-187.
- Williamson, G., Belshaw, N. J. and Williamson, M. P. (1992a) *Biochem. J.*, **282**, 423-428.
- Williamson, G., Belshaw, N. J., Noel, T. R., Ring, S. G. and Williamson, M. P. (1992b) *Eur. J. Biochem.*, **207**, 661-670.

Table I. GA specific activities at 50°C and pH 4.4 using 4% maltose substrate, as well as GA kinetic parameters on maltose at 45°C and pH 4.4, both in 0.05 M NaOAc buffer.

| GA form | Specific activity (IU/mg GA) | k_{cat} (s ⁻¹) | K_{M} (mM) | $k_{\text{cat}}/K_{\text{M}}$ (s ⁻¹ mM ⁻¹) |
|---------------------|---------------------------------|--|------------------------|--|
| Wild-type | 20.6 ± 0.2 ^a | 8.67 ± 0.17 | 0.72 ± 0.03 | 12.0 |
| Asp71→Asn | 19.6 ± 0.6 | 7.59 ± 0.11 | 0.78 ± 0.04 | 9.7 |
| Gln409→Pro | 21.9 ± 0.8 | 8.71 ± 0.22 | 0.68 ± 0.02 | 12.8 |
| Gly447→Ser | 21.5 ± 0.9 | 9.13 ± 0.19 | 0.71 ± 0.04 | 12.9 |
| 23-30 replacement | 23.4 ± 0.7 | 9.32 ± 0.29 | 0.70 ± 0.05 | 13.3 |
| 297-301 replacement | 22.2 ± 0.8 | 9.38 ± 0.20 | 0.76 ± 0.06 | 12.3 |
| Δ439-441 | 18.4 ± 0.5 | 8.13 ± 0.13 | 0.69 ± 0.05 | 11.8 |


^aStandard error.

Table II. Changes of the activation free energy for thermoinactivation ($\Delta\Delta G^\ddagger$) at 65 and 75°C for mutated GAs relative to wild-type GA.


| GA form | $\Delta\Delta G^\ddagger$ (kJ/mol) | |
|---------------------|------------------------------------|------|
| | 65°C | 75°C |
| Asp71→Asn | 0.5 | 0.3 |
| Gln409→Pro | 1.8 | 1.2 |
| Gly447→Ser | 0.7 | 1.5 |
| 23-30 replacement | 4.4 | 3.5 |
| 297-301 replacement | 1.2 | 0.7 |
| Δ 439-441 | 1.0 | 0.7 |

Figure captions

1. Stereoscopic view of *A. awamori* GA, showing the mutations made in the enzyme. Replacement and deletion mutations are denoted by black shading. The mutated residues Asp71, Gln409, and Gly447 are located on a short loop between α -helices 3 and 4, a short β -strand between α -helices 12 and 13, and the belt, respectively.
2. Effect of temperature on thermostability of *A. awamori* GA at pH 4.4. Symbols: wild-type (○, _____), Asp71→Asn (△, _____), Gln409→Pro (□, _____), Gly447→Ser (◇, _____), 23-30 replacement (▲,), 297-301 replacement (◆, _____), Δ 439-441 (■, _____).



Liu et al.
Fig. 1



Liu et al.
Fig. 2