

9-13-2013

# Iowa Farmer Perception of Corn Rootworm Resistance

Erin W. Hodgson

*Iowa State University*, ewh@iastate.edu

Kristine J. P. Schaefer

*Iowa State University*, schaefer@iastate.edu

Aaron J. Gassmann

*Iowa State University*, aaronjg@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

---

## Recommended Citation

Hodgson, Erin W.; Schaefer, Kristine J. P.; and Gassmann, Aaron J., "Iowa Farmer Perception of Corn Rootworm Resistance" (2013). *Integrated Crop Management News*. 19.  
<http://lib.dr.iastate.edu/cropnews/19>

**The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.**

---

# Iowa Farmer Perception of Corn Rootworm Resistance

## **Abstract**

Every year Iowa State University (ISU) Extension and Outreach provides private pesticide applicator training covering a variety of pesticide safety and pest management issues. In response to reported incidences of Bt-resistant corn rootworm, a series of remote “clicker” questions were developed to survey current corn rootworm management practices and assess the degree to which farmers perceived resistance in 2012. The questions were asked at 153 pesticide applicator training sessions facilitated by extension field agronomists. More than 8,100 meeting participants responded to the six multiple-choice questions. Responses were anonymous and not everyone answered every question. Here are the statewide summaries of selected questions from the survey based on percentages of respondents:

## **Keywords**

Entomology

## **Disciplines**

Agricultural Science | Agriculture | Entomology

Subscribe to Crop  
News

#### Archives

**2014**

**2013**

**2012**

**2011**

**2010**

**2009**

**2008**

[Previous Years](#)

#### ISU Crop Resources

[Extension Field  
Agronomists](#)

[Crop & Soils Info](#)

[Pesticide Applicator  
Training](#)

[Agronomy Extension](#)

[Entomology Extension](#)

[Plant Pathology  
Extension](#)

[Ag and Biosystems  
Engineering Extension](#)

[Agribusiness  
Education Program](#)

[Iowa Grain Quality  
Initiative](#)

[College of Agriculture  
and Life Sciences](#)

[ISU Extension](#)

## Integrated Crop Management NEWS

 PRINT STORY

 EMAIL STORY

 ADD TO DELICIOUS

 ATOM FEED


 FOLLOW ON TWITTER

## Iowa Farmer Perception of Corn Rootworm Resistance

By Erin Hodgson, Kristine Schaefer and Aaron Gassmann, Department of Entomology

Every year Iowa State University (ISU) Extension and Outreach provides private pesticide applicator training covering a variety of pesticide safety and pest management issues. In response to reported incidences of Bt-resistant corn rootworm, a series of remote “clicker” questions were developed to survey current corn rootworm management practices and assess the degree to which farmers perceived resistance in 2012. The questions were asked at 153 pesticide applicator training sessions facilitated by extension field agronomists. More than 8,100 meeting participants responded to the six multiple-choice questions. Responses were anonymous and not everyone answered every question. Here are the statewide summaries of selected questions from the survey based on percentages of respondents:

To ascertain current management practices, participants were asked how they manage corn rootworm (Figure 1). Of the various management options for corn rootworm, 17 percent of respondents used Bt traits, 25 percent used crop rotation and 38 percent responded they used three or more methods.



**Figure 1. How do you manage corn rootworm?**

When asked if they had fields in 2012 where they suspected Bt trait failure, almost one-fourth (23 percent) of participants responded “yes” and 43 percent responded that they reported the suspected failure to someone. The majority of respondents who suspected trait failure either dug and rated root injury (38 percent) or noted goosenecking/lodging symptoms (33 percent) to assess the injury (Figure 2). About 13 percent of people who suspected a Bt trait failure were able to verify the larval injury (Figure 3). Although lodging can occur as a result of rootworm feeding injury, there may be other causes such as strong wind events. The only way to verify Bt trait performance issues is to evaluate root injury. Thus the rate of verified trait failures from this survey may be overestimated because larval root injury was not confirmed for all cases.

Additionally, the level of root injury that a grower may have considered sufficiently high to warrant a trait failure is unknown. Generally, a trait failure would be considered to be present if there is one node of root injury for Bt corn with a single trait targeting rootworm and 0.5 nodes for corn that has two Bt traits targeting rootworm. A [root rating scale](#) developed by the Iowa State University Department of Entomology is available.




Figure 2. What method did you use to assess corn rootworm injury?






Figure 3. What was the outcome of the assessment?

To understand where most of the perceived Bt trait failures were occurring within Iowa, we grouped responses by region. About one-third of people in northeast Iowa suspected a Bt trait failure in 2012 (Figure 4). Most of the positive verifications were in north central Iowa, with approximately one-third of people reporting Bt trait failures (Figure 5).



**Figure 4. Percent of respondents who suspected a corn rootworm trait failure in at least one of their cornfields in 2012.**



**Figure 5. Percent of respondents who positively verified corn rootworm trait failure.**

If you find injury to Bt corn on more than one node, consider diversifying pest management for that field. Severe injury may be due to western corn rootworm that has developed Bt resistance, and resistance should be suspected in these cases. The Department of Entomology at Iowa State University is working to understand the extent of resistance within the state and to develop management recommendations for Bt-resistant populations of western corn rootworm. Please contact Aaron Gassmann or Erin Hodgson to report fields with suspected resistance.

*Erin Hodgson is an associate professor of entomology with extension and research responsibilities; contact her at [ewh@iastate.edu](mailto:ewh@iastate.edu) or 515-294-2847. Kristine Schaefer is program manager of the pest management and the environment program and can be reached at [schaefer@iastate.edu](mailto:schaefer@iastate.edu) or 515-294-4286. Aaron Gassmann is an assistant professor of entomology with research and teaching responsibilities in insect pest management; contact him at [aaronjq@iastate.edu](mailto:aaronjq@iastate.edu) or 515-294-7623.*

This article was published originally on 9/13/2013. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.

---

Copyright ©2014 [Iowa State University Extension](#) | [Iowa State University](#)  
[Contact us](#) | [For Staff](#) | [Nondiscrimination and Information Disclosures](#) | [CMS Admin](#)  
Last Updated 9/14/2013