

11-2014

Monitoring Sheep and Culicoides Midges in Montana for Evidence of Bunyamwera Serogroup Virus Infection


Gregory D. Johnson
Montana State University - Bozeman

Charlie S. Bahnson
Iowa State University

Patricia Ishii
Iowa State University

Zachary N. Cochrane
Iowa State University, zcochran@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/vdpam_pubs

 [Part of the Entomology Commons](#), [Large or Food Animal and Equine Medicine Commons](#), [Veterinary Infectious Diseases Commons](#), and the [Veterinary Preventive Medicine, Epidemiology, and Public Health Commons](#)
See next page for additional authors

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/vdpam_pubs/42. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

Monitoring Sheep and Culicoides Midges in Montana for Evidence of Bunyamwera Serogroup Virus Infection

Abstract

A serological and entomological investigation was performed to monitor for potential Bunyamwera (BUN) serogroup virus activity in Montana. To facilitate the serological investigation, sera were collected from 104 sheep in 2013 and 2014 and assayed by plaque reduction neutralization test using all six BUN serogroup viruses known to occur in the United States: Cache Valley virus (CVV), Lokern virus (LOKV), Main Drain virus (MDV), Northway virus, Potosi virus and Tensaw virus. BUN serogroup virus-specific antibodies were detected in 41 (39%) sheep. Of these, three were seropositive for MDV, one was seropositive for CVV, one was seropositive for LOKV and 36 had antibodies to an undetermined BUN serogroup virus. Additionally, 30,606 *Culicoides sonorensis* were collected in 2013 using Centers for Disease Control and Prevention (CDC) light traps and assayed for cytopathic virus by virus isolation in African Green Monkey kidney (Vero) cells. All midges were negative. Almost one-third of the midges were further tested by reverse transcription-polymerase chain reaction using BUN serogroup virus-reactive primers and all were negative. We provide evidence of BUN serogroup virus infection in sheep but not *C. sonorensis* in Montana in 2013-2014. This study also provides the first evidence of CVV, MDV and LOKV activity in Montana.

Disciplines

Entomology | Large or Food Animal and Equine Medicine | Veterinary Infectious Diseases | Veterinary Preventive Medicine, Epidemiology, and Public Health

Comments

This article is from *Veterinary Record Open* 1 (2014): 71, doi:[10.1136/vetreco-2014-000071](https://doi.org/10.1136/vetreco-2014-000071). Posted with permission.

Rights

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Authors

Gregory D. Johnson, Charlie S. Bahnson, Patricia Ishii, Zachary N. Cochrane, D. Grant Hokit, Paul J. Plummer, Lyric Bartholomay, and Bradley J. Blitvich

Monitoring sheep and *Culicoides* midges in Montana for evidence of Bunyamwera serogroup virus infection

Gregory D. Johnson,¹ Charlie S. Bahnson,² Patricia Ishii,² Zachary N. Cochrane,² D. Grant Hokit,³ Paul J. Plummer,⁴ Lyric C. Bartholomay,⁵ Bradley J. Blitvich²

To cite: Johnson GD, *et al.* Monitoring sheep and *Culicoides* midges in Montana for evidence of Bunyamwera serogroup virus infection. *Vet Rec Open* 2014;**0**:e000071. doi:10.1136/vetreco-2014-000071

► Prepublication history and additional material for this paper are available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/vetreco-2014-000071>).

Received 14 July 2014
Revised 8 October 2014
Accepted 22 October 2014

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 3.0 Licence; see <http://vroepn.bmj.com>

ABSTRACT

Introduction: A serological and entomological investigation was performed to monitor for potential Bunyamwera (BUN) serogroup virus activity in Montana.

Results: To facilitate the serological investigation, sera were collected from 104 sheep in 2013 and 2014 and assayed by plaque reduction neutralization test using all six BUN serogroup viruses known to occur in the United States: Cache Valley virus (CVV), Lokern virus (LOKV), Main Drain virus (MDV), Northway virus, Potosi virus and Tensaw virus. BUN serogroup virus-specific antibodies were detected in 41 (39%) sheep. Of these, three were seropositive for MDV, one was seropositive for CVV, one was seropositive for LOKV and 36 had antibodies to an undetermined BUN serogroup virus. Additionally, 30,606 *Culicoides* sonorensis were collected in 2013 using Centers for Disease Control and Prevention (CDC) light traps and assayed for cytopathic virus by virus isolation in African Green Monkey kidney (Vero) cells. All midges were negative. Almost one-third of the midges were further tested by reverse transcription-polymerase chain reaction using BUN serogroup virus-reactive primers and all were negative.

Conclusions: We provide evidence of BUN serogroup virus infection in sheep but not *C. sonorensis* in Montana in 2013-2014. This study also provides the first evidence of CVV, MDV and LOKV activity in Montana.

INTRODUCTION

All viruses in the Bunyamwera (BUN) serogroup (genus *Orthobunyavirus*, family *Bunyaviridae*) are maintained in transmission cycles involving haematophagous arthropods and vertebrate hosts ([Schmaljohn and Nichol 2007](#)). Six BUN serogroup viruses occur in the USA: Cache Valley virus (CVV), Lokern virus (LOKV), Main Drain virus (MDV), Northway virus (NORV), Potosi virus (POTV) and Tensaw virus (TENV) ([Calisher and others 1986](#), [Francy and others 1990](#)). CVV infections in sheep can result in embryonic and fetal

death, stillbirths and congenital malformations ([Edwards and others 1989](#), [Chung and others 1990a, b](#)). MDV induces severe musculoskeletal and nervous system malformations in ovine fetuses infected in utero by injection into the amniotic vesicle and is a cause of equine encephalomyelitis but has not been associated with naturally occurring disease in sheep ([Emmons and others 1983](#), [Edwards and others 1997](#)). LOKV, POTV, NORV and TENV are not recognised ovine pathogens; however, their ability to cause disease has not been widely investigated. Because most veterinary diagnostic laboratories do not routinely test for BUN serogroup viruses, information on their ability to cause disease and data on their true disease incidence and seroprevalence in livestock are limited.

Most viruses in the BUN serogroup are transmitted primarily by mosquitoes ([Calisher and others 1986](#)). Several species of biting midges in the genus *Culicoides* also play an important role in BUN serogroup virus transmission. LOKV and MDV have been repeatedly isolated from *Culicoides* species, and *Culicoides variipennis* is a competent vector of MDV ([Mellor and others 1974](#), [Calisher and others 1986](#)). *Culicoides* species are also important vectors of several medically and veterinary important orthobunyaviruses in the Simbu serogroup (e.g. Akabane virus, Oropouche virus and Schmallenberg virus (SBV)) ([Mellor and others 2000](#), [Rasmussen and others 2012](#), [Veronesi and others 2013](#)). For instance, field studies in Denmark ([Rasmussen and others 2012](#)), Italy ([Goffredo and others 2013](#)) and Belgium ([De Regge and others 2012](#)) have implicated biting midges of the *C. obsoletus* group as relevant vectors of SBV and *Culicoides sonorensis* was demonstrated to be a suitable model vector species for SBV transmission ([Veronesi and others 2013](#)). *C. sonorensis* is an abundant species in Montana (Johnson, unpublished data), produced in the wet,



CrossMark

For numbered affiliations see end of article.

Correspondence to
Dr Bradley J Blitvich;
blitvich@iastate.edu

manure-contaminated soil surrounding stock ponds and other lentic water sources on farms and ranches.

There is no recent information on the seroprevalence of BUN serogroup viruses in livestock in Montana, nor are there any recent data of the significance of *Culicoides* species in BUN serogroup virus transmission in this region. Therefore, the overall goal of this study was to perform a serological and entomological investigation to determine the seroprevalence of BUN serogroup viruses in sheep and the prevalence of these viruses in *C. sonorensis* temporally and spatially associated with sheep in Montana.

MATERIALS AND METHODS

Sera collections

Sera were collected from sheep from a ranch in south central Montana in 2013 and a ranch in eastern Montana in 2014 (Fig 1). Several ranchers were contacted in the study area where *Culicoides* were collected regarding participation in the study. Some declined and others were unable to participate because of other commitments.

Plaque reduction neutralisation tests

Plaque reduction neutralisation tests (PRNTs) were performed using CVV (strain CVV-478), LOKV (strain FMS 4332), MDV (strain BFS 5015), NORV (strain 0234), POTV (strain BeAr7272) and TENV (strain A9-171b). CVV-478 was originally isolated from mosquitoes collected in Mexico (Farfan-Ale and others 2009, Blitvich and others 2012). All other viruses were obtained from the World Arbovirus Reference Collection at the University of Texas Medical Branch in Galveston, Texas. PRNTs were performed using all BUN serogroup viruses that occur in the USA (as opposed to only those associated with disease) because antibodies to one BUN serogroup virus often cross-react with other viruses in this serogroup due to their close antigenic relatedness (Hunt and Calisher 1979); thus, the exclusion of one or more viruses may result in serological misdiagnosis. PRNTs were performed in six-well plates containing confluent monolayers of African Green Monkey kidney (Vero) cells following published protocols (Beatty and others 1995). Initially, all sera were screened at a single dilution of 1:20. All sera that tested positive for BUN serogroup-specific antibodies were further diluted and tested by PRNT in order to identify the viruses responsible for these infections. Titres were expressed as the reciprocal of highest serum dilutions yielding ≥ 90 per cent reduction in the number of plaques (PRNT₉₀). For aetiological diagnosis, the PRNT₉₀ antibody titre to the respective virus was required to be at least four-fold greater than that to the other viruses tested.

Arthropod collections

Arthropods were collected at 10 trap sites in eight Montana counties from May to August 2013 (Fig 1). Five

arthropod collection sites were adjacent to or within 5 km of summer-pastured sheep. Collections were made using Centers for Disease Control and Prevention (CDC) light traps placed near the edges of stock ponds or other lentic water sources. Traps were baited with dry ice in the late afternoon, and arthropods were collected early the following morning and transported to the laboratory on dry ice. *C. sonorensis* were identified according to morphological characteristics and sorted into pools of up to 50 according to date and study site.

Homogenisations

C. sonorensis were placed in polypropylene, round-bottom 5 ml tubes with 1.8 ml phosphate-buffered saline (pH 7.4) supplemented with 100 units/ml penicillin, 100 µg/ml streptomycin and 2.5 µg/ml fungizone. Four 4.5 mm diameter copper-clad steel beads (BB-calibre airgun shot) were added to each tube, and midge pools were homogenised by vortexing for 60 seconds. Homogenates were centrifuged (3000 rpm, 10 minutes, 4°C) and supernatants were collected.

Virus isolation in cell culture

An aliquot (100 µl) of each homogenate was added to 0.5 ml of maintenance medium which consisted of Dulbecco's modified Eagle medium (DMEM; Invitrogen, Carlsbad, California, USA) supplemented with 2 per cent fetal bovine serum, 2 mM L-glutamine, 100 units/ml penicillin, 100 µg/ml streptomycin and 2.5 µg/ml fungizone. Supernatants were filtered using a 0.22 µm filter and inoculated onto subconfluent monolayers of Vero cells in 6-well plates. Cells were incubated for one hour at room temperature on an orbital shaker and then 5 ml of maintenance medium was added to each well. Cells were incubated at 37°C in the presence of 5 per cent CO₂ for 14 days and monitored regularly for cytopathic effect.

Reverse transcription PCR

Total RNA was extracted from homogenates using the QIAamp viral RNA extraction kit (QIAGEN, Valencia, California, USA) and analysed by RT-PCR using primers BCS82C and BCS332V, which are specific to orthobunyaviruses in the BUN and California serogroups (including CVV, MDV, POTV and TENV (Kuno and others 1996) and LOKV and NORV (B.J. Blitvich, unpublished data)). Complementary DNAs were generated using Superscript III reverse transcriptase (Invitrogen), and PCRs were performed using *Taq* polymerase (Invitrogen).

RESULTS

BUN serogroup virus-specific antibodies were identified in 41 (39.4 per cent) sheep (Table 1). Of these, three were seropositive for MDV, one was seropositive for CVV, one was seropositive for LOKV and 36 had antibodies to an undetermined BUN serogroup virus. Sixteen sheep that had antibodies to an undetermined BUN serogroup virus had low PRNT₉₀ titres to MDV and no detectable




FIG 1: Locations of *Culicoides* sampling sites (Δ) and premises for sheep sera collections (O)

antibody titres to any of the other viruses. Antibodies to NORV, POTV and TENV were not identified in any sheep.

A total of 30,606 *C. sonorensis* were collected and sorted into 629 pools. All were homogenised and an aliquot of each homogenate was tested by virus isolation in Vero cells. All homogenates were negative for cytopathic virus. A subset of homogenates (n=200) was further tested by RT-PCR using primers specific to BUN and California serogroup viruses. All homogenates were negative.

DISCUSSION

BUN serogroup virus-specific antibodies were identified in 41 (39.4 per cent) sheep. At least three viruses were responsible for these infections: CVV, LOKV and MDV. None of these viruses have been previously reported in Montana. CVV has been detected throughout much of North America including North and South Dakota which border Montana to the east, and Alberta and Saskatchewan which border Montana to the north (Iversen and others 1979, Calisher and others 1986, Pabbaraju and others 2009) but there are no published data demonstrating the occurrence of this virus in Montana. Previously, LOKV activity had only been reported in California, Colorado, New Mexico, Texas and Utah while MDV is known to occur in all of the aforementioned states as well as Arizona (Crane and others 1983, Calisher and others 1986, Kramer and others 1990). Our findings suggest that the geographic distribution of CVV, MDV and LOKV is wider than previously reported.

Because our data indicate that BUN serogroup viruses commonly infect sheep in Montana, it is likely that some of the sheep analysed in this study had been exposed to more than one BUN serogroup virus. However, information on the antibody responses in

vertebrates sequentially infected with BUN serogroup viruses is limited and research should be performed to address this issue. There is currently only one report that describes the antibody responses in vertebrates experimentally inoculated with different BUN serogroup viruses. The study was performed using white-tailed deer that were first inoculated with CVV or POTV then sequentially challenged with the alternate virus five to seven months later (Blackmore and Grimstad 1998).

As already noted, 36 sheep had antibodies to an undetermined BUN serogroup virus. Of these, 16 had low PRNT₉₀ titres to MDV and no detectable antibody titres to any of the other viruses. This could indicate that the sheep had been infected with MDV several years ago and, because neutralising antibody levels steadily decline over time (Gibbs and others 2005), only trace amounts of MDV-specific antibodies remained. Alternatively, the sheep could have been infected with a novel BUN serogroup virus that is more closely related to MDV than it is to the other viruses included in the PRNTs. In this regard, the three sheep seropositive for MDV had MDV PRNT₉₀ titres that could be considered low (all titres were 80); thus, we cannot dismiss the possibility that these sheep had instead been infected with an unrecognised MDV-like virus and that the MDV PRNT₉₀ titres are a consequence of serological cross-reactivity. Likewise, the sheep seropositive for LOKV had a LOKV PRNT₉₀ titre of 80 and, therefore, an unknown LOKV-like virus may have been responsible for this infection.

BUN serogroup viruses were not isolated from any *C. sonorensis* collected in this study. Our rationale for testing the midges by virus isolation in cell culture is because this technique is not restricted to the isolation of BUN serogroup viruses; other arthropod-transmitted viruses can also be detected. However, viral RNA is more stable than

**TABLE 1:** PRNT data for sheep with Bunyamwera serogroup virus-specific antibodies

Serum ID	PRNT ₉₀ titre						PRNT diagnosis
	CVV	LOKV	MDV	NORV	POTV	TENV	
A03	*—	—	40	—	—	—	†Und
A13	—	—	40	—	—	—	Und
A15	—	—	20	—	—	—	Und
A17	—	—	20	—	—	—	Und
A21	—	—	20	—	—	—	Und
A25	≥1280	160	40	160	—	20	CVV
A30	—	—	20	—	—	—	Und
A34	—	—	20	—	—	—	Und
A35	—	—	—	—	20	—	Und
A36	—	—	40	—	—	—	Und
A37	—	—	40	—	—	—	Und
A39	—	20	—	—	—	—	Und
A41	—	—	20	—	—	—	Und
A42	—	—	80	—	—	—	MDV
A44	—	40	—	—	—	—	Und
A45	—	—	80	—	—	—	MDV
A49	—	—	20	—	—	—	Und
A53	—	—	20	—	—	—	Und
A60	—	—	—	40	—	—	Und
B01	—	20	—	—	—	—	Und
B05	—	—	40	—	—	—	Und
B07	—	—	20	—	—	—	Und
B13	640	320	160	40	20	160	Und
B14	—	80	—	—	—	20	LOKV
B17	—	—	20	—	—	—	Und
B18	—	—	20	20	20	—	Und
B19	—	—	—	—	20	—	Und
B20	—	—	80	20	20	—	MDV
B21	—	20	—	—	—	—	Und
B22	—	—	—	20	—	—	Und
B24	40	20	20	80	20	40	Und
B25	—	—	—	—	20	—	Und
B29	—	—	—	—	20	—	Und
B30	—	20	—	—	—	—	Und
B31	—	—	—	—	20	—	Und
B32	—	—	—	20	—	—	Und
B35	—	20	20	—	—	—	Und
B36	—	—	20	—	—	—	Und
B37	—	—	20	—	20	—	Und
B38	—	20	20	—	20	—	Und
B39	—	—	—	—	20	—	Und

* <20

† Undetermined Bunyamwera serogroup virus

CVV, Cache Valley virus; LOKV, Lokern virus; MDV, Main Drain virus; NORV, Northway virus; POTV, Potosi virus; PRNT, plaque reduction neutralisation test; TENV, Tensaw virus

infectious viral particles and therefore a subset of *C. sonorensis* was further tested by RT-PCR using BUN serogroup-reactive primers. Another reason why we also decided to perform RT-PCRs is because this technique has the potential to detect novel BUN serogroup viruses that lack the capacity to replicate in Vero cells. However, as already noted, viral RNA was not detected in any midges.

In summary, we provide evidence of BUN serogroup virus infection in sheep but not *C. sonorensis* in Montana. At least three viruses were responsible for these infections: CVV, LOKV and MDV. This study provides the first

evidence of CVV, MDV and LOKV activity in Montana. Additional research is needed to determine the impact of these viruses on ovine health.

Author affiliations

¹Department of Animal and Range Sciences, Montana State University, Bozeman, Montana, USA

²Department of Veterinary Microbiology and Preventive Medicine, College of Veterinary Medicine, Iowa State University, Ames, Iowa, USA

³Department of Natural Science, Carroll College, Helena, Montana, USA

⁴Department of Veterinary and Diagnostic Production Animal Medicine, College of Veterinary Medicine, Iowa State University, Ames, Iowa, USA

⁵Department of Entomology, College of Agriculture and Life Sciences, Iowa State University, Ames, Iowa, USA

Acknowledgements The authors thank the undergraduate researchers at Aaniiih Nakoda College, Carroll College, Little Big Horn College and Salish Kootenai College who performed the trapping efforts in Montana, Marni Rolston for processing *Culicoides* collections, and Hayes Goosey and Lisa Surber for collecting sheep sera. The authors also thank Robert Tesh for providing isolates of LOKV, MDV, POTV, NORV and TENV. Financial support was provided by Boehringer Ingelheim, the Iowa Livestock Health Advisory Committee, the National Institute of General Medical Sciences of the National Institutes of Health (award number P20GM103474) and the Howard Hughes Medical Institute (award number 52007534).

Contributors GDJ and DGH coordinated the collection of sera and midges, CSB and ZNC performed the PRNTs, PI tested midges by virus isolation and RT-PCR, PJP and LCB participated in the design and coordination of the study, and BJB conceived of the study, participated in its design and coordination and drafted the manuscript. All authors read and approved the final manuscript.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Beatty B. J., Calisher C. H., Shope R. E. (1995) Arboviruses. In *Diagnostic Procedures for Viral and Rickettsial Diseases*. Ed E. Lennette. Washington, DC: American Public Health Association. 189–212
- Blackmore C. G., Grimstad P. R. (1998) Cache Valley and Potosi viruses (*Bunyaviridae*) in white-tailed deer (*Odocoileus virginianus*): experimental infections and antibody prevalence in natural populations. *The American Journal of Tropical Medicine and Hygiene* 59, 704–709
- Blitvich B. J., Loroño-Pino M. A., Garcia-Rejon J. E., Farfan-Ale J. A., Dorman K. S. (2012) Nucleotide sequencing and serologic analysis of Cache Valley virus isolates from the Yucatan Peninsula of Mexico. *Virus Genes* 45, 176–180
- Calisher C. H., Francy D. B., Smith G. C., Muth D. J., Lazuick J. S., Karabatsos N., Jakob W. L., Mclean R. G. (1986) Distribution of Bunyamwera serogroup viruses in North America, 1956–1984. *The American Journal of Tropical Medicine and Hygiene* 35, 429–443
- Chung S. I., Livingston C. W. Jr., Edwards J. F., Crandell R. W., Shope R. E., Shelton M. J., Collisson E. W. (1990a) Evidence that Cache Valley virus induces congenital malformations in sheep. *Veterinary Microbiology* 21, 297–307
- Chung S. I., Livingston C. W., Jr., Edwards J. F., Gauer B. B., Collisson E. W. (1990b) Congenital malformations in sheep resulting from in utero inoculation of Cache Valley virus. *American Journal of Veterinary Research* 51, 1645–1648
- Crane G. T., Elbel R. E., Francy D. B., Calisher C. H. (1983) Arboviruses from western Utah, USA, 1967–1976. *Journal of Medical Entomology* 20, 294–300
- De Regge N., Deblauwe I., De Deken R., Vantieghem P., Madder M., Geysen D., Smeets F., Losson B., Van Den Berg T., Cay A. B. (2012) Detection of Schmallenberg virus in different *Culicoides* spp. by real-time RT-PCR. *Transboundary and Emerging Diseases* 59, 471–475
- Edwards J. F., Karabatsos N., Collisson E. W., De La Concha Bermejillo A. (1997) Ovine fetal malformations induced by in utero inoculation with Main Drain, San Angelo, and LaCrosse viruses. *The American Journal of Tropical Medicine and Hygiene* 56, 171–176
- Edwards J. F., Livingston C. W., Chung S. I., Collisson E. C. (1989) Ovine arthrogyposis and central nervous system malformations associated with in utero Cache Valley virus infection: spontaneous disease. *Veterinary Pathology* 26, 33–39
- Emmons R. W., Woodie J. D., Laub R. L., Oshiro L. S. (1983) Main Drain virus as a cause of equine encephalomyelitis. *Journal of the American Veterinary Medical Association* 183, 555–558
- Farfan-Ale J. A., Loroño-Pino M. A., Garcia-Rejon J. E., Hovav E., Powers A. M., Lin M., Dorman K. S., Platt K., Bartholomay L. C., Soto V., Beatty B. J., Lanciotti R. S., Blitvich B. J. (2009) Detection of RNA from a novel West Nile-like virus and high prevalence of an insect-specific flavivirus in mosquitoes in the Yucatan Peninsula of Mexico. *The American Journal of Tropical Medicine and Hygiene* 80, 85–95
- Francy D. B., Karabatsos N., Wesson D. M., Moore C. G. Jr, Lazuick J. S., Niebylski M. L., Tsai T. F., Craig G. B., Jr (1990) A new arbovirus from *Aedes albopictus*, an Asian mosquito established in the United States. *Science* 250, 1738–1740
- Gibbs S. E., Hoffman D. M., Stark L. M., Marlenee N. L., Blitvich B. J., Beatty B. J., Stallknecht D. E. (2005) Persistence of antibodies to West Nile virus in naturally infected rock pigeons (*Columba livia*). *Clinical and Diagnostic Laboratory Immunology* 12, 665–667
- Goffredo M., Monaco F., Capelli G., Quaglia M., Federici V., Catalani M., Montarsi F., Polci A., Pinoni C., Calistri P., Savini G. (2013) Schmallenberg virus in Italy: a retrospective survey in *Culicoides* stored during the bluetongue Italian surveillance program. *Preventive Veterinary Medicine* 111, 230–236
- Hunt A. R., Calisher C. H. (1979) Relationships of Bunyamwera group viruses by neutralization. *The American Journal of Tropical Medicine and Hygiene* 28, 740–749
- Iversen J. O., Wagner R. J., Leung M. K., Hayles L. B., McIntock J. R. (1979) Cache Valley virus: isolations from mosquitoes in Saskatchewan, 1972–1974. *Canadian Journal of Microbiology* 25, 760–764
- Kramer W. L., Jones R. H., Holbrook F. R., Walton T. E., Calisher C. H. (1990) Isolation of arboviruses from *Culicoides* midges (Diptera: Ceratopogonidae) in Colorado during an epizootic of vesicular stomatitis New Jersey. *Journal of Medical Entomology* 27, 487–493
- Kuno G., Mitchell C. J., Chang G. J., Smith G. C. (1996) Detecting bunyaviruses of the Bunyamwera and California serogroups by a PCR technique. *Journal of Clinical Microbiology* 34, 1184–1188
- Mellor P. S., Boorman J., Baylis M. (2000) *Culicoides* biting midges: their role as arbovirus vectors. *Annual Review of Entomology* 45, 307–340
- Mellor P. S., Boorman J., Loke R. (1974) The multiplication of main drain virus in two species of *Culicoides* (Diptera, Ceratopogonidae). *Archiv für die gesamte Virusforschung* 46, 105–110
- Pabbaraju K., Ho K. C., Wong S., Fox J. D., Kaplen B., Tyler S., Drebot M., Tilley P. A. (2009) Surveillance of mosquito-borne viruses in Alberta using reverse transcription polymerase chain reaction with generic primers. *Journal of Medical Entomology* 46, 640–648
- Rasmussen L. D., Kristensen B., Kirkeby C., Rasmussen T. B., Belsham G. J., Bødker R., Bøtner A. (2012) *Culicoides* as vectors of Schmallenberg virus. *Emerging Infectious Diseases* 18, 1204–1206
- Schmaljohn C. S., Nichol S. T. (2007) Bunyaviridae. In *Fields Virology*. 5th edn. Ed D. M. Knipe. Philadelphia: Lippincott Williams and Wilkins. 1741–1789
- Veronesi E., Henstock M., Gubbins S., Batten C., Manley R., Barber J., Hoffmann B., Beer M., Attoui H., Mertens P. P., Carpenter S. (2013) Implicating *Culicoides* biting midges as vectors of Schmallenberg virus using semi-quantitative RT-PCR. *PLoS ONE* 8, e57747



Monitoring sheep and *Culicoides* midges in Montana for evidence of Bunyamwera serogroup virus infection

Gregory D. Johnson, Charlie S. Bahnson, Patricia Ishii, Zachary N. Cochrane, D. Grant Hokit, Paul J. Plummer, Lyric C. Bartholomay and Bradley J. Blitvich

Vet Rec Open 2014 1:
doi: 10.1136/vetreco-2014-000071

Updated information and services can be found at:
<http://vetrecordopen.bmj.com/content/1/1/e000071>

These include:

References

This article cites 23 articles, 12 of which you can access for free at:
<http://vetrecordopen.bmj.com/content/1/1/e000071#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections
[Open access](#) (47)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>