

6-10-2013

Nitrogen Loss—Spring 2013

John E. Sawyer

Iowa State University, jsawyer@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Soil Science Commons](#)

Recommended Citation

Sawyer, John E., "Nitrogen Loss—Spring 2013" (2013). *Integrated Crop Management News*. 35.
<http://lib.dr.iastate.edu/cropnews/35>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Nitrogen Loss—Spring 2013

Abstract

Dry soils across Iowa from deficit precipitation following the drought in 2012 have changed to excess wetness this spring. Unfortunately, precipitation to replenish subsoil moisture has been much more than needed. As I wrote in an [ICM News article](#) earlier this spring, one downside to the excess precipitation has been movement of carryover nitrate deeper into the profile and to tile lines, resulting in high nitrate-N concentrations in surface waters. This means there is less nitrate-N carryover to be utilized by 2013 crops, although as noted in the previous article, not all of the carryover N moved below the three-foot depth.

Keywords

Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Soil Science

[ICM Home](#)[ISU Extension Calendar](#)[Publications](#)[Extension News](#)[County Offices](#)[Contact Us](#)
[Subscribe to Crop News](#)

Archives

[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)

ISU Crop Resources

[Extension Field Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology Extension](#)[Ag and Biosystems Engineering Extension](#)[Agribusiness Education Program](#)[Iowa Grain Quality Initiative](#)[College of Agriculture and Life Sciences](#)[ISU Extension](#)

Integrated Crop Management NEWS

[PRINT STORY](#)
[EMAIL STORY](#)
[ADD TO DELICIOUS](#)
[ATOM FEED](#)
[FOLLOW ON TWITTER](#)

Nitrogen Loss - Spring 2013

By John Sawyer, Department of Agronomy

Dry soils across Iowa from deficit precipitation following the drought in 2012 have changed to excess wetness this spring. Unfortunately, precipitation to replenish subsoil moisture has been much more than needed. As I wrote in an [ICM News article](#) earlier this spring, one downside to the excess precipitation has been movement of carryover nitrate deeper into the profile and to tile lines, resulting in high nitrate-N concentrations in surface waters. This means there is less nitrate-N carryover to be utilized by 2013 crops, although as noted in the previous article, not all of the carryover N moved below the three-foot depth.

As in past wet springs, excess wetness can lead to loss of N applied for the current corn crop and result in greater than normal N response. I wrote several ICM News articles in the spring of 2008 and 2011 covering several aspects about N transformations in soil and N loss potential, and estimating the need for additional N application. Links to those articles are listed here.

www.extension.iastate.edu/CropNews/2013/0604Sawyer.htm
www.extension.iastate.edu/CropNews/2011/0613sawyer4.htm
www.extension.iastate.edu/CropNews/2011/0613sawyer3.htm
www.extension.iastate.edu/CropNews/2011/0613sawyer2.htm
www.extension.iastate.edu/CropNews/2011/0613sawyer1.htm

A web-based program to estimate sidedress N application for corn, called Adapt-N, is currently being promoted across the Corn Belt. It was developed at Cornell University in New York and is a combination of models that simulate soil N processes, influence of precipitation and corn N uptake. It has also recently been promoted as a way to estimate N losses from fall applied N. My evaluation the last three years indicates the program consistently and significantly underestimates needed N application (compared to economic optimal N rates at response trials). This brings into question if the program has been adequately calibrated and tested with Midwest conditions. If you try the program this spring, be cautious and consider use on only a limited trial basis.

The spring this year has been cold. Nitrification and denitrification are slow in cold soils, which will decrease potential for loss of applied N. But cold temperatures will not slow leaching of nitrate present in the soil. With slow conversion of ammonium to nitrate, there should be less loss of applied N – especially for recently applied ammonium fertilizers, such as anhydrous ammonia. Nitrification will also be less for applications such as late fall applied anhydrous ammonia and inclusion of a nitrification inhibitor. Early fall applied N, such as liquid manure, would have significant conversion to nitrate last fall. The nitrate component of UAN solution (28 or 32 percent N) is immediately subject to loss, and the nitrification rate of urea and ammonium in UAN will be more rapid than anhydrous ammonia. Therefore, loss potential is greater with such applications.

An aspect of this spring that is somewhat different from past wet springs is the late corn planting and slow growth. From my observations in north central

Iowa, the corn crop at this time is the smallest I remember in the 2000s. Corn N demand has been very low so far, and for late planted fields, season-long demand may be less due to lower productivity.

When warm, wet soils provide a good environment for microbial mineralization, the conversion of organic N to ammonium. Ammonium will accumulate under anaerobic conditions. While corn may die in ponded field areas, if replanted, the accumulated ammonium will supply crop-available N and perhaps an amount adequate to meet crop needs.

For corn fields with intended sidedress or split/sidedress N application, major loss conditions this spring have been avoided. For fields that had fall or early spring N applied, a visual way to check if corn will respond to additional N is to apply sidedress N fertilizer strips as soon as possible across several fields and watch the corn response. This can give a visual clue to potential for more N need and provide reference areas for mid-vegetative stage crop greenness measurement or canopy sensing.

John Sawyer is a professor of agronomy with research and extension responsibilities in soil fertility and nutrient management. He can be contacted at jsawyer@iastate.edu.

This article was published originally on 6/10/2013. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.