

10-28-2013

# Antimicrobial Effect of 7-O-Butylnaringenin, a Novel Flavonoid, and Various Natural Flavonoids against *Helicobacter pylori* Strains

Sun Hee Moon  
*Iowa State University*

Jae Hoon Lee  
*Konkuk University*

Kee-Tae Kim  
*Konkuk University*

Yong-Suhn Park  
*Konkuk University*

Seung-Yeol Nah  
*Konkuk University*

Follow this and additional works at: [http://lib.dr.iastate.edu/ans\\_pubs](http://lib.dr.iastate.edu/ans_pubs)

 next page for additional authors  
Part of the [Agriculture Commons](#), [Bioinformatics Commons](#), and the [Meat Science Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/ans\\_pubs/64](http://lib.dr.iastate.edu/ans_pubs/64). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

---

# Antimicrobial Effect of 7-O-Butylnaringenin, a Novel Flavonoid, and Various Natural Flavonoids against *Helicobacter pylori* Strains

## Abstract

Abstract: The antimicrobial effect of a novel flavonoid (7-O-butylnaringenin) on *Helicobacter pylori* 26695, 51, and SS1 strains and its inhibitory effect on the urease activity of the strains were evaluated and compared with those of several natural flavonoids. First, various flavonoids were screened for antimicrobial activities using the paper disc diffusion method. Hesperetin and naringenin showed the strongest antimicrobial effects among the natural flavonoids tested, and thus hesperetin and naringenin were selected for comparison with 7-O-butylnaringenin. The antimicrobial effect of 7-O-butylnaringenin was greater than that of the hesperetin and naringenin. *H. pylori* 51 was more sensitive to 7-O-butylnaringenin (2 log reduction of colony forming units,  $p < 0.05$ ) than the other two strains at 200  $\mu\text{M}$ . 7-O-Butylnaringenin also showed the highest inhibitory effect against urease activity of *H. pylori*. Morphological changes of *H. pylori* 26695 treated with these flavonoids indicated that both hesperetin and 7-O-butylnaringenin at 200  $\mu\text{M}$  damaged the cell membranes.

## Keywords

flavonoid, *Helicobacter pylori*, 7-O-butylnaringenin, hesperetin, antimicrobial effect, urease activity

## Disciplines

Agriculture | Animal Sciences | Bioinformatics | Meat Science

## Comments

This article is from *International Journal of Environmental Research and Public Health* 10 (2013): 5459–5469, doi:[10.3390/ijerph10115459](https://doi.org/10.3390/ijerph10115459). Posted with permission.

## Rights

This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Authors

Sun Hee Moon, Jae Hoon Lee, Kee-Tae Kim, Yong-Suhn Park, Seung-Yeol Nah, Dong U. Ahn, and Hyun-Dong Paik

Article

## Antimicrobial Effect of 7-*O*-Butylnaringenin, a Novel Flavonoid, and Various Natural Flavonoids against *Helicobacter pylori* Strains

Sun Hee Moon <sup>1,2</sup>, Jae Hoon Lee <sup>2</sup>, Kee-Tae Kim <sup>3</sup>, Yong-Sun Park <sup>2,3,4</sup>, Seung-Yeol Nah <sup>2,3,5</sup>, Dong Uk Ahn <sup>1,6</sup> and Hyun-Dong Paik <sup>2,3,\*</sup>

<sup>1</sup> Department of Animal Science, Iowa State University, Ames 50011, USA;  
E-Mails: smoon@iastate.edu (S.H.M.); duahn@iastate.edu (D.U.A.)

<sup>2</sup> Division of Animal Life Science, Konkuk University, Seoul 143-701, Korea;  
E-Mails: jokko77@naver.com (J.H.L.); parkyong@konkuk.ac.kr (Y.-S.P.);  
synah@konkuk.ac.kr (S.-Y.N.)

<sup>3</sup> Bio/Molecular Informatics Center, Konkuk University, Seoul 143-701, Korea;  
E-Mail: richard44@hanmail.net

<sup>4</sup> Department of Chemistry, Konkuk University, Seoul 143-701, Korea

<sup>5</sup> Department of Physiology, College of Veterinary Medicine, Konkuk University, Seoul 143-701, Korea

<sup>6</sup> Department of Animal Science and Technology, Sunchon National University,  
Sunchon 540-742, Korea

\* Author to whom correspondence should be addressed; E-Mail: hdpaik@konkuk.ac.kr;  
Tel.: +82-2-2049-6011; Fax: +82-2-455-3082.

Received: 5 August 2013; in revised form: 16 October 2013 / Accepted: 21 October 2013 /  
Published: 28 October 2013

---

**Abstract:** The antimicrobial effect of a novel flavonoid (7-*O*-butylnaringenin) on *Helicobacter pylori* 26695, 51, and SS1 strains and its inhibitory effect on the urease activity of the strains were evaluated and compared with those of several natural flavonoids. First, various flavonoids were screened for antimicrobial activities using the paper disc diffusion method. Hesperetin and naringenin showed the strongest antimicrobial effects among the natural flavonoids tested, and thus hesperetin and naringenin were selected for comparison with 7-*O*-butylnaringenin. The antimicrobial effect of 7-*O*-butylnaringenin was greater than that of the hesperetin and naringenin. *H. pylori* 51 was more sensitive to 7-*O*-butylnaringenin (2 log reduction of colony forming units,  $p < 0.05$ ) than the other two strains at 200  $\mu$ M. 7-*O*-Butylnaringenin also showed the

highest inhibitory effect against urease activity of *H. pylori*. Morphological changes of *H. pylori* 26695 treated with these flavonoids indicated that both hesperetin and 7-*O*-butylnaringenin at 200  $\mu$ M damaged the cell membranes.

**Keywords:** flavonoid; *Helicobacter pylori*; 7-*O*-butylnaringenin; hesperetin; antimicrobial effect; urease activity

---

## 1. Introduction

*Helicobacter pylori* is a micro-aerophilic, Gram-negative, spiral-shaped flagellated bacterium [1]. This microorganism lives in the stomach and duodenum and is known to be one of the most common chronic bacterial pathogens of humans. *H. pylori* is associated with several diseases, including chronic gastritis, peptic ulcers, and gastric mucosa associated lymphoid tissue lymphoma [2–5].

*H. pylori* is resistant to stomach acid because it is protected by the mucous cells and its urease activity [2]. Urease, which is the most characteristic feature of *H. pylori*, catalyzes the hydrolysis of urea to produce ammonia and CO<sub>2</sub>, and protects the bacteria from the acidic environment of the stomach [6,7]. Ammonia itself may cause tissue injury or be toxic to intercellular junctions [8]. However, ammonia generated by the urease buffers gastric acid, and thus is also important for the colonization of the bacteria in the stomach. The urease of *H. pylori* has been described as a highly active enzyme that may be associated with virulence [9] and is considered as a constitutive and permanently active enzyme [10]. The urease in *H. pylori* is a high-molecular weight enzyme that has a high affinity to urea and rapidly hydrolyzes it, but is highly sensitive to urease inhibitors. To treat the patients with gastro-duodenal diseases by *H. pylori*, therefore, inhibiting the infection, growth, and urease activity of *H. pylori* is important. Antimicrobial drugs have been used to treat *H. pylori* infections in recent years, and the successful eradication of this bacterium has been demonstrated to prevent the relapse of duodenal and gastric ulcers [10–12].

Many naturally occurring compounds found in dietary and medicinal plants, herbs and fruit extracts have been shown to possess antimicrobial activities [13–16]. Flavonoids are natural compounds ubiquitous in green plant cells [17]. Flavonoids appear to have antimicrobial, antioxidative, anti-inflammatory and anti-carcinogenic effects, and have played major roles in successful medical treatments since ancient times and their use has continued to these days [18–20]. There have been various studies on the functional effects of flavonoids with regard to their use by the health food and pharmaceutical industries [21–23]. In particular, it has been shown that certain flavonoids have antimicrobial effects against *H. pylori* [13,24,25]. Although the Minimum Inhibitory Concentration (MIC) of some flavonoids against the growth of *H. pylori* has been determined, the nature of the inhibitory effects has not been sufficiently studied [14]. In addition, a new chemically-derived flavonoid has recently been evaluated for its functional activities as a medicinal compound [19]. With this approach, the protective mechanism of some popularly used flavonoids (naringenin and hesperetin), and 7-*O*-butylnaringenin, a new modified flavonoid from naringenin, on the growth as well as the urease activity of *H. pylori* was studied.

## 2. Experimental Section


### 2.1. Bacterial Strains

*H. pylori* 26695, 51, and SS1 were purchased from the *H. pylori* Korean-Type Culture Collection (HpKTCC, Jinju, Korea). The strains were activated in brucella agar (Difco Laboratories, Detroit, MI, USA) plates supplemented with 5% (v/v) horse serum and was cultured under micro-aerophilic conditions (10% CO<sub>2</sub> atmosphere) for 3 days. For these studies, the strains were then inoculated in brucella broth supplemented with 5% horse serum and were cultured for 1 day at 37 °C before use.

### 2.2. Flavonoids

Nine different flavonoids were used for comparison in this study; kaempferol, and quercetin as flavonols, apigenin, luteolin, and 5,4'-dihydroxy-7-methoxyflavone (genkwanin) as flavones, and naringenin, hesperetin, and hesperidin as flavanones [26] (Figure 1).

**Figure 1.** Chemical structures of flavonoids used in this study. (A) kaempferol, (B) quercetin, (C) apigenin, (D) naringenin, (E) luteolin, (F) hesperetin, (G) hesperidin, (H) genkwanin, and (I) 7-*O*-butylnaringenin.



All flavonoids were purchased from Sigma Chemical Co. (St. Louis, MO, USA). 7-*O*-Butylnaringenin was supplied by the Chemistry Department in Konkuk University. 7-*O*-Butylnaringenin solution was prepared by mixing and stirring naringenin (500 mg, 1.84 mM), butyl bromide (1.5 equiv.) and K<sub>2</sub>CO<sub>3</sub> (1.0 equiv.) in dimethylformamide (DMF, 10 mL) for 12 h at 25 °C. The mixture was quenched by adding saturated aqueous NH<sub>4</sub>Cl and then extracted with ethyl acetate.

The extracts were then dried over anhydrous  $\text{MgSO}_4$ . After evaporation, the resulting crude product was purified using a flash column chromatography, which produced a colorless oil in 35% yield [19].

### 2.3. Screening by Paper Disc Diffusion Method

The paper disc diffusion method was used as a screening test for the antibacterial activity of flavonoids [2]. *H. pylori* was incubated as described above. Forty microliters of flavonoid sample were applied to a paper disc (8 mm in diameter) and the concentrations of flavonoids were 2.5, 5, 10, and 20 mM in dimethylsulfoxide (DMSO), respectively. The DMSO was removed by drying at 20 °C for 10 min, and the paper discs were placed on brucella agar plates supplemented with 5% horse serum inoculated with  $2.0 \times 10^7$  CFU/mL of each *H. pylori* strain. The zone of inhibition was determined after incubating the plates at 37 °C for 3 days under 10%  $\text{CO}_2$  incubator (MCO-18AIC; Sanyo, Oizumi-Machi, Japan).

### 2.4. Assay of Antimicrobial Effects on *H. pylori*

The initial cell numbers of *H. pylori* strains were adjusted to  $2.0 \times 10^5$  CFU/mL in broth. Four milliliters of brucella supplemented with 5% (v/v) horse serum, 1 mL of culture broth, and 50  $\mu\text{L}$  of flavonoid solution were added to each well and cultured at 37 °C under 10%  $\text{CO}_2$  atmosphere. The concentration of flavonoid was adjusted to 100 and 200  $\mu\text{M}$  in total broth per well. For the blank and control, 50  $\mu\text{L}$  of distilled water and DMSO were added instead of flavonoid solutions, respectively. After 24 h incubation, culture samples including the blank and control, were serially diluted in 0.1% peptone water and spread on brucella agar supplemented with 5% (v/v) horse serum. Plates were incubated for 3 days at 37 °C under 10%  $\text{CO}_2$  atmosphere [21]. The effect of flavonoids on the *H. pylori* strains was determined using the standard cell counting method.

### 2.5. Flavonoid Inhibition of Urease in *H. pylori*

Urease activity was measured by the method of Malekzadeh *et al.* [27] and Mobley *et al.* [28]. One thousand eight hundred microliters of urea broth (urea 20 g, NaCl 5 g,  $\text{KH}_2\text{PO}_4$  2 g, peptone 1 g, glucose 1 g, and phenol red 0.012 g per L solution), 200  $\mu\text{L}$  of the cultured broth, and 20  $\mu\text{L}$  of the flavonoid solutions were mixed in a test tube. The initial cell number of *H. pylori* was adjusted to  $2.0 \times 10^5$  CFU/mL reaction mixture, and the concentration of flavonoid was adjusted to 200  $\mu\text{M}$  for each reaction mixture. For control, 20  $\mu\text{L}$  of DMSO instead of flavonoid solution was added. After 3 h of incubation at 37 °C, the changes of optical density (pink red color) in urea broth by the ammonia produced were measured at 560 nm with a spectrophotometer (EL311; Bio-Tek Instruments Inc., Seoul, Korea).

### 2.6. Preparation of *H. pylori* for Scanning Electron Microscopy (SEM)

Bacterial cells were prepared for SEM examination using the methods of Tsugawa *et al.* [29]. Cell broth was cultured in brucella broth supplemented with 5% horse serum for 24 h at 37 °C under 10%  $\text{CO}_2$  and were treated with 200  $\mu\text{M}$  of hesperetin, naringenin, or 7-*O*-butylnaringenin. The cells were harvested by centrifugation at  $10,000 \times g$  for 10 min and fixed overnight with 25% (v/v)

glutaraldehyde at 4 °C. The cells were then washed with 0.1 M phosphate buffer (pH 7.0) three times and dehydrated by washing with ethanol. The ethanol concentration for washing was gradually increased from 10% to 100% in a sequential manner. The dehydrated specimens were treated with ion-spotter-coating at 7 V and 15 mA in a vacuum evaporator (E-1010, Hitachi Science System Ltd., Hitachinaka, Japan) and subjected to SEM (S-3000N, Hitachi Science System Ltd.).

### 2.7. Statistical Analysis

All results are presented as mean  $\pm$  standard error (SE), and statistical analysis was performed using the SPSS (Chicago, IL, USA) package for Windows (Ver. 18.0). The mean values were compared using one-way analysis of variance (ANOVA) followed by Duncan's multiple range tests with a significance defined at  $p < 0.05$ . Correlations were determined using Pearson's correlation coefficient.

## 3. Results and Discussion

### 3.1. Screening by Paper Disc Diffusion Method

Quercetin, kaempferol, naringenin, and hesperetin had stronger antimicrobial effects than the other flavonoids tested. The antimicrobial effects of naringenin and hesperetin were greater than those of other flavonoids used in this study. Apigenin, hesperidin, and genkwanin had no effects, even at 20 mM, against all *H. pylori* strains studied (Table 1).

**Table 1.** Zone of growth inhibition of *H. pylori* 26695, 51, and SS1.

Flavonoid	Conc. (mM)	<i>H. pylori</i> 26695	<i>H. pylori</i> 51	<i>H. pylori</i> SS1
Kaempferol	2.5	-	-	-
	5	++	-	+
	10	++	+	+
	20	++	+	++
Quercetin	2.5	-	-	-
	5	-	-	-
	10	-	-	+
	20	+	-	+
Apigenin	2.5	-	-	-
	5	-	-	-
	10	-	-	-
	20	-	-	-
Naringenin	2.5	-	-	-
	5	+++	+++	+++
	10	+++	+++	+++
	20	+++	+++	+++
Luteolin	2.5	-	-	-
	5	+	+	-
	10	++	++	+
	20	++	++	+

Table 1. Cont.

Flavonoid	Conc. (mM)	<i>H. pylori</i> 26695	<i>H. pylori</i> 51	<i>H. pylori</i> SS1
Hesperetin	2.5	-	-	-
	5	+++	+++	+++
	10	+++	+++	+++
	20	+++	+++	+++
Hesperidin	2.5	-	-	-
	5	-	-	-
	10	-	-	-
	20	-	-	-
Genkwanin	2.5	-	-	-
	5	-	-	-
	10	-	-	-
	20	-	-	-
7- <i>O</i> -Butylnaringenin	2.5	-	-	-
	5	-	++	+
	10	+	+	+
	20	++	++	++

-: No inhibition ( $\leq 8.9$  mm); +: Slight inhibition (9.0–14.9 mm); ++: Moderate inhibition (15–19.9 mm); +++: Strong inhibition ( $\geq 20$  mm).

In addition, the antimicrobial effects of flavonoids were slightly different among the *H. pylori* strains tested.

### 3.2. Evaluation of Antimicrobial Effects on *H. pylori* Strains


The antimicrobial effects of naringenin, hesperetin, and 7-*O*-butylnaringenin against three *H. pylori* strains indicated that naringenin below 200 mM was the least effective among the three flavonoids and the effect was not significantly different from with blank and control ( $p > 0.05$ , Figure 2(A)).

Hesperetin has a stronger antimicrobial effect than naringenin against *H. pylori*, although the effect differed between the strains (Figure 2(B)). Hesperetin (200  $\mu$ M) showed stronger antimicrobial effects to *H. pylori* SS1 than other two strains and reduced the number by 2.5 log colony forming units ( $p < 0.05$ ). 7-*O*-Butylnaringenin at 100–200  $\mu$ M showed a stronger antimicrobial effect than the other two flavonoids tested (Figure 2(C)). The antimicrobial effect of 7-*O*-butylnaringenin was >10-fold of that of unmodified naringenin, and 7-*O*-butylnaringenin at 200  $\mu$ M was highly effective against *H. pylori* 51 ( $p < 0.05$ ). The results indicated that 7-*O*-butylnaringenin was the most effective of all the flavonoids tested at any concentration. However, the effects of 7-*O*-butylnaringenin in the broth culture method were different from those in the disc diffusion method (Table 1). These observations are likely due to the differences in the diffusivity of flavonoids on agar culture and paper disc [22]. The inhibitory effects of flavonoids on the growth of microorganism can vary depending upon the type and size of molecular groups and the positions of their side chains on the flavonoid backbone structure [16,22]. Since many different assays are employed in flavonoid research, including the paper disk diffusion method, the agar dilution assay, the broth microdilution method, the results appear



widely conflicting. In particular, assays depending on the diffusion of flavonoids may not give a reliable quantitative measure of antibacterial activity because a strong antibacterial flavonoid may have a low rate of diffusion [23].

**Figure 2.** The antimicrobial effects of three flavonoids on growth of *H. pylori* 26695, 51, and SS1. Initial cell density was adjusted to be  $2.0 \times 10^5$  CFU/mL. (A) naringenin, (B) hesperetin, and (C) 7-*O*-butylnaringenin.



Values are mean with standard error. \*  $p < 0.05$  compared with control.

### 3.3. Urease Inhibition by Flavonoids in *H. pylori*

The urease of *H. pylori* is known to be one of the major factors for its resistance to the strong acidic conditions of the stomach [1,7]. *H. pylori* urease produces ammonia from urea and raises the pH of the stomach. Since urea is a small uncharged molecule, it can passively diffuse through the membrane of ureolytic bacteria [2,7]. Naringenin had little effect on the urease activity at  $<200 \mu\text{M}$ , while both hesperetin and 7-*O*-butylnaringenin had significant inhibitory effects on the urease activities of three *H. pylori* strains (Table 2). 7-*O*-Butylnaringenin has a stronger inhibitory effect on urease than the other two flavonoids, showing the highest inhibitory effect on *H. pylori* 26695, 51, and SS1 with the inhibition ratios of about 27%, 62%, and 71% at  $200 \mu\text{M}$  concentration, respectively.

Paulo *et al.* [30] demonstrated that resveratrol and red wines showed an inhibitory effect on *H. pylori* urease activity, which is considered a virulence factor of this organism. Masuda *et al.* [13] reported that gochoonangi (*Wasabia japonica*) has urease inhibitory activity. They showed that the inhibition rates of roots (44.6, 45.8, and 39.7%) and leaves (36.7, 40.2, and 48.7%) on urease activities

of *H. pylori* NCTC11637, YS27, and YS50, respectively. Although the anti-*H.pylori* activity of flavonoids has been described by several authors [2,5,6,13,14,16,19,20], there has been little study on the inhibition of urease activity on *H. pylori* strains.

**Table 2.** The inhibitory effect (%) of various flavonoids at 200  $\mu$ M on urease activity of *H. pylori* strains after incubation for 3 h at 37 °C.


Flavonoids	Naringenin	Hesperetin	7-O-Butylnaringenin
<i>H. pylori</i> 26695	1.33 $\pm$ 0.13 <sup>*,ax</sup>	0.93 $\pm$ 0.15 <sup>ax</sup>	27.49 $\pm$ 0.32 <sup>bx</sup>
<i>H. pylori</i> 51	16.55 $\pm$ 1.23 <sup>ax</sup>	7.31 $\pm$ 0.23 <sup>ax</sup>	61.93 $\pm$ 1.45 <sup>by</sup>
<i>H. pylori</i> SS1	13.64 $\pm$ 0.98 <sup>ax</sup>	6.32 $\pm$ 0.11 <sup>ax</sup>	70.75 $\pm$ 3.56 <sup>by</sup>

\* Inhibition effect (%) =  $(OD_{\text{control}} - OD_{\text{sample}}/OD_{\text{control}}) \times 100$ ; <sup>a-b</sup> Different letters within a row differ significantly ( $p < 0.05$ ); <sup>x-y</sup> Different letters within a column differ significantly ( $p < 0.05$ ).

### 3.4. Morphological Study of *H. pylori* Using SEM

The treatment of *H. pylori* with flavonoids at 200  $\mu$ M resulted in morphological alterations of the membrane compared with the control (Figure 3). *H. pylori* cells in the control treatment had regular shapes and smooth surfaces (Figure 3(A)), but the cells treated with hesperetin (Figure 3(C)) and 7-O-butyl naringenin (Figure 3(D)) at 200  $\mu$ M were damaged and had irregular shapes and rough surfaces.

**Figure 3.** SEM micrographs of *H. pylori* 26695 treated with naringenin, hesperetin, and 7-O-butyl naringenin at 200  $\mu$ M concentration. (A) control, (B) naringenin, (C) hesperetin, and (D) 7-O-butyl naringenin.



In addition, hesperetin and 7-*O*-butylnaringenin generated different cell damage patterns. Although the morphological changes in 7-*O*-butylnaringenin-treated cells were less pronounced than those with hesperetin-treated cells, 7-*O*-butylnaringenin reduced the viable cell count more and had a stronger inhibitory effect on urease activity than hesperetin (Figure 2 and Table 2). These results indicated that the mechanisms of cell damages may be quite different depending on the kind of flavonoids. On the other hand, naringenin at the same concentration did not cause detectable cell damages (Figure 3(B)).

The paper disc diffusion results indicated that the cells incubated with hesperetin and 7-*O*-butylnaringenin showed severe membrane damages with membrane integrity disruptions (Table 1). The initial exposure of hesperetin and 7-*O*-butylnaringenin to *H. pylori* caused large surface collapses in the cells and produced wrinkled abnormalities in the cells of *H. pylori* with numerous small clefts, regularly distributed on the bacterial cell surface. Such morphological features in bacterial cells may be due to the lysis of the outer membrane followed by the loss of cellular electron-dense materials on the surface of treated cells [31,32]. Although the mechanisms by which the effects observed in this study are not yet known, it likely involves with the perturbation or inhibition of metabolism such as nucleic acid synthesis, cytoplasmic membrane function, and energy metabolism [22].

#### 4. Conclusions

In conclusion, the greater inhibitory effect of 7-*O*-butylnaringenin compared with natural naringenin may be helpful in investigating the antimicrobial effects of other flavonoids against *H. pylori*, and in the development of new drugs for patients with gastric diseases caused by *H. pylori*. In addition, 7-*O*-butylnaringenin can be chemically synthesized from orange or tangerine waste, increasing its industrial value as an antimicrobial agent.

#### Acknowledgments

This work was supported by Priority Research Centers Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2012-0006686).

#### Conflicts of Interest

The authors declare no conflict of interest.

#### References

1. Marshall, B.J. *Campylobacter pyloridis* and gastritis. *J. Infect. Dis.* **1986**, *153*, 650–657.
2. Fukai, T.; Marumo, A.; Kaitou, K.; Kanda, T.; Terada, S.; Nomura, T. Anti-*Helicobacter pylori* flavonoids from licorice extract. *Life Sci.* **2002**, *71*, 1449–1463.
3. Jiang, X.; Doyle, M.P. Optimizing enrichment culture conditions for detecting *Helicobacter pylori* in foods. *J. Food Prot.* **2002**, *65*, 1949–1954.
4. Smith, J.L. The role of gastric acid in preventing foodborne disease and how bacteria overcome acid conditions. *J. Food Prot.* **2003**, *66*, 1292–1303.

5. Stamatis, G.; Kyriazopoulos, P.; Golegou, S.; Basayiannis, A.; Skaltsas, S.; Skaltsa, H. *In vitro* anti-*Helicobacter pylori* activity of Greek herbal medicines. *J. Ethnopharmacol.* **2003**, *88*, 175–179.
6. Shin, J.E.; Kim, J.M.; Bae, E.A.; Hyun, Y.J.; Kim, D.H. *In vitro* inhibitory effects of flavonoids on growth, infection and vacuolation of *Helicobacter pylori*. *Planta Med.* **2005**, *71*, 197–201.
7. Stingel, K.; Reuse, H.D. Staying alive overdose: How does *Helicobacter pylori* control urease activity? *Int. J. Med. Microbiol.* **2005**, *295*, 307–315.
8. Hazell, S.L.; Lee, A. *Campylobacter pyloridis*, urease, hydrogen ion back diffusion and gastric ulcers. *Lancet* **1986**, *5*, 15–17.
9. Hazell, S.L.; Borody, T.J.; Lee, A. *Campylobacter pyloridis* gastritis. I: Detection of urease as a marker of bacterial colonization and gastritis. *Am. J. Gastroenterol.* **1987**, *82*, 292–296.
10. Muhammad, A.; Farooq, A.; Muhammad, R.J.; Muhammad, A.I.; Umer, R. Green synthesis of silver nanoparticles through reduction with *Solanum xanthocarpum* L. Berry extract: Characterization, antimicrobial and urease inhibitory activities against *Helicobacter pylori*. *Int. J. Mol. Sci.* **2012**, *13*, 9923–9941.
11. Jiang, X.; Doyle, M.P. Effect of environmental and substrate factors on survival and growth of *Helicobacter pylori*. *J. Food Prot.* **1998**, *61*, 929–933.
12. Nicoline, F.T.; Lucy, M.N.; Roland, N.N. DNA sequence analysis of South African *Helicobacter pylori* vacuolating cytotoxin gene (*vacA*). *Int. J. Mol. Sci.* **2011**, *12*, 7459–7468.
13. Masuda, H.; Naohide, K.; Woo, G.J.; Shin, I.S. Inhibitory effects of gochoonangi (*Wasabia japonica*) against *Helicobacter pylori* and its urease activity. *Food Sci. Biotechnol.* **2004**, *13*, 191–196.
14. Silvia, M.; Claudia, D.A.; Andrea, C.; Silvia, F.; Francesca, B.; Annalisa, S.; Natale, F.; Claudio, R. Antimicrobial activity against *Helicobacter pylori* strains and antioxidant properties of blackberry leaves (*Rubus ulmifolius*) and isolated compounds. *Int. J. Antimicrob. Agric.* **2009**, *34*, 50–59.
15. Muhammad, A.; Farooq, A.; Fauqia, N.; Tahir, M.; Nazamid, S. Anti-*Helicobacter pylori* and urease inhibition activities of some traditional medicinal plants. *Molecules* **2013**, *18*, 2135–2149.
16. Stojanovic, G.; Radulovic, N.; Hashimoto, T.; Palic, R. *In vitro* antimicrobial activity of extracts of four *Achillea* species: The composition of *Achillea clavennae* L. (Asteraceae) extract. *J. Ethnopharmacol.* **2005**, *101*, 185–190.
17. Maria, H.; Concepción, S.-M.; Sonia, P.T. Flavonoid-flavonoid interaction and its effect on their antioxidant activity. *Food Chem.* **2010**, *121*, 691–696.
18. Bang, M.H.; Jung, Y.J.; Kim, W.K. Effect of several flavonoids on cancer cell motility in human fibrosarcoma HT 1080 cells. *Food Sci. Biotechnol.* **2004**, *13*, 739–743.
19. Kim, K.T.; Moon, S.H.; Yeo, E.J.; Han, Y.S.; Nah, S.Y.; Paik, H.-D. Inhibitory effect of 7-*O*-butyl naringenin on growth of *Helicobacter pylori* ATCC 26695. *Food Sci. Biotechnol.* **2006**, *15*, 466–468.
20. Ghaleb, A.; Bassam, A.-S.; Kamel, A. Antibacterial activities of some plant extracts alone and in combination with different antimicrobials against multidrug-resistant *Pseudomonas aeruginosa* strains. *Asian Pac. J. Trop. Med.* **2010**, *3*, 266–269.

21. Son, D.J.; Lee, S.E.; Park, B.S. Inhibitory effects of naturally occurring flavonoids on a human intestinal bacterium, *Clostridium perfringens*. *Food Sci. Biotechnol.* **2003**, *12*, 180–182.
22. Tim, C.T.P.; Lamb, A.J. Antimicrobial activity of flavonoids. *Int. J. Antimicrob. Agric.* **2005**, *26*, 343–356.
23. Tim, C.T.P.; Hamilton, V.E.S.; Lamb, A.J. Assessment of the antibacterial activity of selected flavonoids and consideration of discrepancies between previous reports. *Microbial. Res.* **2003**, *158*, 281–289.
24. Christy, E.M.L.; Clarke, A.M.; Munzhelele, T.; Ezekiel, G.; Noxolo, F.M.; Roland, N.N. Selected South African honeys and their extracts possess *in vitro* anti-*Helicobacter pylori* activity. *Arch. Med. Res.* **2010**, *41*, 324–331.
25. Yerra, K.R.; Lien, H.-M.; Lin, Y.H.; Hsu, Y.-M.; Yeh, C.-T.; Chen, C.C.; Lai, C.H.; Tzeng, Y.M. Antibacterial activities of *Anisomeles indica* constituents and their inhibition effect on *Helicobacter pylori*-induced inflammation in human gastric epithelial cells. *Food Chem.* **2012**, *132*, 780–787.
26. Kim, K.T.; Paik, H.-D. Propolis as a Multi-Functional Natural Material for Health. In *Natural Products as Future Medicinal Agents*; Hawthorne, S., Ed.; Transworld Research Network: Kerala, India, **2008**; pp. 33–48.
27. Malekzadeh, F.; Ehsanifar, H.; Shahamat, M.; Levin, M.; Colwell, R.R. Antibacterial activity of black myrobalan (*Terminalia chebula*) against *Helicobacter pylori*. *Int. J. Antimicrob. Agric.* **2001**, *18*, 85–88.
28. Mobley, H.L.T.; Cortesia, M.J.; Roseuthal, L.E.; Jones, B.D. Characterization of urease from *Campylobacter pylori*. *J. Clin. Microbiol.* **1988**, *26*, 831–836.
29. Tsugawa, H.; Suzuki, H.; Nakagawa, I.; Nishizawa, T.; Saito, Y.; Suematsu, M.; Hibi, T. Alpha-ketoglutarate oxidoreductase, an essential salvage enzyme of energy metabolism, in coccoid form of *Helicobacter pylori*. *Biochem. Biophys. Res. Commun.* **2008**, *376*, 46–51.
30. Paulo, L.; Oleastro, M.; Gallardo, E.; Queiroz, J.A.; Domi, F. Anti-*Helicobacter pylori* and urease inhibitory activities of resveratrol and red wine. *Food Res. Int.* **2011**, *44*, 964–969.
31. Becker-Ritta, A.B.; Martinelli, A.H.S.; Mitidierib, S.; Federa, V.; Wassermann, G.E.; Santia, L.; Vainsteina, M.H.; Oliveirad, J.T.A.; Fiuzae, L.M.; Pasqualia, G.; *et al.* Antifungal activity of plant and bacterial ureases. *Toxicon* **2007**, *50*, 971–983.
32. Shin, S.Y.; Bajpai, V.K.; Kim, H.R.; Kang, S.C. Antibacterial activity of eicosapentaenoic acid (EPA) against foodborne and food spoilage microorganisms. *LWT-Food Sci. Technol.* **2007**, *40*, 1515–1519.