

10-2003

The Role of Dietary Vitamin E in Experimental *Listeria monocytogenes* Infections in Turkeys

M. Zhu

United States Department of Agriculture

I. V. Wesley

United States Department of Agriculture

R. Nannapaneni

University of Arkansas at Fayetteville

M. Cox

University of Arkansas at Fayetteville

Aubrey F. Mendonca

Iowa State University, amendon@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/ans_pubs



Part of the [Agriculture Commons](#), [Meat Science Commons](#), and the [Poultry or Avian Science Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/ans_pubs/66. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

The Role of Dietary Vitamin E in Experimental *Listeria monocytogenes* Infections in Turkeys

Abstract

The current study was designed to determine if dietary vitamin E influenced either the gut clearance or levels of peripheral blood CD4+ and CD8+ T lymphocytes in adult turkeys experimentally infected with *Listeria monocytogenes*. Turkeys were fed vitamin E (0, 100, or 200 IU) from day of hatch to time of necropsy. After 6 wk on the experimental diet, turkeys were orally inoculated with *L. monocytogenes* (~ 10⁹ cfu). To monitor infection status, cloacal swabs were taken on selected days post-inoculation (DPI). At necropsy, samples of viscera, including liver, spleen, cecum, duodenum, ileum, and colon were collected and cultured for *L. monocytogenes*. In experiments 1 and 2, recovery of *L. monocytogenes* from cloacal swabs, tissues, and intestines from turkeys fed vitamin E was generally lower than that from turkeys fed the control diet, although these differences were not statistically significant. When data from both trials were combined, *L. monocytogenes* was cultured less frequently from cloacal swabs of the vitamin E-treated group (200 IU) on 2 and 3 DPI, when compared to controls (0 IU, $P < 0.01$). There were no changes in virulence characteristics of *L. monocytogenes* cells, as measured by in vitro killing of Ped-2E9 cells, recovered from cloacal swabs or tissues of experimentally infected turkeys fed the control or a vitamin E treatment diet. Flow cytometric analysis indicated that CD4+ and CD8+ peripheral blood T lymphocytes were elevated at 6 and 8 DPI in infected turkeys given 200 IU vitamin E.

Keywords

CD4+ and CD8+, immune response, *Listeria*, turkey, vitamin E

Disciplines

Agriculture | Animal Sciences | Meat Science | Poultry or Avian Science

Comments

This article is from *Poultry Science* 82 (2003): 1559, doi:[10.1093/ps/82.10.1559](https://doi.org/10.1093/ps/82.10.1559).

Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

Authors

M. Zhu, I. V. Wesley, R. Nannapaneni, M. Cox, Aubrey F. Mendonca, M. G. Johnson, and Dong U. Ahn

The Role of Dietary Vitamin E in Experimental *Listeria monocytogenes* Infections in Turkeys¹

M. Zhu,^{*†} I. V. Wesley,^{*2} R. Nannapaneni,[‡] M. Cox,[‡] A. Mendonca,[§] M. G. Johnson,[‡] and D. U. Ahn[§]

^{*}Pre-Harvest Food Safety and Enteric Diseases Research Unit, National Animal Disease Center, USDA, Agricultural Research Service, Ames, Iowa 50010; [†]Department of Food Science and Human Nutrition, Iowa State University, Ames, Iowa 50011; [‡]Department of Food Science and Center for Food Safety and Quality, IFSE, University of Arkansas, Fayetteville, Arkansas 72704; and [§]Department of Animal Science, Iowa State University, Ames, Iowa 50011

ABSTRACT The current study was designed to determine if dietary vitamin E influenced either the gut clearance or levels of peripheral blood CD4⁺ and CD8⁺ T lymphocytes in adult turkeys experimentally infected with *Listeria monocytogenes*. Turkeys were fed vitamin E (0, 100, or 200 IU) from day of hatch to time of necropsy. After 6 wk on the experimental diet, turkeys were orally inoculated with *L. monocytogenes* (~ 10⁹ cfu). To monitor infection status, cloacal swabs were taken on selected days post-inoculation (DPI). At necropsy, samples of viscera, including liver, spleen, cecum, duodenum, ileum, and colon were collected and cultured for *L. monocytogenes*. In experiments 1 and 2, recovery of *L. monocytogenes* from cloacal swabs, tissues, and intestines from turkeys fed

vitamin E was generally lower than that from turkeys fed the control diet, although these differences were not statistically significant. When data from both trials were combined, *L. monocytogenes* was cultured less frequently from cloacal swabs of the vitamin E-treated group (200 IU) on 2 and 3 DPI, when compared to controls (0 IU, *P* < 0.01). There were no changes in virulence characteristics of *L. monocytogenes* cells, as measured by in vitro killing of Ped-2E9 cells, recovered from cloacal swabs or tissues of experimentally infected turkeys fed the control or a vitamin E treatment diet. Flow cytometric analysis indicated that CD4⁺ and CD8⁺ peripheral blood T lymphocytes were elevated at 6 and 8 DPI in infected turkeys given 200 IU vitamin E.

(Key words: CD4⁺ and CD8⁺, immune response, *Listeria*, turkey, vitamin E)

2003 Poultry Science 82:1559–1564

INTRODUCTION

Listeria monocytogenes is a major human bacterial food-borne pathogen that annually accounts for ~2,500 cases (meningitis, encephalitis, sepsis, fetal death, prematurity) and 504 deaths (Mead et al., 1999). Sporadic human cases of listeriosis have been epidemiologically linked to the consumption of undercooked poultry products (Schwartz et al., 1988). Analysis of risk factors associated with sporadic human listeriosis in the U.S. indicated that cancer patients and immunocompromised patients, in whom 69% of listeriosis cases occur, were more likely than controls to have eaten undercooked poultry (odds ratio = 3.3; Schuchat et al., 1992). A 1998 multistate outbreak of human listeriosis, ascribed to serotype 4 (101 cases,

resulting in 22 deaths), was linked to delicatessen meats, including turkey (MMWR, 1998). A 2002 outbreak involving 46 cases, seven deaths, and three stillbirths was linked to contaminated delicatessen turkey meats (MMWR, 2002). The recall of 26 million pounds of turkey meat in 2002 indicates the economic consequences of ready-to-eat meats contaminated with *L. monocytogenes* (U.S. Department of Health and Human Services, 2002).

Adult turkeys may be transiently colonized by consuming contaminated feed or water (Husu et al., 1990). Thus, *L. monocytogenes* may enter the packing plant at low levels in the intestine of recently infected birds, survive in biofilms, and ultimately contribute to both environmental and ready-to-eat product contamination (Genigeorgis et al., 1990; Ojeniyi et al., 1996). In the United States, *L. monocytogenes* was found on 5.9% of turkey carcass rinses and in 31% of ground turkey meat examined in the Nationwide Young Turkey Microbiological Baseline Data Collection Program (U.S. Department of Agriculture, 1998).

Vitamin E is required for normal development and function of the immune system in poultry (Boa-Am-

©2003 Poultry Science Association, Inc.
Received for publication January 2, 2003.
Accepted for publication May 21, 2003.

¹Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture.

²To whom correspondence should be addressed: iwesley@nadc.ars.usda.gov.

Abbreviation Key: DPI = days post-inoculation.

ponsem et al., 2000; Leshchinsky and Klasing, 2000, 2001). In chickens, vitamin E supplement increased the number of lymphocytes in the bursa and the thymus gland and stimulated the proliferation and differentiation of T cells (Chang et al., 1994). In broilers, vitamin E selectively increased the percentage of mature CD4⁺ T helper cells in the thymus and spleen but did not alter the percentage of thymic and splenic B cells and macrophages in total immune cell (Gore and Qureshi, 1997; Erf et al., 1998). Vitamin E enhanced immunity of birds to *Escherichia coli* infection, coccidiosis, infectious bursal disease, and Newcastle disease and altered cytokine expression in broilers (Tengerdy and Brown, 1977; Colnago et al., 1984; Erf et al., 1998; Leshchinsky and Klasing, 2000). However, vitamin E, although increasing serum levels of α -tocopherol, did not reduce the severity of *Eimeria maxima* infections in broilers (Allen and Fretterer, 2002).

Besides stimulating immune parameters, dietary vitamin E also enhances meat quality. Dietary vitamin E contributes to oxidative stability, extends shelf life and prevents oxidative off-odor of poultry meat, thus preserves the sensory quality of both frozen and refrigerated turkey breast meat (Ahn et al., 1997, 1998; Sheldon et al., 1997).

As part of the immune response during acute listeriosis, the host marshalls neutrophils, macrophages, natural killer cells, and T lymphocytes, especially CD4⁺ and CD8⁺ (Unanue, 2002). In vitro transfer experiments have shown that CD4⁺ and CD8⁺ cells are required to eliminate infection with wild-type strains of *L. monocytogenes* in mice (Kaufmann, 1993).

The current study was designed primarily to assess the effectiveness of dietary vitamin E, using doses previously shown to enhance meat quality, in accelerating the gut clearance of *L. monocytogenes* in experimentally infected adult turkeys. Secondarily, we monitored CD4⁺ and CD8⁺ T lymphocytes to evaluate the role of vitamin E as an immune potentiator.

MATERIALS AND METHODS

Bacterial Inoculum

L. monocytogenes (ATCC 700301) was obtained from American Type Culture Collection.³ The stock cultures were maintained (−70°C) in 50% glycerol. For experimental inoculations, cultures of *L. monocytogenes* were grown on brain heart infusion agar with 20% bovine blood and 0.5% yeast extract (30°C, 24 h, in 5% O₂, 10% CO₂, 85% N₂) and harvested in PBS buffer (5 mL/plate). After centrifugation (5,000 × g, 10 min, 4°C), the pellet was washed twice with PBS and resuspended in 20 mL PBS. A 1-mL aliquot of the suspension was serially diluted in PBS. *L. monocytogenes* (cfu) were enumerated after incubation of

brain heart infusion agar plates seeded with 0.1 mL of each serial 10-fold dilution (30°C for 24 h, in 5% O₂, 10% CO₂, 85% N₂).

Dietary Vitamin E

DL- α -Tocopherol acetate (500 IU/g Rovimix E-50%),⁴ was used in the corn-soybean meal-based diet formulation, as described by Nam et al. (2003).

Turkeys

Experiment 1. One-day-old mixed sex Large White turkeys (n = 90) were obtained from a local hatchery and allotted to six rooms. Two rooms each (30 turkeys total) were randomly assigned to one of the three dietary treatments containing 0, 100, or 200 IU vitamin E/kg feed. After 5 wk, cloacal swabs were taken with sterile cotton-tipped applicators⁵ to ensure that birds were culture-negative for *Listeria*. Any turkeys positive for *Listeria* were eliminated prior to experimental inoculation. No attempt was made to select birds, which were innately resistant to *Listeria*. One week later, only *Listeria*-negative turkeys were orally challenged with 1 mL of *L. monocytogenes* (1 × 10⁹ cfu/mL). To monitor infection status, cloacal swabs were taken at 1, 4, 5, and 6 d post-inoculation (DPI). Turkeys (4 to 5 per group) were necropsied at 5, 8, 11, 14, and 25 DPI. Liver, spleen, cecum, duodenum, ileum, and colon from each bird were collected and processed for *Listeria* isolation as described below.

Experiment 2. One-day-old mixed sex Large White turkeys (n = 70) were obtained from a local hatchery and allotted to 4 rooms. Two rooms (35 turkeys total) were randomly assigned to diets containing either 0 or 200 IU vitamin E/kg feed. Prior to infection at 5 wk, cloacal swabs were taken and cultured for *Listeria*. At 6 wk, 30 *Listeria*-negative turkeys in each diet group were orally challenged with 1 mL of *L. monocytogenes* (1 × 10⁹ cfu/mL). The remaining five *Listeria*-negative turkeys in each diet group were moved to the clean pens and served as non-infected controls. To monitor infection status, cloacal swabs were taken at 1, 2, 3, 4, 6, 8, and 10 DPI. Turkeys (~ 5 per group) were necropsied at 2, 4, 6, 8, and 10 DPI. Liver, spleen, cecum, duodenum, ileum, and colon from each bird were collected and processed for *Listeria* isolation as described below.

Bacterial Isolation

To culture *Listeria*, cloacal swabs were placed in UVM I (10 mL)⁶ and incubated (2 to 3 d, 30°C, in 5% O₂, 10% CO₂, and 85% N₂). After enrichment, 100- μ L UVM I enrichment was transferred into 10-mL *Listeria* secondary enrichment broth (UVM II) and incubated (30°C, in 5% O₂, 10% CO₂, and 85% N₂). After ~48 h, 100 μ L of UVM II was plated to PALCAM *L. monocytogenes* selective agar⁶ (30°C, 5% O₂, 10% CO₂, and 85% N₂ for 48 h). At necropsy, the liver, spleen, cecum, duodenum, ileum, and colon from each bird were sampled, enriched in UVM I (10% wt/vol), and cultured as described above.

³American Type Culture Collection, Manassas, VA.

⁴Roche Vitamins, Inc., Ames, IA.

⁵Puritan Hardwood Products, Guilford, ME.

⁶Oxoid Ltd., Basingstoke, Hampshire, UK.

Two presumptive *Listeria* spp. colonies were recovered from each PALCAM agar plate and were verified as *L. monocytogenes* by a multiplex PCR assay as described (Wesley et al., 2002). A total of 245 isolates from cloacal swabs and viscera were stored (4°C) on tryptic soy agar slants supplemented with 0.6% yeast extract and assayed for virulence.

Virulence Assay for *L. Monocytogenes* Isolates

Single microcolonies from each of 245 *Listeria* isolates recovered from infected turkeys were tested in vitro for virulence, as described (Bhunja et al., 1994, 1995). The ratio of *L. monocytogenes* cells per each target hybridoma cell was approximately 1,000:1. Tissue culture plates were incubated (37°C and 7% CO₂) for 6 h prior to scoring Ped-2E9 cell death using a trypan blue⁷ exclusion assay.

Percent hybridoma cell death at 6 h after microcolony challenge was calculated as follows: $[(L_{NC} - L_{LC}) / L_{NC}] \times 100$, where L_{NC} = number of total live Ped-2E9 cells observed in untreated or unchallenged control wells, and L_{LC} = number of live Ped-2E9 cells in wells challenged with *Listeria* strain. Isolates that killed >70 to 90% Ped-2E9 cells within 6 h of challenge, which was comparable to the reference strains of *L. monocytogenes*, were scored as highly virulent. Isolates that killed <10% of Ped-2E9 cells, which was comparable to the 8 to 10.6% cell death observed for *L. innocua* (negative control), were scored as avirulent.

Serum Vitamin E (α -Tocopherol) Analysis

Blood samples (10 mL) were collected in serum separation vacutainer tubes.⁷ Serum vitamin E analyses were performed with a Hewlett Packard (HP) 6890 GC equipped with an on-column capillary injector and a FID detector.⁸ Serum vitamin E was calculated using an internal standard, 5 α -cholestane, and expressed as micrograms per milliliter (Du and Ahn, 2002).

Flow Cytometric Analysis of Lymphocyte Population

Five milliliters of blood was collected from the wing vein into a vacutainer tube containing sodium heparin.⁷ The heparinized whole blood was transferred into a 15-mL conical centrifuge tube containing 5 mL of fluorescence buffer (FB is PBS containing 1% heat-inactivated fetal blood serum and 0.05% NaN₃). The contents were mixed and centrifuged (200 g for 15 min, 4°C). The buffy coat was collected, washed three times with FB, and resuspended in 0.5 mL FB, as described (Stabel et al., 2000).

TABLE 1. Serum vitamin E levels (μ g/mL) in turkeys fed 0, 100, 200 IU of vitamin E (experiments 1 and 2)

Days post-infection	0 IU	100 IU	200 IU
Experiment 1			
5	0.11 \pm 0.29 ^c	2.08 \pm 0.33 ^b	2.88 \pm 0.29 ^a
8	0.16 \pm 0.20 ^c	1.51 \pm 0.20 ^b	3.10 \pm 0.20 ^a
11	0.09 \pm 0.22 ^c	1.75 \pm 0.22 ^b	3.88 \pm 0.22 ^a
14	0.15 \pm 0.21 ^c	1.89 \pm 0.21 ^b	3.67 \pm 0.22 ^a
Experiment 2			
0	0.10 \pm 0.09 ^b		3.42 \pm 0.24 ^a
2	0.10 \pm 0.08 ^b		3.31 \pm 0.39 ^a
4	0.12 \pm 0.04 ^b		3.19 \pm 0.29 ^a
6	0.09 \pm 0.06 ^b		3.41 \pm 0.35 ^a

^{a-c}Means data in same row with significant difference ($P \leq 0.05$). Four to five birds were analyzed at each sampling point for each dietary regimen.

Direct dual color immunofluorescence staining was performed as described previously (Stabel et al., 2000). Briefly, 50 μ L of buffy coat ($\sim 1 \times 10^6$ viable cells) was incubated (20 min at room temperature) with 10 μ L each of 1:20 diluted CT4-FITC (fluorescein isothiocyanate conjugate mouse anti-chicken CD4 monoclonal antibody; catalog number 8210-02)⁹ and CT8-PE (phycoerythrin mouse anti-chicken CD8a antibody; catalog number 8405-09).⁹ After fixation with 1.5% formalin, the fluorescence intensities were measured with a Becton-Dickson FACScan flow cytometer.¹⁰ Cells incubated with the fluorescently labeled isotype served as controls.

Statistical Analysis

A completely randomized design with unequal numbers of subjects was used to examine the effects of *L. monocytogenes* recovery from cloacal swabs of infected turkeys on vitamin E supplemented (200 IU) and the control (0 IU) diets. Regression equations were fit for percentage of *L. monocytogenes* recovery as a function of DPI. A general linear models F-test for full and reduced models was used for comparison. If a significant test statistic was found, indicating unequal swab responses of the two diet groups, 95% confidence intervals were calculated for each group on each day to highlight where differences were occurring (Neter et al., 1990). The mean and standard deviations of serum vitamin E as well as immune cells were analyzed statistically by the general linear models procedure using SAS software.¹¹ Student-Newman-Keuls' multiple range test was used to compare differences among mean values ($P < 0.05$). Means and SEM are reported.

RESULTS

Serum α -Tocopherol (Vitamin E)

Dietary vitamin E resulted in an increase in serum α -tocopherol (vitamin E) for experiments 1 and 2 (Table 1). At the time of experimental inoculation with *L. monocytogenes* (wk 6), serum vitamin E levels in both the 100 IU and 200 IU vitamin E treatment groups were significantly

⁷Becton Dickinson, Franklin Lakes, NJ.

⁸Hewlett Packard Co., Wilmington, DE.

⁹Southern Biotechnology Associates, Inc., Birmingham, AL.

¹⁰Becton Dickinson Co., Cockeysville, MD.

¹¹SAS Institute, Cary, NC.




FIGURE 1. Comparison of the recovery of *Listeria monocytogenes* from cloacal swabs of turkeys fed control (0 IU) or a vitamin E (200 IU) diet. Significantly fewer turkeys harbored *L. monocytogenes* on d 2 and 3 post-infection (DPI) when fed vitamin E (200 IU). An asterisk (*) indicates a statistically significant difference between 0 IU (curve a) and 200 IU vitamin E (curve b) treatments at 2 and 3 DPI.

elevated when compared to birds on the control diet (0 IU). For both trials 1 and 2, serum vitamin E titers were significantly elevated in birds receiving 200 IU when compared to controls (0 IU). Because serum vitamin E levels, at 8, 11, and 14 DPI, were consistently higher in turkeys fed 200 IU than that in turkeys receiving 100 IU vitamin E, only the 200 IU diet was evaluated in experiment 2.

Recovery of *L. monocytogenes* in Cloacal Swabs

In experiments 1 and 2, pre-inoculation cloacal swabs were negative for *L. monocytogenes*. In experiment 1 at 1 DPI, *L. monocytogenes* was detected in cloacal swabs of turkeys (22/24, 91.7%) fed the control diet (0 IU vitamin E), as well as birds fed 100 IU (21/24, 87.5%) and 200 IU (17/25, 68%) vitamin E. At 4 DPI, *L. monocytogenes* was

detected in swabs of 8.3% of control diet turkeys (2/24), whereas only one bird each was positive in the 100 IU (1/24, 4.2 %) and 200 IU (1/25, 4 %) treatment groups. On d 5 PI, three control birds (3/24, 12.5%) were positive, in contrast to none of the turkeys in the 100 IU and 200 IU groups. In experiment 2, at all days of post-infection sampling, *L. monocytogenes* was recovered more frequently from cloacal swabs of turkeys fed the control diet (0 IU) than in turkeys fed 200 IU vitamin E. When data from the two trials were combined (Figure 1), *L. monocytogenes* was cultured more frequently ($P < 0.01$) on d 2 and 3 in birds receiving 200 IU vitamin E when compared to control birds (0 IU).

Recovery of *L. monocytogenes* in Tissues

In experiment 1, *L. monocytogenes* was recovered more often in the ceca and ileum of control diet turkeys versus vitamin E-treated birds at 5 DPI. *L. monocytogenes* was not recovered from the intestine or tissue samples of turkeys on either control or vitamin E diets after 8 DPI (data not shown). For experiment 2, as summarized in Table 2, there were fewer tissue samples positive for *Listeria* in vitamin E-treated birds at 2, 4, or 6 DPI. At 8 DPI, *L. monocytogenes* was not recovered from any group (control and vitamin E).

Virulence Assays for *L. monocytogenes* Isolates

A total of 240 out of 245 isolates (98%) recovered from infected turkeys and submitted for virulence testing killed >70% to 90% of target hybridoma cells within 6 h of challenge. These were scored as highly virulent as were the inoculating strain (ATCC 700301) and the reference *L. monocytogenes* strains (positive controls). Of these 245 isolates, 156 were recovered from cloacal swabs and the remaining 89 were from intestinal tissues of turkeys. Four remaining isolates killed 3.8 to 12.8% of Ped-2E9 cells. This finding was comparable to the 8 to 10.6% cell death observed for *L. innocua* (negative control). These were scored as avirulent, and later were confirmed by PCR as isolates that were not *L. monocytogenes*.

TABLE 2. Recovery of *Listeria monocytogenes* from tissues of turkeys fed 0 and 200 IU of vitamin E (experiment 2)

Diet	Liver ¹	Spleen	Cecal	Small intestine		
				Duodenum	Ileum	Colon
Day 2						
0 IU	20%	0%	60%	40%	60%	40%
200 IU	40%	0%	60%	20%	20%	0%
Day 4						
0 IU	0%	0%	60%	20%	0%	0%
200 IU	0%	0%	60%	0%	0%	0%
Day 6						
0 IU	20%	0%	0%	0%	0%	40%
200 IU	0%	0%	0%	0%	0%	20%

¹Percentage of infected birds sampled for each day for each group, n = 5.

Flow Cytometric Analysis of Lymphocytes

As summarized in Figure 2, for experimentally infected birds, CD4⁺ populations of turkeys fed 200 IU vitamin E were increased ($P < 0.05$) at 6, 8, 10, and 31 DPI when compared to infected turkeys fed control diets (0 IU vitamin E) (Figure 2a). At 6 and 8 DPI, the CD8⁺ T lymphocytes were higher ($P < 0.05$) in infected turkeys given 200 IU vitamin E than in infected turkeys on control diets (0 IU vitamin E) (Figure 2b). CD4⁺CD8⁺ double positive lymphocytes of experimentally infected turkeys on the 200 IU vitamin E diet were also markedly elevated ($P < 0.05$) at 6 and 8 DPI when compared to infected birds on the control (0 IU vitamin E) diet (Figure 2c).

DISCUSSION

The impact of dietary vitamin E on both gut colonization as well as on CD4⁺ and CD8⁺ T lymphocyte populations was evaluated in turkeys experimentally infected with *L. monocytogenes*. The vitamin E doses used were those previously shown to improve meat quality (Ahn et al., 1997, 1998). Serum vitamin E increased proportionately with dietary vitamin E content. This reflects effective gut absorption of vitamin E by the time of experimental challenge at the sixth week of dietary treatment. When cloacal swab data for the two trials were combined, *L. monocytogenes* was cultured more frequently ($P < 0.01$) on d 2 and 3 in birds receiving 200 IU vitamin E when compared to control birds (0 IU). Vitamin E supplement was previously reported to have increased the resistance of mice to influenza virus as well as of chickens to Newcastle disease virus, and increased antibody production and phagocytosis in chickens infected with *E. coli* (Tengerdy and Brown, 1977; Franchini et al., 1991; Han et al., 2000).

No virulence differences of *L. monocytogenes* recovered from in birds receiving vitamin E were detected using the PedE9 assay. The inoculating strains (ATCC 700301) and nearly 99% of the isolates recovered from cloacal swabs as well as from tissues, including spleen, liver, and intestine, were pathogenic for Ped-2E9 hybridoma cells after 6 h of incubation.

In order to determine the role of dietary vitamin E on immune parameters, the lymphocytes of infected turkeys were analyzed. In this current study, dietary vitamin E (200 IU) was associated with elevation of CD4⁺ (6, 8, and 31 DPI), CD8⁺, as well as CD4⁺CD8⁺ T lymphocytes (6 and 8 DPI) in *Listeria*-infected turkeys, when compared with infected turkeys on control diets. This observation was in concert with the requirement of CD4⁺ and CD8⁺ T cells to eliminate listeriosis (Unanue, 2002).

ACKNOWLEDGMENTS

We acknowledge Norm Lyon, Gary Buck, and Roger Ezarski for providing care for the turkeys. We thank Chunyan Wang, Sharon Franklin, Angelia Pinkerton, Sandhya Boyapalle, Laura B. Byl, and Karla Fenton for




FIGURE 2. The response of CD4⁺, CD8⁺, and CD4⁺CD8⁺ populations, measured as percentage of lymphocytes, of turkeys fed 0 or 200 IU of vitamin E and experimentally infected with *Listeria monocytogenes*. The response was monitored for CD4⁺ (a), CD8⁺ (b), and CD4⁺CD8⁺ (c) lymphocytes. An asterisk (*) indicates a statistically significant difference between 0 IU and 200 IU vitamin E treatments. CD4⁺ populations of turkeys fed 200 IU vitamin E were higher at 6, 8, 10, and 31 d post-infection (DPI) when compared to infected turkeys fed a control diet (0 IU vitamin E) (b). At 6 and 8 DPI, the CD8⁺ lymphocytes were significantly higher in infected turkeys given 200 IU vitamin E than in infected turkeys on control diets (0 IU vitamin E). CD4⁺CD8⁺ double positive lymphocytes of experimentally infected turkeys on 200 IU vitamin E diet were also markedly elevated ($P < 0.05$) at 6 and 8 DPI when compared to infected birds on the control (0 IU vitamin E) diet.

processing samples and bacterial isolation. We thank Bruce Pesch, Randy Sacco, Thomas Stabel, and Jolita Januenaite for their valuable advice in flow cytometry. We are indebted to Deb Palmquist for her guidance in statistical analysis. The proofreading skills of Sandy Johnson are appreciated. This study was supported by a USDA, ARS, National Alliance for Food Safety grant.

REFERENCES

- Ahn, D. U., J. L. Sell, M. Jeffery, C. Jo, X. Chen, C. Wu, and J. I. Lee. 1997. Dietary vitamin E affects lipid oxidation and total volatiles of irradiated raw turkey meat. *J. Food Sci.* 62:954–958.
- Ahn, D. U., J. L. Sell, C. Jo, X. Chen, C. Wu, and J. I. Lee. 1998. Effects of dietary vitamin E supplementation on lipid oxidation and volatiles content of irradiated, cooked turkey meat patties with different packaging. *Poult. Sci* 77:912–920.
- Allen, P. C., and R. H. Fetterer. 2002. Interaction of dietary vitamin E with *Eimeria maxima* infections in chickens. *Poult. Sci* 81:41–48.
- Bhunia, A. K., P. J. Steele, D. G. Westbrook, T. P. Maloney, and M. G. Johnson. 1994. A six hour in vitro virulence assay for *Listeria monocytogenes* using myeloma and hybridoma cells from murine and human sources. *Microb. Pathog.* 16:99–110.
- Bhunia, A. K., D. G. Westbrook, R. Story, and M. G. Johnson. 1995. Frozen stored murine hybridoma cells can be used to determine the virulence of *Listeria monocytogenes*. *J. Clin. Microbiol.* 33:3349–3351.
- Boa-Amponsem, K., S. E. Price, M. Picard, A. Geraert, and P. B. Siegel. 2000. Vitamin E and immune responses of broiler pureline chickens. *Poult. Sci* 79:466–470.
- Chang, W. P., J. S. Hom, R. R. Dietert, G. F. Combs, and J. A. Marsh. 1994. Effect of dietary vitamin E and selenium deficiency on chicken splenocyte proliferation and cell surface marker expression. *Immunopharmacol. Immunotoxicol.* 16:203–223.
- Colnago, G. L., L. S. Jensen, and P. L. Long. 1984. Effect of selenium and vitamin E on the development of immunity to coccidiosis in chickens. *Poult. Sci* 63:1136–1143.
- Du, M., and D. U. Ahn. 2002. Simultaneous analysis of tocopherols, cholesterol and phytosterols by gas chromatography. *J. Food Sci.* 67:1696–1700.
- Erf, G. F., W. G. Bottje, T. K. Bersi, M. D. Headrick, and C. A. Fritts. 1998. Effects of dietary vitamin E on the immune system in broilers: altered proportions of CD4 T cells in the thymus and spleen. *Poult. Sci* 77:529–537.
- Franchini, A., M. Canti, G. Manfreda, S. Bertuzzi, G. Asdrubali, and C. Franciosi. 1991. Vitamin E as adjuvant in emulsified vaccine for chicks. *Poult. Sci* 70:1709–1715.
- Genigeorgis, C. A., P. Oanca, and D. Dutulescu. 1990. Prevalence of *Listeria* spp. in turkey meat at the supermarket and slaughterhouse level. *J. Food Prot.* 53:282–288.
- Gore, A. B., and M. A. Qureshi. 1997. Enhancement of humoral and cellular immunity by vitamin E after embryonic exposure. *Poult. Sci* 76:984–991.
- Han, S. N., D. Wu, W. K. Ha, A. Beharka, D. E. Smith, B. S. Bender, and S. N. Meydani. 2000. Vitamin E supplementation increases T helper 1 cytokine production in old mice infected with influenza virus. *Immunology* 100:487–493.
- Husu, J. R., J. T. Beery, E. Nurmi, and M. P. Doyle. 1990. Fate of *Listeria monocytogenes* in orally dosed chicks. *Int. J. Food Microbiol.* 11:259–269.
- Kaufmann, S. H. E. 1993. Immunity to intracellular bacteria. *Annu. Rev. Immunol.* 11:19–163.
- Leshchinsky, T. V., and K. C. Klasing. 2000. Vitamin E and leukocytic cytokine expression in broilers. *Poult. Sci* 79(Suppl. 1):37. (Abstr.)
- Leshchinsky, T. V., and K. C. Klasing. 2001. Relationship between the level of dietary vitamin E and the immune response of broiler chickens. *Poult. Sci* 80:1590–1599.
- Mead, P. S., L. Slutsker, V. Dietz, L. F. McCaig, J. S. Bresse, C. Shapiro, P. M. Griffin, and R. V. Tauxe. 1999. Food-related illness and death in the United States. *Emerg. Infect. Dis.* 5:607–625.
- MMWR. 1998. Multistate outbreak of listeriosis—United States. *Morb. Mortal. Wkly. Rep.* 47:1085–1086.
- MMWR. 2002. Outbreak of listeriosis—Northeastern United States, 2002. *Morb. Mortal. Wkly. Rep.* 51:950–951.
- Nam, K. C., B. E. R. Min, H. Yan, E. J. Lee, A. Mendonca, I. Wesley and D. U. Ahn. 2003. Effect of dietary vitamin E and irradiation on lipid oxidation, color, and volatiles of fresh and previously frozen turkey breast patties. *Meat Sci.* (in press)
- Neter, J., W. Wasserman, and M. H. Kutner. 1990. *Applied Linear Statistical Models*. 3rd ed. Richard D. Irwin, Homewood, IL.
- Ojeniyi, B., H. C. Wegener, N. E. Jensen, and M. Bisgaard. 1996. *Listeria monocytogenes* in poultry and poultry products: epidemiological investigations in seven Danish abattoirs. *J. Appl. Bacteriol.* 80:395–401.
- Schuchat, A., K. A. Deaver, J. D. Wenger, B. D. Plikaytis, L. Mascola, R. W. Pinner, A. L. Reingold, and C. V. Broome. 1992. Role of foods in sporadic listeriosis. I. Case-control study of dietary risk factors. The Listeria Study Group. *J. Am. Med. Assoc.* 267:2041–2045.
- Schwartz, B., C. A. Ciesielski, C. V. Broome, S. Gaventa, G. R. Brown, B. G. Gellin, A. W. Hightower, and L. Mascola. 1988. Association of sporadic listeriosis with consumption of uncooked hot dogs and undercooked chicken. *Lancet* 2:779–782.
- Sheldon, B. W., P. A. Curtis, P. L. Dawson, and P. R. Ferket. 1997. Effect of dietary vitamin E on the oxidative stability, flavor, color, and volatile profiles of refrigerated and frozen turkey breast meat. *Poult. Sci* 76:634–641.
- Stabel, T. J., S. R. Bolin, B. A. Pesch, and T. E. Rahner. 2000. A simple and rapid flow cytometric method for detection of porcine cell surface markers. *J. Immunol. Methods.* 245:147–152.
- Tengerdy, R. P., and J. C. Brown. 1977. Effect of vitamin E and A on humoral immunity and phagocytosis in *E. coli* infected chicken. *Poult. Sci* 56:957–963.
- U.S. Department of Agriculture—Food Safety and Inspection Service. 1998. Subject: Nationwide young turkey microbiological baseline data collection program. <http://www.fsis.usda.gov/ophs/baseline/youngturk.pdf>. Accessed May 2003.
- U.S. Department of Health and Human Services. 2002. Update: Listeriosis outbreak investigation. <http://www.cdc.gov/od/oc/media/pressrel/r021104.htm>. Accessed May 2003.
- Unanue, E. R. 2002. Innate immunity in bacterial infections. Pages 93–103 in *Immunology of Infectious Diseases*. S. H. E. Kaufmann, A. Sher, and R. Ahmed, ed. ASM Press, Washington, D.C.
- Wesley, I. V., K. M. Harmon, J. S. Dickson, and A. R. Schwartz. 2002. Application of multiplex polymerase chain reaction (PCR) assay for the simultaneous confirmation of *Listeria* species and *Listeria monocytogenes*. *J. Food Prot.* 65:780–785.