

7-6-2016

# On the linear convergence of the alternating direction method of multipliers


Mingyi Hong

*Iowa State University*, [mingyi@iastate.edu](mailto:mingyi@iastate.edu)

Zhi-Quan Luo

*University of Minnesota - Twin Cities*

Follow this and additional works at: [http://lib.dr.iastate.edu/imse\\_pubs](http://lib.dr.iastate.edu/imse_pubs)

 Part of the [Industrial Engineering Commons](#), [Systems Engineering Commons](#), and the [Theory and Algorithms Commons](#)

The complete bibliographic information for this item can be found at [http://lib.dr.iastate.edu/imse\\_pubs/89](http://lib.dr.iastate.edu/imse_pubs/89). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

This Article is brought to you for free and open access by the Industrial and Manufacturing Systems Engineering at Iowa State University Digital Repository. It has been accepted for inclusion in Industrial and Manufacturing Systems Engineering Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact [digirep@iastate.edu](mailto:digirep@iastate.edu).

# On the Linear Convergence of the Alternating Direction Method of Multipliers\*

Mingyi Hong and Zhi-Quan Luo<sup>†</sup>

*Dedicated to the fond memories of a close friend and collaborator, Paul Y. Tseng*

August 13, 2012; Revised March 20, 2013

## Abstract

We analyze the convergence rate of the alternating direction method of multipliers (ADMM) for minimizing the sum of two or more nonsmooth convex separable functions subject to linear constraints. Previous analysis of the ADMM typically assumes that the objective function is the sum of only *two* convex functions defined on *two* separable blocks of variables even though the algorithm works well in numerical experiments for three or more blocks. Moreover, there has been no rate of convergence analysis for the ADMM without strong convexity in the objective function. In this paper we establish the global linear convergence of the ADMM for minimizing the sum of *any* number of convex separable functions. This result settles a key question regarding the convergence of the ADMM when the number of blocks is more than two or if the strong convexity is absent. It also implies the linear convergence of the ADMM for several contemporary applications including LASSO, Group LASSO and Sparse Group LASSO without any strong convexity assumption. Our proof is based on estimating the distance from a dual feasible solution to the optimal dual solution set by the norm of a certain proximal residual, and by requiring the dual stepsize to be sufficiently small.

**KEY WORDS:** Linear convergence, alternating directions of multipliers, error bound, dual ascent.

**AMS(MOS) Subject Classifications:** 49, 90.

---

\*The research is supported by the National Science Foundation, grant number DMS-1015346.

<sup>†</sup>Department of Electrical and Computer Engineering, University of Minnesota, Minneapolis, MN 55455, USA.

Email: {mhong, luozq}@umn.edu

# 1 Introduction

Consider the problem of minimizing a separable nonsmooth convex function subject to linear equality constraints:

$$\begin{aligned}
 & \text{minimize} && f(x) = f_1(x_1) + f_2(x_2) + \cdots + f_K(x_K) \\
 & \text{subject to} && Ex = E_1x_1 + E_2x_2 + \cdots + E_Kx_K = q \\
 & && x_k \in X_k, \quad k = 1, 2, \dots, K,
 \end{aligned} \tag{1.1}$$

where each  $f_k$  is a nonsmooth convex function (possibly with extended values),  $x = (x_1^T, \dots, x_K^T)^T \in \mathfrak{R}^n$  is a partition of the optimization variable  $x$ ,  $X = \prod_{k=1}^K X_k$  is the feasible set for  $x$ , and  $E = (E_1, E_2, \dots, E_K) \in \mathfrak{R}^{m \times n}$  is an appropriate partition of matrix  $E$  (consistent with the partition of  $x$ ) and  $q \in \mathfrak{R}^m$  is a vector. Notice that the model (1.1) can easily accommodate general linear inequality constraints  $Ex \geq q$  by adding one extra block. In particular, we can introduce a slack variable  $x_{K+1} \geq 0$  and rewrite the inequality constraint as  $Ex - x_{K+1} = q$ . The constraint  $x_{K+1} \geq 0$  can be enforced by adding a new convex component function  $f_{K+1}(x_{K+1}) = i_{\mathfrak{R}_+^m}(x_{K+1})$  to the objective function  $f(x)$ , where  $i_{\mathfrak{R}_+^m}(x_{K+1})$  is the indicator function for the nonnegative orthant  $\mathfrak{R}_+^m$

$$i_{\mathfrak{R}_+^m}(x_{K+1}) = \begin{cases} 0, & \text{if } x_{K+1} \geq 0 \text{ (entry wise),} \\ \infty, & \text{otherwise.} \end{cases}$$

In this way, the inequality constrained problem with  $K$  blocks is reformulated as an equivalent equality constrained convex minimization problem with  $K + 1$  blocks.

Optimization problems of the form (1.1) arise in many emerging applications involving structured convex optimization. For instance, in compressive sensing applications, we are given an observation matrix  $A$  and a noisy observation vector  $b \approx Ax$ . The goal is to estimate the sparse vector  $x$  by solving the following  $\ell_1$  regularized linear least squares problem:

$$\begin{aligned}
 & \text{minimize} && \|y\|^2 + \lambda \|x\|_1 \\
 & \text{subject to} && Ax + y = b,
 \end{aligned}$$

where  $\lambda > 0$  is a penalty parameter. Clearly, this is a structured convex optimization problem of the form (1.1) with  $K = 2$ . If the variable  $x$  is further constrained to be nonnegative, then the corresponding compressive sensing problem can be formulated as a three block ( $K = 3$ ) convex separable optimization problem (1.1) by introducing a slack variable. Similarly, in the stable version of robust principal component analysis (PCA) [59], we are given an observation matrix  $M \in \mathfrak{R}^{m \times n}$  which is a noise-corrupted sum of a low rank matrix  $L$  and a sparse matrix  $S$ . The goal is recover  $L$  and  $S$  by solving the following nonsmooth convex optimization problem

$$\begin{aligned}
 & \text{minimize} && \|L\|_* + \rho \|S\|_1 + \lambda \|Z\|_F^2 \\
 & \text{subject to} && L + S + Z = M
 \end{aligned}$$

where  $\|\cdot\|_*$  denotes the matrix nuclear norm (defined as the sum of the matrix singular eigenvalues), while  $\|\cdot\|_1$  and  $\|\cdot\|_F$  denote, respectively, the  $\ell_1$  and the Frobenius norm of a matrix (equal to the standard  $\ell_1$  and  $\ell_2$  vector norms when the matrix is viewed as a vector). In the above formulation,  $Z$  denotes the noise matrix, and  $\rho, \lambda$  are some fixed penalty parameters. It is easily seen that the stable robust PCA problem corresponds to the three block case  $K = 3$  in the problem (1.1) with  $x = (L, S, Z)$  and

$$f_1(L) = \|L\|_*, \quad f_2(S) = \|S\|_1, \quad f_3(Z) = \|Z\|_F^2, \quad (1.2)$$

while the coupling linear constraint is given  $L + S + Z = M$ . In image processing applications where the low rank matrix  $L$  is additionally constrained to be nonnegative, then the above problem can be reformulated as

$$\begin{aligned} & \text{minimize} && \|L\|_* + \rho\|S\|_1 + \lambda\|Z\|_F^2 + i_{\mathfrak{R}_+^{mn}}(C) \\ & \text{subject to} && L + S + Z = M, \quad L - C = 0, \end{aligned}$$

where  $C$  is a slack matrix variable of the same size as  $L$ , and  $i_{\mathfrak{R}_+^{mn}}(\cdot)$  is the indicator function for the nonnegative orthant  $\mathfrak{R}_+^{mn}$ . In this case, the stable robust PCA problem is again in the form of (1.1). In particular, it has 4 block variables  $(L, S, Z, C)$  and the first three convex functions are the same as in (1.2), while the fourth convex function is given by  $f_4(C) = i_{\mathfrak{R}_+^{mn}}(C)$ . The coupling linear constraints are  $L + S + Z = M, L - C = 0$ . Other applications of the form (1.1) include the latent variable Gaussian graphical model selection problem, see [9].

A popular approach to solving the separable convex optimization problem (1.1) is to attach a Lagrange multiplier vector  $y$  to the linear constraints  $Ex = q$  and add a quadratic penalty, thus obtaining an augmented Lagrangian function of the form

$$L(x; y) = f(x) + \langle y, q - Ex \rangle + \frac{\rho}{2} \|q - Ex\|^2, \quad (1.3)$$

where  $\rho \geq 0$  is a constant. The augmented dual function is given by

$$d(y) = \min_x f(x) + \langle y, q - Ex \rangle + \frac{\rho}{2} \|q - Ex\|^2 \quad (1.4)$$

and the dual problem (equivalent to (1.1) under mild conditions) is

$$\max_y d(y). \quad (1.5)$$

Moreover, if  $\rho > 0$ , then  $Ex$  is constant over the set of minimizers of (1.4) (see Lemma 2.1 in Section 2). This implies that the dual function  $d(y)$  is differentiable with

$$\nabla d(y) = q - Ex(y)$$

where  $x(y)$  is a minimizer of (1.4). Given the differentiability of  $d(y)$ , it is natural to consider the following dual ascent method to solve the primal problem (1.1)

$$y := y + \alpha \nabla d(y) = y + \alpha(q - Ex(y)), \quad (1.6)$$

where  $\alpha > 0$  is a suitably chosen stepsize. Such a dual ascent strategy is well suited for structured convex optimization problems that are amenable to decomposition. For example, if the objective function  $f$  is separable (i.e., of the form given in (1.1)) and if we select  $\rho = 0$ , then the minimization in (1.4) decomposes into  $K$  independent minimizations whose solutions frequently can be obtained in a simple form. In addition, the iterations can be implemented in a manner that exploits the sparsity structure of the problem and, in certain network cases, achieve a high degree of parallelism. Popular choices for the ascent methods include (single) coordinate ascent (see [4, 8, 10, 34, 40, 42, 51, 52, 58]), gradient ascent (see [34, 42, 53]) and gradient projection [23, 32]. (See [5, 34, 49] for additional references.)

For large scale optimization problems, it is numerically advantageous to select  $\rho > 0$ . Unfortunately, this also introduces variable coupling in the augmented Lagrangian (1.3), which makes the exact minimization step in (1.4) no longer decomposable across variable blocks even if  $f$  has a separable structure. In this case, it is more economical to minimize (1.4) inexactly by updating the components of  $x$  cyclically via the coordinate descent method. In particular, we can apply the Gauss-Seidel strategy to inexactly minimize (1.4), and then update the multiplier  $y$  using an approximate optimal solution of (1.4) in a manner similar to (1.6). The resulting algorithm is called the Alternating Direction Method of Multipliers (ADMM) and is summarized as follows (see [17–20]). In the general context of sums of monotone operators, the work of [16] describes a large family of splitting methods for  $K \geq 3$  blocks which, when applied to the dual, result in similar but not identical methods to the ADMM algorithm (1.7) below.

**Alternating Direction Method of Multipliers (ADMM)**

At each iteration  $r \geq 1$ , we first update the primal variable blocks in the Gauss-Seidel fashion and then update the dual multiplier using the updated primal variables:

$$\begin{cases} x_k^{r+1} = \arg \min_{x_k \in X_k} L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k, x_{k+1}^r, \dots, x_K^r; y^r), & k = 1, 2, \dots, K, \\ y^{r+1} = y^r + \alpha(q - Ex^{r+1}) = y^r + \alpha \left( q - \sum_{k=1}^K E_k x_k^{r+1} \right), \end{cases} \quad (1.7)$$

where  $\alpha > 0$  is the step size for the dual update.

Notice that if there is only one block ( $K = 1$ ), then the ADMM reduces to the standard augmented Lagrangian method of multipliers for which the global convergence is well understood (see e.g., [1]). In particular, it is known that, under mild assumptions on the problem, this type of dual gradient ascent methods generate a sequence of iterates whose limit points must be optimal solutions of the original problem (see [8, 49, 51]). For the special case of ordinary network flow

problems, it is further known that an associated sequence of dual iterates converges to an optimal solution of the dual (see [4]). The rate of convergence of dual ascent methods has been studied in the reference [37] which showed that, under mild assumptions on the problem, the distance to the optimal dual solution set from any  $y \in \mathfrak{R}^m$  near the set is bounded above by the dual optimality ‘residual’  $\|\nabla d(y)\|$ . By using this bound, it can be shown that a number of ascent methods, including coordinate ascent methods and a gradient projection method, converge at least linearly when applied to solve the dual problem (see [35, 36]; also see [2, 11, 29] for related analysis). (Throughout this paper, by ‘linear convergence’ we mean root-linear convergence (denoted by R-linear convergence) in the sense of Ortega and Rheinboldt [41].)

When there are two blocks ( $K = 2$ ), the convergence of the ADMM was studied in the context of Douglas-Rachford splitting method [13–15] for finding a zero of the sum of two maximal monotone operators. It is known that in this case every limit point of the iterates is an optimal solution of the problem. The recent work of [21, 22, 26] have shown that, under some additional assumptions, the objective values generated by the ADMM algorithm and its accelerated version (which performs some additional line search steps for the dual update) converge at a rate of  $O(1/r)$  and  $O(1/r^2)$  respectively. Moreover, if the objective function  $f(x)$  is strongly convex and the constraint matrix  $E$  is row independent, then the ADMM is known to converge linearly to the unique minimizer of (1.1) [33]. [One notable exception to the strong convexity requirement is in the special case of linear programming for which the ADMM is linearly convergent [14].] More recent convergence rate analysis of the ADMM still requires at least one of the component functions ( $f_1$  or  $f_2$ ) to be strongly convex and have a Lipschitz continuous gradient. Under these and additional rank conditions on the constraint matrix  $E$ , some linear convergence rate results can be obtained for a subset of primal and dual variables in the ADMM algorithm (or its variant); see [6, 12, 24]. However, when there are more than two blocks involved ( $K \geq 3$ ), the convergence (or the rate of convergence) of the ADMM method is unknown, and this has been a key open question for several decades. The recent work [38] describes a list of novel applications of the ADMM with  $K \geq 3$  and motivates strongly for the need to analyze the convergence of the ADMM in the multi-block case. The recent monograph [7] contains more details of the history, convergence analysis and applications of the ADMM and related methods.

A main contribution of this paper is to establish the global (linear) convergence of the ADMM method for a class of convex objective functions involving any number of blocks ( $K$  is arbitrary). The key requirement for the global (linear) convergence is the satisfaction of a certain error bound condition that is similar to that used in the analysis of [37]. This error bound estimates the distance from an iterate to the optimal solution set in terms of a certain proximity residual. The class of objective functions that are known to satisfy this error bound condition include many of the compressive sensing applications, such as LASSO [48], Group LASSO [56] or Sparse Group LASSO [57].

## 2 Technical Preliminaries

Let  $f$  be a closed proper convex function in  $\mathfrak{R}^n$ , let  $E$  be an  $m \times n$  matrix, let  $q$  be a vector in  $\mathfrak{R}^m$ . Let  $\text{dom } f$  denote the effective domain of  $f$  and let  $\text{int}(\text{dom } f)$  denote the interior of  $\text{dom } f$ . We make the following standing assumptions regarding  $f$ :

### Assumption A.

(a) The global minimum of (1.1) is attained and so is its dual optimal value. The intersection  $X \cap \text{int}(\text{dom } f) \cap \{x \mid Ex = q\}$  is nonempty.

(b)  $f = f_1(x_1) + f_2(x_2) + \cdots + f_K(x_K)$ , with each  $f_k$  further decomposable as

$$f_k(x_k) = g_k(A_k x_k) + h_k(x_k)$$

where  $g_k$  and  $h_k$  are both convex and continuous over their domains, and  $A_k$ 's are some given matrices (not necessarily full column rank, and can be zero).

(c) Each  $g_k$  is strictly convex and continuously differentiable on  $\text{int}(\text{dom } g_k)$  with a uniform Lipschitz continuous gradient

$$\|\nabla A_k^T g_k(Ax_k) - A_k^T \nabla g_k(Ax'_k)\| \leq L \|x_k - x'_k\|, \quad \forall x_k, x'_k \in X_k$$

where  $L > 0$  is a constant.

(d) Each  $h_k$  satisfies either one of the following conditions

1. The epigraph of  $h_k(x_k)$  is a polyhedral set.
2.  $h_k(x_k) = \lambda_k \|x_k\|_1 + \sum_J w_J \|x_{k,J}\|_2$ , where  $x_k = (\cdots, x_{k,J}, \cdots)$  is a partition of  $x_k$  with  $J$  being the partition index.
3. Each  $h_k(x_k)$  is the sum of the functions described in the previous two items.

(e) For any fixed and finite  $y$  and  $\xi$ ,  $\sum_k h_k(x_k)$  is finite for all  $x \in \{x : L(x; y) \leq \xi\} \cap X$ .

(f) Each submatrix  $E_k$  has full column rank.

(g) The feasible sets  $X_k$ ,  $k = 1, \cdots, K$  are compact polyhedral sets.

We have the following remarks regarding to the assumptions made.

1. Each  $f_k$  may only contain convex function  $h_k$ . That is, the strongly convex part  $g_k$  can be absent. Also, since the matrices  $A_k$ 's are not required to have full column rank, the overall objective function  $f(x)$  is not necessarily strongly convex. In fact, under Assumption A, the optimization problem (1.1) can still have multiple primal or dual optimal solutions. This makes the convergence (and rate of convergence) analysis of ADMM difficult.

2. Assumption (e) does allow  $h_k(\cdot)$  to be an indicator function, as in this case the set  $\{x_k \mid \sum_k h_k(x_k) = \infty\} \cap X$  is not a subset of  $\{x : L(x; y) \leq \xi\} \cap X$  for any given  $y$  and  $\xi$ .
3. Linear term of the form  $\langle b_k, x_k \rangle$  is already included in  $h_k$ , as its ephigraph is polyhedral. Moreover, from the assumption that  $X_k$  is polyhedral, the feasibility constraint  $x_k \in X_k$  can be absorbed into  $h_k$  by adding to it an indicator function  $i_{X_k}(x_k)$ . To simplify notations, we will not explicitly write  $x_k \in X_k$  in the ADMM update (1.7) from now on.
4. Assumption (f) is made to ensure that the subproblems for each  $x_k$  is strongly convex. This assumption will be relaxed later when the subproblems are solved inexactly; see Section 4.1.
5. Assumption (g) requires the feasible set of the variables to be compact, which is needed to ensure that certain error bounds of the primal and dual problems of (1.1) hold. This assumption is usually satisfied in practical applications (e.g. the consensus problems) whenever *a priori* knowledge on the variable domain is available. This assumption can be further relaxed; see the discussion at the end of Section 3.

Under Assumption A, both the primal optimum and the dual optimum values of (1.1) are attained and are equal (i.e., the strong duality holds for (1.1)) so that

$$d^* = \max_y L(x; y) = \max_y \left( f(x) + \langle y, q - Ex \rangle + \frac{\rho}{2} \|Ex - q\|^2 \right) = \min_{Ex=q} f(x),$$

where  $d^*$  is the optimal value of the dual of (1.1).

Under Assumption A, there may still be multiple optimal solutions for both the primal problem (1.1) and its dual problem. We first claim that the dual functional

$$d(y) = \min_x L(x; y) = \min_x f(x) + \langle y, q - Ex \rangle + \frac{\rho}{2} \|q - Ex\|^2, \quad (2.1)$$

is differentiable everywhere. Let  $X(y)$  denote the set of optimal solutions for (2.1).

**Lemma 2.1** *For any  $y \in \mathfrak{R}^m$ , both  $Ex$  and  $A_k x_k$ ,  $k = 1, 2, \dots, K$ , are constant over  $X(y)$ . Moreover, the dual function  $d(y)$  is differentiable everywhere and*

$$\nabla d(y) = q - Ex(y),$$

where  $x(y) \in X(y)$  is any minimizer of (2.1).

**Proof.** Fix  $y \in \mathfrak{R}^m$ . We first show that  $Ex$  is invariant over  $X(y)$ . Suppose the contrary, so that there exist two optimal solutions  $x$  and  $x'$  from  $X(y)$  with the property that  $Ex \neq Ex'$ . Then, we have

$$d(y) = L(x; y) = L(x'; y).$$



Due to the convexity of  $L(x; y)$  with respect to the variable  $x$ , the solution set  $X(y)$  must be convex, implying  $\bar{x} = (x + x')/2 \in X(y)$ . By the convexity of  $f(x)$ , we have

$$\frac{1}{2} [(f(x) + \langle y, q - Ex \rangle) + (f(x') + \langle y, q - Ex' \rangle)] \geq f(\bar{x}) + \langle y, q - E\bar{x} \rangle.$$

Moreover, by the strict convexity of  $\|\cdot\|^2$  and the assumption  $Ex \neq Ex'$ , we have

$$\frac{1}{2} (\|Ex - q\|^2 + \|Ex' - q\|^2) > \|E\bar{x} - q\|^2.$$

Multiplying this inequality by  $\rho/2$  and adding it to the previous inequality yields

$$\frac{1}{2} [L(x; y) + L(x'; y)] > L(\bar{x}; y),$$

which further implies

$$d(y) > L(\bar{x}; y).$$

This contradicts the definition  $d(y) = \min_x L(x; y)$ . Thus,  $Ex$  is invariant over  $X(y)$ . Notice that  $d(y)$  is a concave function and its subdifferential is given by [1]

$$\partial d(y) = \text{Closure of the convex hull } \{ q - Ex(y) \mid x(y) \in X(y) \}.$$

Since  $Ex(y)$  is invariant over  $X(y)$ , the subdifferential  $\partial d(y)$  is a singleton. By Danskin's Theorem, this implies that  $d(y)$  is differentiable and the gradient is given by  $\nabla d(y) = q - Ex(y)$ , for any  $x(y) \in X(y)$ .

A similar argument (and using the strict convexity of  $g_k$ ) shows that  $A_k x_k$  is also invariant over  $X(y)$ . The proof is complete. **Q.E.D.**

By using Lemma 2.1, we show below a Lipschitz continuity property of  $\nabla d(y)$ , for  $y$  over any level set of  $d$ .

**Lemma 2.2** *Fix any scalar  $\eta \leq f^*$  and let  $\mathcal{U} = \{ y \in \mathfrak{R}^m \mid d(y) \geq \eta \}$ . Then there holds*

$$\|\nabla d(y') - \nabla d(y)\| \leq \frac{1}{\rho} \|y' - y\|, \quad \forall y' \in \mathcal{U}, y \in \mathcal{U}.$$

**Proof.** Fix any  $y$  and  $y'$  in  $\mathcal{U}$ . Let  $x = x(y)$  and  $x' = x(y')$  be two minimizers of  $L(x; y)$  and  $L(x; y')$  respectively. By convexity, we have

$$z - E^T y + \rho E^T (Ex - q) = 0 \quad \text{and} \quad z' - E^T y' + \rho E^T (Ex' - q) = 0,$$

where  $z$  and  $z'$  are some subgradient vectors in the subdifferential  $\partial f(x)$  and  $\partial f(x')$  respectively. Thus, we have

$$\langle z - E^T y + \rho E^T (Ex - q), x' - x \rangle = 0$$

and

$$\langle z' - E^T y' + \rho E^T (Ex' - q), x - x' \rangle = 0.$$

Adding the above two equalities yields

$$\langle z - z' + E^T (y' - y) - \rho E^T E(x' - x), x' - x \rangle = 0.$$

Upon rearranging terms and using the convexity property

$$\langle z' - z, x' - x \rangle \geq 0,$$

we get

$$\langle y' - y, E(x' - x) \rangle = \langle z' - z, x' - x \rangle + \rho \|E(x' - x)\|^2 \geq \rho \|E(x' - x)\|^2.$$

Thus,  $\rho \|E(x' - x)\| \leq \|y' - y\|$  which together with  $\nabla d(y') - \nabla d(y) = E(x - x')$  (cf. Lemma 2.1) yields

$$\|\nabla d(y') - \nabla d(y)\| = \|E(x' - x)\| \leq \frac{1}{\rho} \|y - y'\|.$$

The proof is complete. **Q.E.D.**

To show the linear convergence of the ADMM method, we need certain local error bounds around the optimal solution set  $X(y)$  as well as around the dual optimal solution set  $Y^*$ . To describe these local error bounds, we first define the notion of a proximity operator. Let  $h : \text{dom}(h) \mapsto \mathfrak{R}$  be a (possibly nonsmooth) convex function. For every  $x \in \text{dom}(h)$ , the *proximity operator* of  $h$  is defined as [44]

$$\text{prox}_h(x) = \underset{u \in \mathfrak{R}^n}{\text{argmin}} \quad h(u) + \frac{1}{2} \|x - u\|^2.$$

Notice that if  $h(x)$  is the indicator function of a closed convex set  $X$ , then

$$\text{prox}_h(x) = \text{proj}_X(x),$$

so the proximity operator is a generalization of the projection operator. In particular, it is known that the proximity operator satisfies the nonexpansiveness property:

$$\|\text{prox}_h(x) - \text{prox}_h(x')\| \leq \|x - x'\|, \quad \forall x, x'. \quad (2.2)$$

The proximity operator can be used to characterize the optimality condition for a nonsmooth convex optimization problem. Suppose a convex function  $f$  is decomposed as  $f(x) = g(Ax) + h(x)$

where  $g$  is strongly convex and differentiable,  $h$  is a convex (possibly nonsmooth) function, then we can define the *proximal gradient* of  $f$  with respect to  $h$  as

$$\tilde{\nabla}f(x) := x - \text{prox}_h(x - \nabla(f(x) - h(x))) = x - \text{prox}_h(x - A^T \nabla g(Ax)).$$

If  $h \equiv 0$ , then the proximal gradient  $\tilde{\nabla}f(x) = \nabla f(x)$ . In general,  $\tilde{\nabla}f(x)$  can be used as the (standard) gradient of  $f$  for the nonsmooth minimization  $\min_{x \in X} f(x)$ . For example,  $\tilde{\nabla}f(x^*) = 0$  iff  $x^*$  is a global minimizer.

For the Lagrangian minimization problem (2.1) and under Assumption A, the work of [37,50,57] suggests that the size of the proximal gradient

$$\begin{aligned} \tilde{\nabla}_x L(x; y) &:= x - \text{prox}_h(x - \nabla_x(L(x; y) - h(x))) \\ &= x - \text{prox}_h(x - A^T \nabla g(Ax) + E^T y - \rho E^T (Ex - q)) \end{aligned} \quad (2.3)$$

can be used to upper bound the distance to the optimal solution set  $X(y)$  of (2.1). Here

$$h(x) := \sum_{k=1}^K h_k(x_k), \quad g(Ax) := \sum_{k=1}^K g_k(A_k x_k)$$

represent the nonsmooth and the smooth parts of  $f(x)$  respectively.

In our analysis of ADMM, we will also need an error bound for the dual function  $d(y)$ . Notice that a  $y \in \Re^m$  solves (1.5) if and only if  $y$  satisfies the system of nonlinear equations

$$\nabla d(y) = 0.$$

This suggests that the norm of the ‘residual’  $\|\nabla d(y)\|$  may be a good estimate of how close  $y$  is from solving (1.5). The next lemma says if the nonsmooth part of  $f_k$  takes certain forms, then the distance to the primal and dual optimal solution sets can indeed be bounded.

**Lemma 2.3** *Suppose assumptions A(a)—A(e) hold.*

1. *If in addition  $X$  is a polyhedral set, then there exists a positive scalar  $\tau$  and  $\delta$  such that the following error bound holds*

$$\text{dist}(x, X(y)) \leq \tau \|\tilde{\nabla}_x L(x; y)\|, \quad (2.4)$$

*for all  $(x, y)$  such that  $\|\tilde{\nabla}_x L(x; y)\| \leq \delta$ , where the proximal gradient  $\tilde{\nabla}_x L(x; y)$  is given by (2.3). Furthermore, if  $X$  is also a compact set, then there exists some  $\tau > 0$  such that the error bound (2.4) holds for all  $x \in X \cap \text{dom}(h)$ .*

2. Similarly, if assumption  $A-(g)$  also holds, then for any scalar  $\zeta$ , there exist positive scalars  $\delta$  and  $\tau$  such that

$$\text{dist}(y, Y^*) = \|y - y^*\| \leq \tau \|\nabla d(y)\|, \text{ whenever } d(y) \geq \zeta \text{ and } \|\nabla d(y)\| \leq \delta. \quad (2.5)$$

Moreover, in both cases the constant  $\tau$  is independent of the choice of  $y$  and  $x$ .

For any fixed  $y$ , the proof for the first part of Lemma 2.3 is identical to those of [37, 50, 57], each of which shows the error bound with different assumptions on the objective function  $f$ . In particular, it was shown that (2.4) holds for all  $x$  with  $\|\tilde{\nabla}_x L(x; y)\| \leq \delta$  (i.e., sufficiently close to  $X(y)$ ). An important new ingredient is the claim that the error bound holds over the compact set  $X \cap \text{dom}(h)$ . This can be seen in two steps as follows: (1) for all  $x \in X \cap \text{dom}(h)$  such that  $\|\tilde{\nabla}_x L(x; y)\| \leq \delta$ , the error bound (2.4) is already known to hold; (2) for all  $x \in X \cap \text{dom}(h)$  such that  $\|\tilde{\nabla}_x L(x; y)\| \geq \delta$ , the ratio

$$\frac{\text{dist}(x, X(y))}{\|\tilde{\nabla}_x L(x; y)\|}$$

is a continuous function and well defined over the compact set  $X \cap \text{dom}(h) \cap \{x \mid \|\tilde{\nabla}_x L(x; y)\| \geq \delta\}$ . Thus, the above ratio must be bounded from above by a constant  $\tau'$  (independent of  $y$ ). Combining (1) and (2) yields the desired error bound over the set  $X \cap \text{dom}(h)$ .

Another new ingredient in Lemma 2.3 is the additional claim that the constants  $\delta, \tau$  are both independent of the choice of  $y$ . This property follows directly from a similar property of Hoffman's error bound [28] (on which the error bounds of [37, 50, 57] are based) for a feasible linear system  $P := \{x \mid Ax \leq b\}$ :

$$\text{dist}(x, P) \leq \tau \| [Ax - b]_+ \|, \quad \forall x \in \mathfrak{R}^m,$$

where  $\tau$  is independent of  $b$ . In fact, a careful checking of the proofs of [37, 50, 57] shows that the corresponding error constants  $\delta$  and  $\tau$  for the augmented Lagrangian function  $L(x; y)$  can be indeed made *independent* of  $y$ . We omit the proof of the first part of Lemma 2.3 for space consideration.

Dual error bounds like the one stated in the second part of the lemma have been studied previously by Pang [43] and by Mangasarian and Shiau [39], though in different contexts. The above error bound is 'local' in that it holds only for those  $y$  that are bounded or near  $Y^*$  (i.e., when  $\|\nabla d(y)\| \leq \delta$  as opposed to a 'global' error bound which would hold for all  $y$  in  $\mathfrak{R}^m$ ). However if in addition  $y$  also lies in some compact set  $Y$ , then the dual error bound hold true for all  $y \in Y$  (using the same argument as in the preceding paragraph). In the appendix, we include a proof showing that the dual error bound holds true, for the case where the epigraph of  $h_k$  is polyhedral (which includes  $\ell_1$  norm and indicator function for polyhedral sets). We note that from this proof it is clear that indeed the value of  $\tau$  in the dual error bound does not depend on the choice of either  $x$  or  $y$ .

Under Assumption A(f), the augmented Lagrangian function  $L(x; y)$  (cf. (1.3)) is strongly convex with respect to each subvector  $x_k$ . As a result, each alternating minimization iteration of ADMM (1.7)

$$x_k^{r+1} = \underset{x_k}{\operatorname{argmin}} L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k, x_{k+1}^r, \dots, x_K^r; y^r), \quad k = 1, \dots, K.$$

has a unique optimal solution. Thus the sequence of iterates  $\{x^r\}$  of the ADMM are well defined. The following lemma shows that the alternating minimization of the Lagrangian function gives a sufficient descent of the Lagrangian function value.

**Lemma 2.4** *Suppose Assumptions A(b) and A(f) hold. Then fix any index  $r$ , we have*

$$L(x^r; y^r) - L(x^{r+1}; y^r) \geq \gamma \|x^r - x^{r+1}\|^2, \quad (2.6)$$

where the constant  $\gamma > 0$  is independent of  $r$  and  $y^r$ .

**Proof.** By assumptions A(b) and A(f), the augmented Lagrangian function

$$L(x; y) = \sum_{k=1}^K (f_k(x_k) + \langle y_k, q_k - E_k x_k \rangle) + \frac{\rho}{2} \left\| \sum_{k=1}^K E_k x_k - q \right\|^2$$

is strongly convex in each variable  $x_k$  and has a uniform modulus  $\rho \lambda_{\min}(E_k^T E_k) > 0$ . Here, the notation  $\lambda_{\min}(\cdot)$  denotes the smallest eigenvalue of a symmetric matrix. This implies that, for each  $k$ , that

$$L(x; y) - L(x_1, \dots, x_{k-1}, \bar{x}_k, x_{k+1}, \dots, x_K; y) \geq \rho \lambda_{\min}(E_k^T E_k) \|x_k - \bar{x}_k\|^2, \quad (2.7)$$

for all  $x$ , where  $\bar{x}_k$  is the minimizer of  $\min_{x_k} L(x; y)$  (when all other variables  $\{x_j\}_{j \neq k}$  are fixed).

Fix any index  $r$ . For each  $k \in \{1, \dots, K\}$ , by ADMM (1.7),  $x_k^{r+1}$  is the minimizer of  $L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k, x_{k+1}^r, x_{k+2}^r, \dots, x_K^r; y^r)$ . It follows from (2.7)

$$L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k^r, \dots, x_K^r; y^r) - L(x_1^{r+1}, \dots, x_k^{r+1}, x_{k+1}^r, \dots, x_K^r; y^r) \geq \gamma \|x_k^r - x_k^{r+1}\|^2, \quad \forall k, \quad (2.8)$$

where

$$\gamma = \rho \min_k \lambda_{\min}(E_k^T E_k)$$

is independent of  $r$  and  $y^r$ . Summing this over  $k$ , we obtain the sufficient decrease condition

$$L(x^r; y^r) - L(x^{r+1}; y^r) \geq \gamma \|x^r - x^{r+1}\|^2.$$

This completes the proof of Lemma 2.4.

**Q.E.D.**

To prove the linear convergence of the ADMM algorithm, we also need the following lemma which bounds the size of the proximal gradient  $\tilde{\nabla} L(x^r; y^r)$  at an iterate  $x^r$ .

**Lemma 2.5** *Suppose assumptions A(b)—A(c) hold. Let  $\{x^r\}$  be generated by the ADMM algorithm (1.7). Then there exists some constant  $\sigma > 0$  (independent of  $y^r$ ) such that*

$$\|\tilde{\nabla}L(x^r; y^r)\| \leq \sigma \|x^{r+1} - x^r\| \quad (2.9)$$

for all  $r \geq 1$ .

**Proof.** Fix any  $r \geq 1$  and any  $1 \leq k \leq K$ . According to the ADMM procedure (1.7), the variable  $x_k$  is updated as follows

$$x_k^{r+1} = \underset{x_k}{\operatorname{argmin}} \left( h_k(x_k) + g_k(A_k x_k) - \langle y^r, E_k x_k \rangle + \frac{\rho}{2} \left\| E_k x_k + \sum_{j < k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right\|^2 \right).$$

The corresponding optimality condition can be written as

$$x_k^{r+1} = \operatorname{prox}_{h_k} \left[ x_k^{r+1} - A_k^T \nabla_{x_k} g_k(A_k x_k^{r+1}) + E_k^T y^r - \rho E_k^T \left( \sum_{j \leq k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right) \right]. \quad (2.10)$$

Therefore, we have

$$\begin{aligned} & \left\| x_k^{r+1} - \operatorname{prox}_{h_k} \left( x_k^r - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r - \rho E_k^T (E x^r - q) \right) \right\| = \\ & \left\| \operatorname{prox}_{h_k} \left[ x_k^{r+1} - A_k^T \nabla_{x_k} g_k(A_k x_k^{r+1}) + E_k^T y^r + \rho E_k^T \left( \sum_{j \leq k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right) \right] \right. \\ & \quad \left. - \operatorname{prox}_{h_k} \left( x_k^r - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r + \rho E_k^T (E x^r - q) \right) \right\| \\ & \leq \left\| (x_k^{r+1} - x_k^r) - A_k^T (\nabla_{x_k} g_k(A_k x_k^{r+1}) - \nabla_{x_k} g_k(A_k x_k^r)) + \rho E_k^T \sum_{j \leq k} E_j (x_j^{r+1} - x_j^r) \right\| \\ & \leq \|x_k^{r+1} - x_k^r\| + L \|A_k^T\| \|A_k\| \|x_k^{r+1} - x_k^r\| + \rho \|E_k^T\| \sum_{j \leq k} \|E_j\| \|x_j^{r+1} - x_j^r\| \\ & \leq c \|x^{r+1} - x^r\|, \quad \text{for some } c > 0 \text{ independent of } y^r, \end{aligned} \quad (2.11)$$

where the first inequality follows from the nonexpansive property of the prox operator (2.2), and the second inequality is due to the Lipschitz property of the gradient vector  $\nabla g_k$  (cf. Assumption A-(c)). Using this relation and the definition of the proximal gradient  $\tilde{\nabla}L(x^r; y^r)$ , we have

$$\begin{aligned} \|\tilde{\nabla}_{x_k} L(x^r; y^r)\| &= \left\| x_k^r - \operatorname{prox}_{h_k} \left( x_k^r - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r - \rho E_k^T (E x^r - q) \right) \right\| \\ &\leq \|x_k^r - x_k^{r+1}\| + \left\| x_k^{r+1} - \operatorname{prox}_{h_k} \left( x_k^r - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r - \rho E_k^T (E x^r - q) \right) \right\| \\ &\leq (c + 1) \|x^{r+1} - x^r\|, \quad \forall k = 1, 2, \dots, K. \end{aligned}$$

This further implies that the entire proximal gradient vector can be bounded by  $\|x^{r+1} - x^r\|$ :

$$\|\tilde{\nabla}L(x^r; y^r)\| \leq (c+1)\sqrt{K}\|x^{r+1} - x^r\|.$$

Setting  $\sigma = (c+1)\sqrt{K}$  (which is independent of  $y^r$ ) completes the proof.

**Q.E.D.**

### 3 Linear Convergence of ADMM

Let  $d^*$  denote the dual optimal value and  $\{x^r, y^r\}$  be the sequence generated by the ADMM method (1.7). Due to assumption A(a),  $d^*$  also equals to the primal optimal value. Further we denote

$$\Delta_d^r = d^* - d(y^r) \tag{3.1}$$

which represents the gap from dual optimality at the  $r$ -th iteration. The primal gap to optimality at iteration  $r$  is defined as

$$\Delta_p^r = L(x^{r+1}; y^r) - d(y^r), \quad r \geq 1. \tag{3.2}$$

Clearly, we have both  $\Delta_d^r \geq 0$  and  $\Delta_p^r \geq 0$  for all  $r$ . To establish the linear convergence of ADMM, we need several lemmas to estimate the sizes of the primal and dual optimality gaps as well as their respective decrease.

Let  $X(y^r)$  denote the set of optimal solutions for the following optimization problem

$$\min_x L(x; y^r) = \min_x f(x) + \langle y^r, q - Ex \rangle + \frac{\rho}{2}\|Ex - q\|^2.$$

We denote

$$\bar{x}^r = \operatorname{argmin}_{\bar{x} \in X(y^r)} \|\bar{x} - x^r\|.$$

We first bound the sizes of the dual and primal optimality gaps.

**Lemma 3.1** *Suppose assumptions A(a)—A(e) and A(g) hold. Then for any scalar  $\delta > 0$ , there exists a positive scalar  $\tau'$  such that*

$$\Delta_d^r \leq \tau' \|\nabla d(y^r)\|^2 = \tau' \|Ex(y^r) - q\|^2, \tag{3.3}$$

for any  $y^r \in \Re^m$  with  $\|\nabla d(y^r)\| \leq \delta$ . Moreover, there exist positive scalars  $\zeta$  and  $\zeta'$  (independent of  $y^r$ ) such that

$$\Delta_p^r \leq \zeta \|x^{r+1} - x^r\|^2 + \zeta' \|x^r - \bar{x}^r\|^2, \quad \text{for all } r \geq 1. \tag{3.4}$$

**Proof.** Fix any  $y^r$ , and let  $y^*$  be the optimal dual solution closest to  $y^r$ . Then it follows from the mean value theorem that there exists some  $\tilde{y}$  in the line segment joining  $y^r$  and  $y^*$  such that

$$\begin{aligned}
\Delta_d^r &= d(y^*) - d(y^r) \\
&= \langle \nabla d(\tilde{y}), y^* - y^r \rangle \\
&= \langle \nabla d(\tilde{y}) - \nabla d(y^*), y^* - y^r \rangle \\
&\leq \|\nabla d(\tilde{y}) - \nabla d(y^*)\| \|y^* - y^r\| \\
&\leq \frac{1}{\rho} \|\tilde{y} - y^*\| \|y^* - y^r\| \\
&\leq \frac{1}{\rho} \|y^r - y^*\| \|y^* - y^r\| \\
&= \frac{1}{\rho} \|y^* - y^r\|^2
\end{aligned}$$

where the second inequality follows from Lemma 2.2. Recall from the second part in Lemma 2.3 that there exists some  $\tau$  such that

$$\text{dist}(y^r, Y^*) = \|y^r - y^*\| \leq \tau \|\nabla d(y^r)\|.$$

Combining the above two inequalities yields

$$\Delta_d^r = d(y^*) - d(y^r) \leq \tau' \|\nabla d(y^r)\|^2,$$

where  $\tau' = \tau^2/\rho$  is a constant. This establishes the bound on the size of dual gap (3.3).

It remains to prove the bound on the primal gap (3.4). For notational simplicity, let us separate the smooth and nonsmooth part of the augmented Lagrangian as follows

$$L(x; y) = g(x) + h(x) + \langle y, q - Ex \rangle + \frac{\rho}{2} \|q - Ex\|^2 := \bar{L}(x; y) + h(x).$$

Let  $x_k^{r+1}$  denote the  $k$ -th subvector of the primal vector  $x^{r+1}$ . From the way that the variables are updated (2.10), we have

$$\begin{aligned}
x_k^{r+1} &= \text{prox}_{h_k} \left[ x_k^{r+1} - \nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right) \right] \\
&= \text{prox}_{h_k} \left[ x_k^r - \nabla_{x_k} \bar{L}(x^r; y^r) - x_k^r + x_k^{r+1} - \nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right) + \nabla_{x_k} \bar{L}(x^r; y^r) \right] \\
&:= \text{prox}_{h_k} \left[ x_k^r - \nabla_{x_k} \bar{L}(x^r; y^r) - e_k^r \right] \tag{3.5}
\end{aligned}$$

where the gradient vector  $\nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right)$  can be explicitly expressed as

$$\nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right) = A_k^T \nabla_{x_k} g(A_k x_k^{r+1}) - E_k^T y^r + \rho E_k^T \left( \sum_{j \leq k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right)$$



and the error vector  $e_k^r$  is defined by

$$e_k^r := x_k^r - x_k^{r+1} + \nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right) - \nabla_{x_k} \bar{L}(x^r; y^r). \quad (3.6)$$

Note that we can bound the norm of  $e_k^r$  as follows

$$\begin{aligned} \|e_k^r\| &\leq \|x_k^r - x_k^{r+1}\| + \|\nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}_{j \leq k}, \{x_j^r\}_{j > k}; y^r \right) - \nabla_{x_k} \bar{L}(x^r; y^r)\| \\ &\leq \|x_k^r - x_k^{r+1}\| + \left\| A_k^T \left( \nabla_{x_k} g(A_k x_k^{r+1}) - \nabla_{x_k} g(A_k x_k^r) \right) + \rho E_k^T \left( \sum_{j \leq k} E_j (x_j^{r+1} - x_j^r) \right) \right\| \\ &\leq c \|x^r - x^{r+1}\|, \end{aligned} \quad (3.7)$$

where the constant  $c > 0$  is independent of  $y^r$ , and can take the same value as in (2.11).

Using (3.5), and by the definition of the proximity operator, we have the following

$$\begin{aligned} h_k(x_k^{r+1}) + \langle x_k^{r+1} - x_k^r, \nabla_{x_k} \bar{L}(x^r; y^r) + e_k^r \rangle + \frac{1}{2} \|x_k^{r+1} - x_k^r\|^2 \\ \leq h_k(\bar{x}_k^r) + \langle \bar{x}_k^r - x_k^r, \nabla_{x_k} \bar{L}(x^r; y^r) + e_k^r \rangle + \frac{1}{2} \|\bar{x}_k^r - x_k^r\|^2. \end{aligned} \quad (3.8)$$

Summing over all  $k = 1, \dots, K$ , we obtain

$$\begin{aligned} h(x^{r+1}) + \langle x^{r+1} - x^r, \nabla_x \bar{L}(x^r; y^r) + e^r \rangle + \frac{1}{2} \|x^{r+1} - x^r\|^2 \\ \leq h(\bar{x}^r) + \langle \bar{x}^r - x^r, \nabla_x \bar{L}(x^r; y^r) + e^r \rangle + \frac{1}{2} \|\bar{x}^r - x^r\|^2. \end{aligned}$$

Upon rearranging terms, we obtain

$$h(x^{r+1}) - h(\bar{x}^r) + \langle x^{r+1} - \bar{x}^r, \nabla_x \bar{L}(x^r; y^r) \rangle \leq \frac{1}{2} \|\bar{x}^r - x^r\|^2 - \langle x^{r+1} - \bar{x}^r, e^r \rangle. \quad (3.9)$$

Also, we have from the mean value theorem that there exists some  $\tilde{x}$  in the line segment joining  $x^{r+1}$  and  $\bar{x}^r$  such that

$$\bar{L}(x^{r+1}; y^r) - \bar{L}(\bar{x}^r; y^r) = \langle \nabla_x \bar{L}(\tilde{x}; y^r), x^{r+1} - \bar{x}^r \rangle.$$

Using the above results, we can bound  $\Delta_p^r$  by

$$\begin{aligned}
\Delta_p^r &= L(x^{r+1}; y^r) - L(\bar{x}^r; y^r) \\
&= \bar{L}(x^{r+1}; y^r) - \bar{L}(\bar{x}^r; y^r) + h(x^{r+1}) - h(\bar{x}^r) \\
&= \langle \nabla_x \bar{L}(\tilde{x}; y^r), x^{r+1} - \bar{x}^r \rangle + h(x^{r+1}) - h(\bar{x}^r) \\
&= \langle \nabla_x \bar{L}(\tilde{x}; y^r) - \nabla_x \bar{L}(x^r; y^r), x^{r+1} - \bar{x} \rangle + \langle \nabla_x \bar{L}(x^r; y^r), x^{r+1} - \bar{x} \rangle + h(x^{r+1}) - h(\bar{x}^r) \\
&\leq \langle \nabla_x \bar{L}(\tilde{x}; y^r) - \nabla_x \bar{L}(x^r; y^r), x^{r+1} - \bar{x} \rangle + \frac{1}{2} \|\bar{x}^r - x^r\|^2 + c\sqrt{K} \|x^{r+1} - x^r\| \|x^{r+1} - \bar{x}^r\| \\
&\leq \left( \sum_{k=1}^K L \|A_k\|^T \|A_k\| + \rho \|E^T E\| \right) \|\tilde{x} - x^r\| \|x^{r+1} - \bar{x}^r\| \\
&\quad + \frac{1}{2} \|\bar{x}^r - x^r\|^2 + c\sqrt{K} \|x^{r+1} - x^r\| \|x^{r+1} - \bar{x}^r\| \\
&\leq \left( \sum_{k=1}^K L \|A_k\|^T \|A_k\| + \rho \|E^T E\| \right) (\|x^{r+1} - x^r\| + \|\bar{x}^r - x^r\|)^2 \\
&\quad + \frac{1}{2} \|\bar{x}^r - x^r\|^2 + c\sqrt{K} \|x^{r+1} - x^r\| (\|x^{r+1} - x^r\| + \|\bar{x}^r - x^r\|) \\
&\leq \zeta \|x^{r+1} - x^r\|^2 + \zeta' \|\bar{x}^r - x^r\|^2, \quad \text{for some } \zeta, \zeta' > 0,
\end{aligned}$$

where the first inequality follows from (3.9) and (3.7), the second inequality is due to the Cauchy-Schwartz inequality and the Lipschitz continuity of  $\nabla \bar{L}_x(x; y^r)$ , while the third inequality follows from the fact that  $\tilde{x}$  lies in the line segment joining  $x^{r+1}$  and  $\bar{x}^r$  so that  $\|\tilde{x} - x^r\| \leq \|x^{r+1} - x^r\| + \|\bar{x}^r - x^r\|$ . This completes the proof. **Q.E.D.**

We then bound the decrease of the dual optimality gap.

**Lemma 3.2** *For each  $r \geq 1$ , there holds*

$$\Delta_d^r - \Delta_d^{r-1} \leq -\alpha (Ex^r - q)^T (E\bar{x}^r - q). \quad (3.10)$$

**Proof.** The reduction of the optimality gap in the dual space can be bounded as follows:

$$\begin{aligned}
\Delta_d^r - \Delta_d^{r-1} &= [d^* - d(y^r)] - [d^* - d(y^{r-1})] \\
&= d(y^{r-1}) - d(y^r) \\
&= L(\bar{x}^{r-1}; y^{r-1}) - L(\bar{x}^r; y^r) \\
&= [L(\bar{x}^r; y^{r-1}) - L(\bar{x}^r; y^r)] + [L(\bar{x}^{r-1}; y^{r-1}) - L(\bar{x}^r; y^{r-1})] \\
&= (y^{r-1} - y^r)^T (q - E\bar{x}^r) + [L(\bar{x}^{r-1}; y^{r-1}) - L(\bar{x}^r; y^{r-1})] \\
&= -\alpha (Ex^r - q)^T (E\bar{x}^r - q) + [L(\bar{x}^{r-1}; y^{r-1}) - L(\bar{x}^r; y^{r-1})] \\
&\leq -\alpha (Ex^r - q)^T (E\bar{x}^r - q), \quad \forall r \geq 1,
\end{aligned}$$

where the last equality follows from the update of the dual variable  $y^{r-1}$ , and the fact that  $\bar{x}^{r-1}$  minimizes  $L(\cdot, y^{r-1})$ . **Q.E.D.**

Lemma 3.2 implies that if  $q - Ex^r$  is close to the true dual gradient  $\nabla d(y^r) = q - E\bar{x}^r$ , then the dual optimal gap is reduced after each ADMM iteration. However, since ADMM updates the primal variable by only one Gauss-Seidel sweep, the primal iterate  $x^r$  is not necessarily close the minimizer  $\bar{x}^r$  of  $L(x; y^r)$ . Thus, unlike the method of multipliers (for which  $x^r = \bar{x}^r$  for all  $r$ ), there is no guarantee that the dual optimality gap  $\Delta_d^r$  is indeed reduced after each iteration of ADMM.

Next we proceed to bound the decrease in the primal gap  $\Delta_p^r$ .

**Lemma 3.3** *Suppose assumptions A(b) and A(f) hold. Then for each  $r \geq 1$ , we have*

$$\Delta_p^r - \Delta_p^{r-1} \leq \alpha \|Ex^r - q\|^2 - \gamma \|x^{r+1} - x^r\|^2 - \alpha (Ex^r - q)^T (E\bar{x}^r - q) \quad (3.11)$$

for some  $\gamma$  independent of  $y^r$ .

**Proof.** Fix any  $r \geq 1$ , we have

$$L(x^r; y^{r-1}) = f(x^r) + \langle y^{r-1}, q - Ex^r \rangle + \frac{\rho}{2} \|Ex^r - q\|^2$$

and

$$L(x^{r+1}; y^r) = f(x^{r+1}) + \langle y^r, q - Ex^{r+1} \rangle + \frac{\rho}{2} \|Ex^{r+1} - q\|^2.$$

By the update rule of  $y^r$  (cf. (1.7)), we have

$$L(x^r; y^r) = f(x^r) + \langle y^{r-1}, q - Ex^r \rangle + \frac{\rho}{2} \|Ex^r - q\|^2 + \alpha \|Ex^r - q\|^2.$$

This implies

$$L(x^r; y^r) = L(x^r; y^{r-1}) + \alpha \|Ex^r - q\|^2.$$

Recall from Lemma 2.4 that the alternating minimization of the Lagrangian function gives a sufficient descent. In particular, we have

$$L(x^{r+1}; y^r) - L(x^r; y^r) \leq -\gamma \|x^{r+1} - x^r\|^2,$$

for some  $\gamma > 0$  that is independent of  $r$  and  $y^r$ . Therefore, we have

$$L(x^{r+1}; y^r) - L(x^r; y^{r-1}) \leq \alpha \|Ex^r - q\|^2 - \gamma \|x^{r+1} - x^r\|^2, \quad \forall r \geq 1.$$

Hence, we have the following bound on the reduction of primal optimality gap

$$\begin{aligned} \Delta_p^r - \Delta_p^{r-1} &= [L(x^{r+1}; y^r) - d(y^r)] - [L(x^r; y^{r-1}) - d(y^{r-1})] \\ &= [L(x^{r+1}; y^r) - L(x^r; y^{r-1})] - [d(y^r) - d(y^{r-1})] \\ &\leq \alpha \|Ex^r - q\|^2 - \gamma \|x^{r+1} - x^r\|^2 - \alpha (Ex^r - q)^T (E\bar{x}^r - q), \quad \forall r \geq 1, \end{aligned}$$

where the last step is due to Lemma 3.2.

**Q.E.D.**

Notice that when  $\alpha = 0$  (i.e., no dual update in the ADMM algorithm), Lemma 3.3 reduces to the sufficient decrease estimate (2.6) in Lemma 2.4. When  $\alpha > 0$ , the primal optimality gap is not necessarily reduced after each ADMM iteration due to the positive term  $\alpha\|Ex^r - q\|^2$  in (3.11). Thus, in general, we cannot guarantee a consistent decrease of either the dual optimality gap  $\Delta_d^r$  or the primal optimality gap  $\Delta_p^r$ . However, somewhat surprisingly, the sum of the primal and dual optimality gaps decreases for all  $r$ , as long as the dual step size  $\alpha$  is sufficiently small. This is used to establish the linear convergence of ADMM method.

**Theorem 3.1** *Suppose the conditions in Assumption A hold. Then the sequence of iterates  $\{(x^r, y^r)\}$  generated by the ADMM algorithm (1.7) converges linearly to an optimal primal-dual solution for (1.1), provided the stepsize  $\alpha$  is sufficiently small. Moreover, the sequence of feasibility violation  $\{\|Ex^r - q\|\}$  also converges linearly.*

**Proof.** We show by induction that the sum of optimality gaps  $\Delta_d^r + \Delta_p^r$  is reduced after each ADMM iteration, as long as the stepsize  $\alpha$  is chosen sufficiently small. For any  $r \geq 1$ , we denote

$$\bar{x}^r = \operatorname{argmin}_{\bar{x} \in X(y^r)} \|\bar{x} - x^r\|. \quad (3.12)$$

By induction, suppose  $\Delta_d^{r-1} + \Delta_p^{r-1} \leq \Delta_d^0 + \Delta_p^0$  for some  $r \geq 1$ . Recall that each  $X_k$  is compact and that the indicator function  $i_{X_k}(x_k)$  is included in  $h_k(x_k)$  (see the discussion after Assumption A), it follows that  $x^r \in X$ , implying the boundedness of  $x^r$ . Thus, we obtain from Lemma 2.3 that

$$\|x^r - \bar{x}^r\| \leq \tau \|\tilde{\nabla}L(x^r; y^r)\| \quad (3.13)$$

for some  $\tau > 0$  (independent of  $y^r$ ). To prove Theorem 3.1, we combine the two estimates (3.10) and (3.11) to obtain

$$\begin{aligned} [\Delta_p^r + \Delta_d^r] - [\Delta_p^{r-1} + \Delta_d^{r-1}] &= [\Delta_p^r - \Delta_p^{r-1}] + [\Delta_d^r - \Delta_d^{r-1}] \\ &\leq \alpha\|Ex^r - q\|^2 - \gamma\|x^{r+1} - x^r\|^2 - 2\alpha(Ex^r - q)^T(E\bar{x}^r - q) \\ &= \alpha\|Ex^r - E\bar{x}^r\|^2 - \alpha\|E\bar{x}^r - q\|^2 - \gamma\|x^{r+1} - x^r\|^2. \end{aligned} \quad (3.14)$$

Now we invoke (3.13) and Lemma 2.5 to lower bound  $\|x^{r+1} - x^r\|$ :

$$\|x^r - \bar{x}^r\| \leq \tau \|\tilde{\nabla}L(x^r; y^r)\| \leq \tau\sigma\|x^{r+1} - x^r\|. \quad (3.15)$$

Substituting this bound into (3.14) yields

$$[\Delta_p^r + \Delta_d^r] - [\Delta_p^{r-1} + \Delta_d^{r-1}] \leq (\alpha\|E\|^2\tau^2\sigma^2 - \gamma)\|x^{r+1} - x^r\|^2 - \alpha\|E\bar{x}^r - q\|^2. \quad (3.16)$$

Thus, if we choose the stepsize  $\alpha$  sufficiently small so that

$$0 < \alpha < \gamma\tau^{-2}\sigma^{-2}\|E\|^{-2}, \quad (3.17)$$

then the above estimate shows that

$$[\Delta_p^r + \Delta_d^r] \leq [\Delta_p^{r-1} + \Delta_d^{r-1}], \quad (3.18)$$

which completes the induction. Moreover, the induction argument shows that if the stepsize  $\alpha$  satisfies the condition (3.17), then the descent condition (3.16) holds for all  $r \geq 1$ .

By the descent estimate (3.16), we have

$$\|x^{r+1} - x^r\| \rightarrow 0, \quad \|\nabla d(y^r)\| = \|E\bar{x}^r - q\| \rightarrow 0. \quad (3.19)$$

We now show that the sum of optimality gaps  $\Delta_d^r + \Delta_p^r$  in fact contracts geometrically after a finite number of ADMM iterations. By (3.19), for any  $\delta > 0$ , there must exist a finite integer  $\bar{r} > 0$  such that for all  $r \geq \bar{r}$ ,  $\|\nabla d(y^r)\| \leq \delta$ . Since  $\Delta_d^r, \Delta_p^r$  are nonnegative and bounded from above (see (3.18)), it follows that  $d(y^r)$  is bounded from below by a constant  $\zeta$  independent of  $r$ . Applying the second part of Lemma 2.3, we have that for all  $r \geq \bar{r}$ , the dual error bound  $\text{dist}(y^r, Y^*) \leq \tau\|\nabla d(y^r)\|$  holds true.

Therefore, it follows from Lemma 3.1 that we have the following cost-to-go estimate

$$\Delta_d^r = d^* - d(y^r) \leq \tau'\|\nabla d(y^r)\|^2 = \tau'\|E\bar{x}^r - q\|^2, \quad (3.20)$$

for some  $\tau' > 0$  and for all  $r \geq \bar{r}$ .

Moreover, we can use Lemma 3.1 to bound  $\|x^{r+1} - x^r\|^2$  from below by  $\Delta_p^r$ . In particular, we have from (3.15) and Lemma 3.1 that

$$\begin{aligned} \Delta_p^r &\leq \zeta\|x^{r+1} - x^r\|^2 + \zeta'\|\bar{x}^r - x^r\|^2 \\ &\leq \zeta\|x^{r+1} - x^r\|^2 + \zeta'\tau^2\sigma^2\|x^{r+1} - x^r\|^2 \\ &= \left(\zeta + \zeta'\tau^2\sigma^2\right)\|x^{r+1} - x^r\|^2. \end{aligned}$$

Substituting this bound and (3.20) into (3.16), and assuming that  $\alpha > 0$  satisfies (3.17), we obtain

$$\begin{aligned} [\Delta_p^r + \Delta_d^r] - [\Delta_p^{r-1} + \Delta_d^{r-1}] &\leq (\alpha\|E\|^2\tau^2\sigma^2 - \gamma)\|x^{r+1} - x^r\|^2 - \alpha\|E\bar{x}^r - q\|^2 \\ &\leq -\frac{(\gamma - \alpha\|E\|^2\tau^2\sigma^2)}{\zeta + \zeta'\tau^2\sigma^2}\Delta_p^r - \alpha(\tau')^{-1}\Delta_d^r \\ &\leq -\min\left\{\frac{(\gamma - \alpha\|E\|^2\tau^2\sigma^2)}{\zeta + \zeta'\tau^2\sigma^2}, \alpha(\tau')^{-1}\right\}[\Delta_p^r + \Delta_d^r]. \end{aligned}$$

Since  $\alpha > 0$  is chosen small enough such that (3.17) holds, we have

$$\lambda := \min \left\{ \frac{\gamma - \alpha \|E\|^2 \tau^2 \sigma^2}{\zeta + \zeta' \tau^2 \sigma^2}, \alpha (\tau')^{-1} \right\} > 0.$$

Consequently, we have

$$[\Delta_p^r + \Delta_d^r] - [\Delta_p^{r-1} + \Delta_d^{r-1}] \leq -\lambda [\Delta_p^r + \Delta_d^r]$$

which further implies

$$0 \leq [\Delta_p^r + \Delta_d^r] \leq \frac{1}{1 + \lambda} [\Delta_p^{r-1} + \Delta_d^{r-1}].$$

This shows that the sequence  $\{\Delta_p^r + \Delta_d^r\}_{r \geq \bar{r}}$  converges to zero Q-linearly<sup>1</sup>. As a result, we conclude that  $\{\Delta_p^r + \Delta_d^r\}$  and hence both  $\Delta_p^r$  and  $\Delta_d^r$  globally converge to zero R-linearly<sup>2</sup>.

We next show that the dual sequence  $\{y^r\}$  is also R-linearly convergent. To this end, notice that the inequalities (3.15) and (3.16) imply

$$[\Delta_p^r + \Delta_d^r] - [\Delta_p^{r-1} + \Delta_d^{r-1}] \leq (\alpha \|E\|^2 - \gamma \tau^{-2} \sigma^{-2}) \|x^r - \bar{x}^r\|^2 - \alpha \|E \bar{x}^r - q\|^2. \quad (3.21)$$

Then by (3.21), we see that both  $\|x^r - \bar{x}^r\| \rightarrow 0$  and  $\|E \bar{x}^r - q\| \rightarrow 0$  R-linearly. This implies that  $E x^r - q \rightarrow 0$  R-linearly and  $\nabla d(y^r) \rightarrow 0$  R-linearly. Using the fact that  $\text{dist}(y^r, Y^*) \leq \tau \|\nabla d(y^r)\|$ , we conclude that  $y^r$  converges R-linearly to an optimal dual solution.

We now argue that the primal iterates  $\{x^r\}$  converge to an optimal solution of (1.1). By the inequality (3.16), we can further conclude that

$$\|x^{r+1} - x^r\|^2 \rightarrow 0, \quad \|E \bar{x}^r - q\| \rightarrow 0$$

R-linearly. Notice that the R-linear convergence of  $\|x^{r+1} - x^r\|^2 \rightarrow 0$  implies that  $\|x^{r+1} - x^r\| \rightarrow 0$  R-linearly. This further shows that  $x^r \rightarrow x^\infty$  R-linearly for some  $x^\infty$ . Denote the limit of dual sequence  $\{y^r\}$  by  $y^\infty$ . By the preceding argument, we know  $y^\infty$  is a dual optimal solution of (1.1). To show that  $x^\infty$  is a primal optimal solution of (1.1), it suffices to prove that  $x^\infty \in X(y^\infty)$ . Using (3.15), and the fact that  $\|x^r - \bar{x}^r\| \rightarrow 0$ , we have

$$\|x^\infty - \bar{x}^r\| \leq \|x^r - x^\infty\| + \|x^r - \bar{x}^r\| \rightarrow 0.$$

Since  $\bar{x}^r \in X(y^r)$ , we have  $L(\bar{x}^r, y^r) \leq L(x, y^r)$  for all  $x \in X$ . Passing limit, we obtain  $L(x^\infty, y^\infty) \leq L(x, y^\infty)$  for all  $x \in X$ , that is,  $x^\infty \in X(y^\infty)$ . It then follows that the sequence  $\{x^r\}$  converges R-linearly to a primal optimal solution. **Q.E.D.**

<sup>1</sup>A sequence  $\{x^r\}$  is said to converge Q-linearly to some  $\bar{x}$  if  $\|x^{r+1} - \bar{x}\| / \|x^r - \bar{x}\| \leq \mu$  for all  $r$ , where  $\mu \in (0, 1)$  is some constant. A sequence  $\{x^r\}$  is said to converge to  $\bar{x}$  R-linearly if  $\|x^r - \bar{x}\| \leq c \mu^r$  for all  $r$  and for some  $c > 0$ .

<sup>2</sup>To see that such R-linear convergence is in fact global, note that  $\bar{r} > 0$  is finite, and  $\Delta_p^r + \Delta_d^r$  is Q-linearly convergent for  $r \geq \bar{r}$ . Then one can always find an appropriate constant  $c$  such that  $\Delta_p^r + \Delta_d^r \leq c(1 + \lambda)^{-r}$  for all  $r = 1, 2, \dots$

The following corollary relaxes the compactness assumption of the feasible set  $X$  (Assumption A-(g) in Theorem 3.1), and replaces it by the boundedness of the primal-dual iterates.

**Corollary 3.1** *Suppose assumptions A(a)—A(f) hold, and that  $X$  is a polyhedral set. Further assume that either one of the following two assumptions holds true*

1. *The sequence of dual iterates  $\{y^r\}$  lies in a compact set, and that the set  $\{x \mid L(x, y) \leq \zeta\}$  is compact for any finite  $y$  and  $\zeta$ .*
2. *The sequence of primal-dual iterates  $\{(x^r, y^r)\}$  lies in a compact set.*

*Then if  $\alpha$  is chosen sufficiently small (cf. (3.17)), all the conclusions stated in Theorem 3.1 still hold true. Moreover, the sequence of function values  $\{f(x^r)\}$  also converges linearly.*

**Proof.** Suppose the first assumption is true, so that all the dual iterates  $\{y^r\}$  lie in a compact set  $\tilde{Y}$ . Let us define

$$\delta := \sup \left\{ \|x\| \mid [L(x, y) - d(y)] + [d^* - d(y)] \leq \Delta_p^0 + \Delta_d^0, \forall y \in \tilde{Y} \right\}.$$

Clearly we have  $[d^* - d(y)] \geq 0$  and  $[L(x, y) - d(y)] \geq 0$  for all  $y \in \tilde{Y}$ . Moreover  $\delta < \infty$  due to the compactness of  $\tilde{Y}$  as well as the compactness of the set  $\{x \mid L(x, y) \leq \zeta\}$  for any finite  $y$ .

Again we show by induction that the sum of optimality gaps  $\Delta_d^r + \Delta_p^r$  is reduced after each ADMM iteration, as long as the stepsize  $\alpha$  is chosen sufficiently small. For any  $r \geq 1$ , by induction we suppose  $\Delta_d^{r-1} + \Delta_p^{r-1} \leq \Delta_d^0 + \Delta_p^0$ . Then  $\|x^r\| \leq \delta < \infty$ , and by the first part of Lemma 2.3, the primal error bound holds. Then we can carry out exactly the same analysis as the proof of Theorem 3.1 to arrive at the same conclusion.

Additionally, since

$$\begin{aligned} f(x^{r+1}) - d^* &= [f(x^{r+1}) - d(y^r)] + [d(y^r) - d^*] \\ &= [L(x^{r+1}; y^r) - d(y^r)] - [d^* - d(y^r)] - \langle y^r, q - Ex^{r+1} \rangle - \frac{\rho}{2} \|Ex^{r+1} - q\|^2 \\ &= \Delta_p^r - \Delta_d^r - \langle y^r, q - Ex^{r+1} \rangle - \frac{\rho}{2} \|Ex^{r+1} - q\|^2 \end{aligned}$$

and

$$\Delta_p^r \rightarrow 0, \quad \Delta_d^r \rightarrow 0, \quad \|q - Ex^r\| \rightarrow 0,$$

linearly, it follows that  $f(x^{r+1}) - d^* \rightarrow 0$  R-linearly (recall that  $y^r$  is now in a compact set). The proof is complete.

Similarly, when the second assumption is true, then the error bound condition in Lemma 2.3 again holds, and by using the same argument above we arrive at the desired conclusion. **Q.E.D.**

As a remark, we point out that the proof of Corollary 3.1 also shows that the same linear convergence of  $f(x^r) \rightarrow d^*$  also holds under the assumptions in Theorem 3.1.

We close this section by providing a few examples that satisfy the assumptions in Theorem 3.1 and Corollary 3.1. First consider the following  $\ell_1$  minimization problem

$$\min_x \|x\|_1, \quad \text{s.t. } Ex = b, \quad a \leq x_k \leq b, \quad k = 1, \dots, K \quad (3.22)$$

which can be equivalently written as a  $K$ -block problem

$$\min_{\{x_k\}} \sum_{k=1}^K |x_k|, \quad \text{s.t. } \sum_{k=1}^K e_k x_k = b, \quad a \leq x_k \leq b, \quad k = 1, \dots, K, \quad (3.23)$$

where  $e_k$  is the  $k$ -th column of  $E$ ,  $a$  and  $b$  are some scalars. It is easy to verify that this problem meets all the conditions listed in Assumption A, hence the linear convergence result in Theorem 3.1 applies. The same is true for the following mixed  $\ell_1/\ell_2$  minimization problem

$$\min_{\{x_k\}} \sum_{k=1}^K \|x_k\|, \quad \text{s.t. } \sum_{k=1}^K E_k x_k = b, \quad a \leq x_k \leq b, \quad k = 1, \dots, K, \quad (3.24)$$

where  $x_k$ ,  $a$  and  $b$  are now  $n$ -dimensional vectors, and  $E_k \in \Re^{m \times n}$  is some matrix with full column rank.

Furthermore, the boundedness assumption in Corollary 3.1 is satisfied by many examples of the two-block ADMM described in [7]. This is because when  $K = 2$ ,  $\alpha/\beta \in (0, \frac{1}{2}(1 + \sqrt{5}))$  and  $E_k$ 's are full column rank, it is known that both the primal and dual iterates generated by the two-block ADMM algorithm indeed lie in a bounded set [19]. Therefore the second assumption made in the Corollary 3.1 holds true, hence we only require assumptions A(a)–A(f), which are in fact quite mild. For example, they are met by the following instance of the consensus problem (see [7, Section 7] for introduction of the consensus problem)

$$\min_{\{x_k\}, z} \sum_{k=1, \dots, K} \|Ax_k - b\|^2 + w \|x_k\|_1 \quad \text{s.t. } x_k - z = 0, \quad k = 1, \dots, K, \quad (3.25)$$

where  $w > 0$  is some constant. Thus, the two block ADMM algorithm converges linearly for (3.25) regardless of the rank of  $A$ . Note that when  $A$  is full column rank, the objective is strongly convex. Consequently the error bound condition in Lemma 2.3 holds true globally, and the coefficient  $\tau$  can be at least greater than  $\lambda_{\min}(A^T A)$ .



## 4 Variants of ADMM

The convergence analysis of Section 3 can be extended to some variants of the ADMM. We briefly describe two of them below.

### 4.1 Proximal ADMM

In the original ADMM (1.7), each block  $x_k$  is updated by solving a convex optimization subproblem *exactly*. For large scale problems, this subproblem may not be easy to solve unless the matrix  $E_k$  is unitary (i.e.,  $E_k^T E_k = I$ ) in which case the variables in  $x_k$  can be further decoupled (assuming  $f_k$  is separable). If the matrix  $E_k$  is not unitary, we can still employ a simple proximal gradient step to *inexactly* minimize  $L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k, x_{k+1}^r, \dots, x_K^r)$ . More specifically, we update each block of  $x_k$  according to the following procedure

$$\begin{aligned} x_k^{r+1} = \operatorname{argmin}_{x_k} & \left\{ h_k(x_k) + \langle y^r, q - E_k x_k \rangle + \langle A_k^T \nabla g_k(A_k x_k^r), x_k - x_k^r \rangle + \frac{\beta}{2} \|x_k - x_k^r\|^2 \right. \\ & \left. + \left\langle \rho E_k^T \left( \sum_{j < k} E_j x_j^{r+1} + \sum_{j \geq k} E_j x_j^r - q \right), x_k - x_k^r \right\rangle \right\} \end{aligned} \quad (4.1)$$

in which the smooth part of the objective function in the  $k$ -th subproblem, namely,

$$g_k(A_k x_k) + \langle y^r, q - E_k x_k \rangle + \frac{\rho}{2} \left\| E_k x_k + \sum_{j < k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right\|^2$$

is linearized locally at  $x_k^r$ , and a proximal term  $\frac{\beta}{2} \|x_k - x_k^r\|^2$  is added. Here,  $\beta > 0$  is a positive constant. With this change, updating  $x_k$  is easy when  $h_k$  (the nonsmooth part of  $f_k$ ) is separable. For example, this is the case for compressive sensing applications where  $h_k(x_k) = \|x_k\|_1$ , and the resulting subproblem admits a closed form solution given by the component-wise soft thresholding (also known as the shrinkage operator). We note that the proximal ADMM algorithm described here is slightly more general than the proximal ADMM algorithm seen in the literature, in which only the penalization term  $\frac{\rho}{2} \left\| E_k x_k + \sum_{j < k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r - q \right\|^2$  is linearized locally at  $x_k^r$ ; see e.g., [54, 55].

We claim that Theorem 3.1 holds for the proximal ADMM algorithm *without requiring assumption A-(f)* (the full rankness of  $E_k$ 's). Indeed, to establish the (linear) convergence of the proximal ADMM (4.1), we can follow the same proof steps as that for Theorem 3.1, with the only changes being in the proof of Lemmas 2.4-2.5 and Lemma 3.1. We first show that Lemma 2.4 holds without assumption A(f). Clearly subproblem (4.1) is now *strongly convex* without the full column rank assumption of  $E_k$ 's made in A(f). In the following, we will show that as long as  $\beta$  is large enough,

there is a sufficient descent:

$$L(x^{r+1}; y^r) - L(x^r; y^r) \leq -\gamma \|x^{r+1} - x^r\|^2, \quad \text{for some } \gamma > 0 \text{ independent of } y^r. \quad (4.2)$$

This property can be seen by bounding the smooth part of  $L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k, x_{k+1}^r, \dots, x_K^r)$ , which is given by

$$\bar{L}_k(x_k) := g_k(A_k x_k) + \langle y^r, q - E_k x_k \rangle + \frac{\rho}{2} \left\| \sum_{j < k} E_j x_j^{r+1} + \sum_{j > k} E_j x_j^r + E_k x_k - q \right\|^2,$$

with the Taylor expansion at  $x_k^r$ :

$$\bar{L}_k(x_k^{r+1}) \leq \bar{L}_k(x_k^r) + \langle \nabla \bar{L}_k(x_k^r), x_k^{r+1} - x_k^r \rangle + \frac{\nu}{2} \|x_k^{r+1} - x_k^r\|^2 \quad (4.3)$$

where

$$\nu := L \|A_k\| \|A_k^T\| + \rho \|E_k^T E_k\|$$

is the Lipschitz constant of  $\bar{L}_k(\cdot)$  and  $L$  is the Lipschitz constant of  $\nabla g_k(\cdot)$ . Making the above inequality more explicit yields

$$\begin{aligned} & L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k^{r+1}, x_{k+1}^r, \dots, x_K^r; y^r) - L(x_1^{r+1}, \dots, x_{k-1}^{r+1}, x_k^r, x_{k+1}^r, \dots, x_K^r; y^r) \\ & \leq h_k(x_k^{r+1}) - h_k(x_k^r) + \langle y^r, E_k(x_k^r - x_k^{r+1}) \rangle + \langle A_k^T \nabla g_k(A_k x_k^r), x_k^{r+1} - x_k^r \rangle \\ & \quad + \left\langle \rho E_k^T \left( \sum_{j < k} E_j x_j^{r+1} + \sum_{j \geq k} E_j x_j^r - q \right), x_k^{r+1} - x_k^r \right\rangle + \frac{\nu}{2} \|x_k^{r+1} - x_k^r\|^2 \\ & \leq -\frac{\beta}{2} \|x_k^{r+1} - x_k^r\|^2 + \frac{\nu}{2} \|x_k^{r+1} - x_k^r\|^2 \\ & = -\gamma \|x_k^{r+1} - x_k^r\|^2, \quad \forall k, \end{aligned} \quad (4.4)$$

provided the regularization parameter  $\beta$  satisfies

$$\gamma := \frac{1}{2} (\beta - \nu) > 0.$$

In the above derivation of (4.4), the first step is due to (4.3), while the second inequality follows from the definition of  $x_k^{r+1}$  (cf. (4.1)). Summing (4.4) over all  $k$  yields the desired estimate of sufficient descent (4.2).

To verify that Lemma 2.5 still holds for the proximal ADMM algorithm, we note from the corresponding optimality condition for (4.1)

$$x_k^{r+1} = \text{prox}_{h_k} \left[ x_k^{r+1} - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r - \rho E_k^T \left( \sum_{j < k} E_j x_j^{r+1} + \sum_{j \geq k} E_j x_j^r - q \right) - \beta (x_k^{r+1} - x_k^r) \right].$$

Using this relation in place of (2.10) and following the same proof steps, we can easily prove that the bound (2.9) in Lemma 2.5 can be extended to the proximal ADMM algorithm. Thus, the convergence results in Theorem 3.1 remain true for the proximal ADMM algorithm (4.1).

It remains to verify that Lemma 3.1 still holds true. In fact the first part of Lemma 3.1 can be shown to be independent of the iterates, thus it trivially holds true for the proximal ADMM algorithm. To show that the second part of Lemma 3.1 is true, note that the optimality condition of the proximal ADMM algorithm implies that

$$\begin{aligned} x_k^{r+1} &= \text{prox}_{h_k} \left[ x_k^{r+1} - \nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}, \{x_j^r\}_{j \geq k}; y^r \right) - \beta(x_k^{r+1} - x_k^r) \right] \\ &:= \text{prox}_{h_k} \left[ x_k^r - \nabla_{x_k} \bar{L}(x^r; y^r) - e_k^r \right] \end{aligned}$$

where in this case  $e_k^r$  is given as

$$e_k^r := x_k^r - x_k^{r+1} + \nabla_{x_k} \bar{L} \left( \{x_j^{r+1}\}, \{x_j^r\}_{j \geq k}; y^r \right) - \nabla_{x_k} \bar{L}(x^r; y^r) + \beta(x_k^{r+1} - x_k^r).$$

It is then straightforward to show that the norm of  $e_k^r$  can be bounded by  $c' \|x^r - x^{r+1}\|$  for some constant  $c' > 0$ . The rest of the proof follows the same steps as in Lemma 3.1.

## 4.2 Jacobi Update

Another popular variant of the ADMM algorithm is to use a Jacobi iteration (instead of a Gauss-Seidel iteration) to update the primal variable blocks  $\{x_k\}$ . In particular, the ADMM iteration (1.7) is modified as follows:

$$x_k^{r+1} = \underset{x_k}{\text{argmin}} \left( h_k(x_k) + g_k(A_k x_k) - \langle y^r, E_k x_k \rangle + \frac{\rho}{2} \left\| E_k x_k + \sum_{j \neq k} E_j x_j^r - q \right\|^2 \right), \quad \forall k. \quad (4.5)$$

The convergence for this direct Jacobi scheme is unclear, as the augmented Lagrangian function may not decrease after each Jacobi update. In the following, we consider a modified Jacobi scheme with an explicit stepsize control. Specifically, let us introduce an intermediate variable  $w = (w_1^T, \dots, w_K^T)^T \in \mathfrak{R}^n$ . The modified Jacobi update is given as follows:

$$w_k^{r+1} = \underset{x_k}{\text{argmin}} \left( h_k(x_k) + g_k(A_k x_k) - \langle y^r, E_k x_k \rangle + \frac{\rho}{2} \left\| E_k x_k + \sum_{j \neq k} E_j x_j^r - q \right\|^2 \right), \quad \forall k, \quad (4.6)$$

$$x_k^{r+1} = x_k^r + \frac{1}{K} (w_k^{r+1} - x_k^r), \quad \forall k. \quad (4.7)$$

where a stepsize of  $1/K$  is used in the update of each variable block.

With this modification, we claim that Lemmas 2.4-2.5 and Lemma 3.1 still hold. In particular, Lemma 2.4 can be argued as follows. The strong convexity of  $L(x; y)$  with respect to the variable block  $x_k$  implies that

$$\begin{aligned} & L(x_1^r, \dots, x_{k-1}^r, x_k^r, x_{k+1}^r \dots, x_K^r; y^r) - L(x_1^r, \dots, x_{k-1}^r, w_k^r, x_{k+1}^r \dots, x_K^r; y^r) \\ & \geq \gamma \|w_k^{r+1} - x_k^r\|^2, \quad \forall k. \end{aligned}$$

Using this inequality we obtain

$$\begin{aligned} & L(x^r; y^r) - L(x^{r+1}; y^r) \\ & = L(x^r; y^r) - L\left(\frac{K-1}{K}x^r + \frac{1}{K}w^{r+1}; y^r\right) \\ & = L(x^r; y^r) - L\left(\frac{1}{K}\sum_{k=1}^K(x_1^r, \dots, x_{k-1}^r, w_k^{r+1}, x_{k+1}^r \dots, x_K^r); y^r\right) \\ & \geq L(x^r; y^r) - \frac{1}{K}\sum_{k=1}^K L(x_1^r, \dots, x_{k-1}^r, w_k^{r+1}, x_{k+1}^r \dots, x_K^r; y^r) \\ & = \frac{1}{K}\sum_{k=1}^K (L(x^r; y^r) - L(x_1^r, \dots, x_{k-1}^r, w_k^{r+1}, x_{k+1}^r \dots, x_K^r; y^r)) \\ & \geq \frac{\gamma}{K}\sum_{k=1}^K \|w_k^{r+1} - x_k^r\|^2 \\ & = \frac{\gamma}{K}\|w^{r+1} - x^r\|^2. \end{aligned}$$

where the first inequality comes from the convexity of the augmented Lagrangian function.

From the update rule (4.7) we have  $K(x_k^{r+1} - x_k^r) = (w_k^{r+1} - x_k^r)$ , which combined with the previous inequality yields

$$L(x^r; y^r) - L(x^{r+1}; y^r) \geq \gamma K \|x^{r+1} - x^r\|^2.$$

The proof of Lemma 2.5 also requires only minor modifications. In particular, we have the following optimality condition for (4.5)

$$w_k^{r+1} = \text{prox}_{h_k} \left[ w_k^{r+1} - A_k^T \nabla_{x_k} g_k(A_k w_k^{r+1}) + E_k^T y^r - \rho E_k^T \left( \sum_{j \neq k} E_j x_j^r + E_k w_k^{r+1} - q \right) \right]$$

Similar to the proof of Lemma 2.5, we have

$$\|w_k^{r+1} - \text{prox}_{h_k} [x_k^r - A_k^T \nabla_{x_k} g_k(A_k x_k^r) + E_k^T y^r - \rho E_k^T (E x^r - q)]\| \leq c \|w^{r+1} - x^r\|.$$

Utilizing the relationship  $K(x_k^{r+1} - x_k^r) = (w_k^{r+1} - x_k^r)$ , we can establish Lemma 2.5 by following similar proof steps (which we omit due to space reason).

Lemma 3.1 can be shown as follows. We first express  $w_k^{r+1}$  as

$$\begin{aligned} w_k^{r+1} &= \text{prox}_{h_k} [w_k^{r+1} - \nabla_{x_k} \bar{L}(\{x_{j \neq k}^r\}, w_k^{r+1}; y^r)] \\ &= \text{prox}_{h_k} [x_k^r - \nabla_{x_k} \bar{L}(x^r; y^r) - e_k^r] \end{aligned}$$

where we have defined

$$e_k^r := \nabla_{x_k} \bar{L}(\{x_{j \neq k}^r\}, w_k^{r+1}; y^r) - \nabla_{x_k} \bar{L}(x^r; y^r) + x_k^r - w_k^{r+1}.$$

Again by using the relationship  $K(x_k^{r+1} - x_k^r) = (w_k^{r+1} - x_k^r)$ , we can bound the norm of  $e_k^r$  by  $c' \|x^{r+1} - x^r\|$ , for some  $c' > 0$ . The remaining proof steps are similar to those in Lemma 3.1.

Since Lemmas 2.4-2.5 and Lemma 3.1 hold for the Jacobi version of the ADMM algorithm with a step size control, we conclude that the convergence results of Theorem 3.1 remain true in this case.

## 5 Concluding Remarks

In this paper we have established the convergence and the rate of convergence of the classical ADMM algorithm when the number of variable blocks are more than two and in the absence of strong convexity. Our analysis is a departure of the conventional analysis of ADMM algorithm which relies on a contraction argument involving a weighted (semi-)norm of  $(x^r - x^*, y^r - y^*)$ , see [17–20, 25, 26, 30, 47]. In our analysis, we require neither the strong convexity of the objective function nor the row independence assumption of the constrained matrix  $E$ . Instead, we use a local error bound to show that when the stepsize of dual update is made sufficiently small, the sum of the primal and the dual optimality gaps decreases after each ADMM iteration, although separately they may individually increase. An interesting issue for further research is to identify good practical stepsize rules for dual update. While (3.17) does suggest a dual stepsize rule in terms of error bound constants, it may be too conservative and cumbersome to compute unless the objective function is strongly convex. One possibility may be to use an adaptive dual stepsize rule to guarantee the decrease of the sum of the primal and dual optimality gaps.

**Acknowledgement:** The authors are grateful to Xiangfeng Wang and Dr. Min Tao of Nanjing University for their constructive comments.

## 6 Appendix

### 6.1 Proof of Dual Error Bound (2.5)

The augmented Lagrangian dual function can be expressed as

$$d(y) = \min_{x \in X} \langle y, q - Ex \rangle + \frac{\rho}{2} \|q - Ex\|^2 + g(Ax) + h(x). \quad (6.1)$$

For convenience, define  $p(Ex) := \frac{\rho}{2} \|q - Ex\|^2$ , and let  $\ell(x) := p(Ex) + g(Ax) + h(x)$ . For simplicity, in this proof we further restrict ourselves to the case where the nonsmooth part has polyhedral level sets, i.e.,  $\{x : h(x) \leq \xi\}$  is polyhedral for each  $\xi$ . More general cases can be shown along similar lines, but the arguments become more involved.

Let us define

$$x(y) \in \arg \min_{x \in X} \ell(x) + \langle y, q - Ex \rangle.$$

Let  $(x^*, y^*)$  denote a primal and dual optimal solution pair. Let  $X^*$  and  $Y^*$  denote the primal and dual optimal solution set. The following properties will be useful in our subsequent analysis.

- (a) There exist positive scalars  $\sigma_g, L_g$  such that  $\forall x(y), x(y') \in X$ 
  - a-1)  $\langle A^T \nabla g(Ax(y')) - A^T \nabla g(Ax(y)), x(y') - x(y) \rangle \geq \sigma_g \|Ax(y') - Ax(y)\|^2$ .
  - a-2)  $g(Ax(y')) - g(Ax(y)) - \langle A^T \nabla g(Ax(y)), x(y') - x(y) \rangle \geq \frac{\sigma_g}{2} \|Ax(y') - Ax(y)\|^2$ .
  - a-3)  $\|A^T \nabla g(Ax(y')) - A^T \nabla g(Ax(y))\| \leq L_g \|Ax(y') - Ax(y)\|$ .
- b) All a-1)–a-3) are true for  $p(\cdot)$  as well, with some constants  $\sigma_p$  and  $L_p$ .
- c)  $\nabla d(y) = q - Ex(y)$ , and  $\|\nabla d(y') - \nabla d(y)\| \leq \frac{1}{\rho} \|y' - y\|$ .

Part (a) is true due to the assumed Lipschitz continuity and strong convexity of the function  $g(\cdot)$ . Part (b) is from the Lipschitz continuity and strong convexity of the quadratic penalization  $p(\cdot)$ . Part (c) has been shown in Lemma 2.1 and Lemma 2.2.

To proceed, let us rewrite the primal problem equivalently as

$$d(y) = \min_{(x,s): x \in X, h(x) \leq s} \langle y, q - Ex \rangle + p(Ex) + g(Ax) + s. \quad (6.2)$$

Let us write the polyhedral set  $\{(x, s) : x \in X, h(x) \leq s\}$  compactly as  $C_x x + C_s s \geq c$  for some matrices  $C_x \in \mathbb{R}^{j \times n}$ ,  $C_s \in \mathbb{R}^{j \times 1}$  and  $c \in \mathbb{R}^{j \times 1}$ , where  $j$  is some integer. For any fixed  $y$ , let  $(x(y), s(y))$  denote one optimal solution for (6.2), note we must have  $h(x(y)) = s(y)$ . Due to equivalence, if  $y^* \in Y^*$ , we must also have  $x(y^*) \in X^*$ .

Define a set-valued function  $\mathcal{M}$  that assigns the vector  $(d, e) \in \mathbb{R}^n \times \mathbb{R}^m$  to the set of vectors  $(x, s, y, \lambda) \in \mathbb{R}^n \times \mathbb{R} \times \mathbb{R}^m \times \mathbb{R}^j$  that satisfy the following system of equations

$$\begin{aligned} E^T y + C_x^T \lambda &= d, \\ C_s^T \lambda &= 1, \\ q - Ex &= e, \\ \lambda &\geq 0, \quad (C_x x + C_s s) \geq c, \quad \langle C_x x + C_s s - c, \lambda \rangle = 0. \end{aligned}$$

It is easy to verify by using the optimality condition for problem (6.2) that

$$\begin{aligned} (x, s, y, \lambda) \in \mathcal{M}(E^T \nabla p(Ex) + A^T \nabla g(Ax), e) \text{ for some } \lambda \\ \text{if and only if } x = x(y), \quad e = \nabla d(y). \end{aligned} \tag{6.3}$$

We can take  $e = 0$ , and use the fact that  $x(y^*) \in X^*$ , we see that  $(x, s, y, \lambda) \in \mathcal{M}(E^T \nabla p(Ex) + A^T \nabla g(Ax), 0)$  if and only if  $x \in X^*$  and  $y \in Y^*$ .

The following result states a well-known local upper Lipschitzian continuity property for the polyhedral multifunction  $\mathcal{M}$ ; see [28, 36, 37].

**Proposition 6.1** *There exists a positive scalar  $\theta$  that depends on  $A, E, C_x, C_s$  only, such that for each  $(\bar{d}, \bar{e})$  there is a positive scalar  $\delta'$  satisfying*

$$\mathcal{M}(d, e) \subseteq \mathcal{M}(\bar{d}, \bar{e}) + \theta \|(d, e) - (\bar{d}, \bar{e})\| \mathcal{B}, \tag{6.4}$$

$$\text{whenever } \|(d, e) - (\bar{d}, \bar{e})\| \leq \delta'. \tag{6.5}$$

where  $\mathcal{B}$  denotes the unit Euclidean ball in  $\mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R} \times \mathbb{R}^j$ .

The following is the main result for this appendix. Note that the scalar  $\tau$  in the claim is independent the choice of  $y, x, s$ , and is independent on the coefficients of the linear term  $s$ .

**Claim 6.1** *Suppose all the assumptions in Assumption A are satisfied. Then there exists positive scalars  $\delta, \tau$  such that for all  $y \in \mathcal{U}$  and  $\|\nabla d(y)\| \leq \delta$ , there holds  $\text{dist}(y, Y^*) \leq \tau \|\nabla d(y)\|$ .*

**Proof.** By the previous claim,  $\mathcal{M}$  is locally Lipschitzian with modulus  $\theta$  at  $(\nabla \ell(x^*), 0) = (E^T \nabla p(Ex^*) + A^T \nabla g(Ax^*), 0)$ .

Let  $\delta \leq \delta'/2$ . We first show that if  $\|\nabla d(y)\| \leq \delta$ , then we must have  $\|\nabla \ell(x(y)) - \nabla \ell(x^*)\| \leq \delta'/2$ . To this end, take a sequence  $y^1, y^2, \dots$ , such that  $e^r := \nabla d(y^r) \rightarrow 0$ . By assumption A(g)  $\{x(y^r)\}$  lies in a compact set. Due to the fact that  $s(y^r) = h(x(y^r))$ , so the sequence  $\{s(y^r)\}$  also lies in

a compact set (cf. Assumption A(e)). By passing to a subsequence if necessary, let  $(x^\infty, s^\infty)$  be a cluster point of  $\{x(y^r), s(y^r)\}$ . In light of the continuity of  $\nabla\ell(\cdot)$ , we have  $(\nabla\ell(x(y^r)), e^r) \rightarrow (\nabla\ell(x^\infty), 0)$ . Now for all  $r$ ,  $\{(x(y^r), s(y^r), \nabla\ell(x(y^r)), e^r)\}$  lies in the set

$$\{(x, s, d, e) \mid (x, s, y, \lambda) \in \mathcal{M}(d, e) \text{ for some } (y, \lambda)\}$$

which is polyhedral and thus is closed. Then we can pass limit to it and conclude (cf. Proposition 6.1)

$$(x^\infty, s^\infty, y^\infty, \lambda^\infty) \in \mathcal{M}(\nabla\ell(x^\infty), 0)$$

for some  $(y^\infty, \lambda^\infty) \in \mathbb{R}^m \times \mathbb{R}^j$ . Thus by (6.3) and the discussions that follow, we have  $x^\infty \in X^*$  and  $y^\infty \in Y^*$ . By Lemma 2.1, we have  $\nabla\ell(x^*) = \nabla\ell(x^\infty)$ , which further implies that  $\nabla\ell(x(y^r)) \rightarrow \nabla\ell(x^*)$ . This shows that the desired  $\delta$  exists.

Then we let  $e = \nabla d(y)$ , and suppose  $\|e\| \leq \delta$ . From the previous argument we have

$$\|\nabla\ell(x(y)) - \nabla\ell(x^*)\| + \|e\| \leq \delta'/2 + \delta'/2 = \delta'.$$

Using the results in Proposition 6.1, we have that there exists  $(x^*, s^*, y^*, \lambda^*) \in \mathcal{M}(\nabla\ell(x^*), 0)$  satisfying

$$\|(x(y), s, y, \lambda) - (x^*, s^*, y^*, \lambda^*)\| \leq \theta (\|\nabla\ell(x^*) - \nabla\ell(x(y))\| + \|e\|).$$

Since  $(x(y), s, y, \lambda) \in \mathcal{M}(\nabla\ell(x(y)), e)$ , it follows from the definition of  $\mathcal{M}$  that

$$E^T y + C_x^T \lambda = \nabla\ell(x(y)), \quad (6.6)$$

$$C_s^T \lambda = 1, \quad (6.7)$$

$$q - Ex(y) = e, \quad (6.8)$$

$$\lambda \geq 0, \quad (C_x x(y) + C_s s(y)) \geq c, \quad \langle C_x x(y) + C_s s(y) - c, \lambda \rangle = 0. \quad (6.9)$$

Since  $(x^*, s^*, y^*, \lambda^*) \in \mathcal{M}(\nabla\ell(x^*), 0)$ , we have from the definition of  $\mathcal{M}$

$$E^T y^* + C_x^T \lambda^* = \nabla\ell(x^*), \quad (6.10)$$

$$C_s^T \lambda^* = 1, \quad (6.11)$$

$$q - Ex^* = 0, \quad (6.12)$$

$$\lambda^* \geq 0, \quad (C_x x^* + C_s s^*) \geq c, \quad \langle C_x x^* + C_s s^* - c, \lambda^* \rangle = 0. \quad (6.13)$$

Moreover, we have

$$\begin{aligned} & \sigma_g \|A(x(y) - x^*)\|^2 + \sigma_p \|E(x(y) - x^*)\|^2 \\ & \leq \langle A^T \nabla g(Ax(y)) - A^T \nabla g(Ax(y^*)), x(y) - x(y^*) \rangle + \langle E^T \nabla p(Ex(y)) - E^T \nabla p(Ex(y^*)), x(y) - x(y^*) \rangle \\ & = \langle \nabla\ell(x(y)) - \nabla\ell(x(y^*)), x(y) - x(y^*) \rangle \\ & = \langle \lambda - \lambda^*, C_x x(y) - C_x x^* \rangle + \langle y - y^*, Ex(y) - Ex^* \rangle \end{aligned}$$



where the first inequality comes from the strong convexity of  $g(\cdot)$  and  $p(\cdot)$ ; the last equality is from (6.6) and (6.10). Moreover, we have

$$\begin{aligned}
& \langle \lambda - \lambda^*, C_x x(y) - C_x x^* \rangle \\
&= \langle \lambda - \lambda^*, C_x x(y) - C_x x^* \rangle + \langle \lambda - \lambda^*, C_s s - C_s s^* \rangle \\
&= \langle \lambda - \lambda^*, (C_x x(y) + C_s s) - (C_x x^* + C_s s^*) \rangle \\
&= -\langle \lambda^*, C_x x(y) + C_s s - c \rangle - \langle \lambda, C_x x^* + C_s s^* - c \rangle \leq 0
\end{aligned} \tag{6.14}$$

where in the first equality we have used the fact that  $C_s^T \lambda - C_s^T \lambda^* = 0$ ; see (6.7) (6.11); in the third equality and in the last inequality we have used the complementary conditions (6.13) and (6.9). As a result, we have

$$\begin{aligned}
& \sigma_g \|A(x(y) - x^*)\|^2 + \sigma_p \|E(x(y) - x^*)\|^2 \\
& \leq \langle y - y^*, (Ex(y) - q) - (Ex^* - q) \rangle \leq \|y - y^*\| \|e\|,
\end{aligned} \tag{6.15}$$

where the last step is due to  $\nabla d(y) = Ex(y) - q$  and  $\nabla d(y^*) = Ex^* - q = 0$ . Finally we have from Proposition 6.1

$$\begin{aligned}
& \|(x(y), s, y, \lambda) - (x^*, s^*, y^*, \lambda^*)\|^2 \\
& \leq \theta^2 (\|\nabla \ell(x^*) - \nabla \ell(x(y))\| + \|e\|)^2 \\
& \leq \theta^2 (2\|\nabla \ell(x^*) - \nabla \ell(x(y))\|^2 + 2\|e\|^2) \\
& \leq 2\theta^2 (2\|\nabla g(x^*) - \nabla g(x(y))\|^2 + 2\|\nabla p(x^*) - \nabla p(x(y))\|^2 + \|e\|^2) \\
& \leq 2\theta^2 (L_g^2 \|A^T(x(y) - x^*)\|^2 + L_p^2 \|E^T(x(y) - x^*)\|^2 + \|e\|^2) \\
& \leq 2\theta^2 \max\left(\frac{2L_g^2}{\sigma_g}, \frac{2L_p^2}{\sigma_p}, 1\right) (\sigma_g \|A^T(x(y) - x^*)\|^2 + \sigma_p \|E^T(x(y) - x^*)\|^2 + \|e\|^2) \\
& \leq 2\theta^2 \max\left(\frac{2L_g^2}{\sigma_g}, \frac{2L_p^2}{\sigma_p}, 1\right) (\|e\| \|y - y^*\| + \|e\|^2) \\
& \leq 2\theta^2 \max\left(\frac{2L_g^2}{\sigma_g}, \frac{2L_p^2}{\sigma_p}, 1\right) (\|e\| \|(x(y), s, y, \lambda) - (x^*, s^*, y^*, \lambda^*)\| + \|e\|^2),
\end{aligned}$$

where the second inequality is due to  $\nabla \ell(x) = \nabla g(x) + \nabla p(x)$  and the fourth step follows from properties a-3) and b).

We see that the above inequality is quadratic in  $\|(x(y), s, y, \lambda) - (x^*, s^*, y^*, \lambda^*)\|/\|e\|$ , so we have

$$\|(x(y), s, y, \lambda) - (x^*, s^*, y^*, \lambda^*)\|/\|e\| \leq \tau$$

for some scalar  $\tau$  depending on  $\theta, L_g, L_p, \sigma_g, \sigma_p$ . It is worth noting that  $\tau$  does not depend on the choice of the coefficients of the linear term  $s$ . We conclude  $\text{dist}(y, Y^*) \leq \tau \|\nabla d(y)\|$ . **Q.E.D.**

## References

- [1] Bertsekas, D.P.: Nonlinear Programming. Athena Scientific (2010).
- [2] Bertsekas, D. P. and Gafni, E.: Projection methods for variational inequalities with application to the traffic assignment problem. *Math. Prog. Study.* 17, 139–159 (1982)
- [3] Bertsekas, D. P. and Gallager, R.: *Data Networks*. Prentice–Hall, Englewood Cliffs, New Jersey. (1987)
- [4] Bertsekas, D. P., Hosein, P. A., and Tseng, P.: Relaxation methods for network flow problems with convex arc costs. *SIAM J. Control Optim.* 25, 1219–1243 (1987)
- [5] Bertsekas, D. P. and Tsitsiklis, J. N.: *Parallel and Distributed Computation: Numerical Methods*. Prentice–Hall, Englewood Cliffs, New Jersey (1989)
- [6] Boley, D.: Linear convergence of ADMM on a model problem. TR 12-009, Department of Computer Science and Engineering, University of Minnesota (2012)
- [7] Boyd, S., Parikh, N., Chu, E., Peleato, B. and Eckstein, J.: Distributed Optimization and Statistical Learning via the Alternating Direction Method of Multipliers. *Foundations and Trends in Machine Learning*, Michael Jordan, Editor in Chief, 3, 1–122 (2011)
- [8] Bregman, L. M.: The relaxation method of finding the common point of convex sets and its application to the solution of problems in convex programming. *USSR Comp. Math. and Math. Physics* 7, 200–217 (1967)
- [9] Chandrasekaran, V., Parrilo, P., and Willsky, A.: Latent variable graphical model via convex optimization. Preprint. (2010)
- [10] Cottle, R. W. Duvall, S. G., and Zikan, K.: A Lagrangian relaxation algorithm for the constrained matrix problem. *Naval Res. Logistics Quarterly.* 33, 55–76 (1986)
- [11] De Pierro, A. R. and Iusem, A. N.: On the convergence properties of Hildreth’s quadratic programming algorithm. *Math. Prog.* 47, 37–51 (1990)
- [12] Deng, W. and Yin, W.: On the global linear convergence of alternating direction methods. Preprint. (2012)
- [13] Douglas, J. and Rachford, H.H.: On the numerical solution of the heat conduction problem in 2 and 3 space variables. *Transactions of the American Mathematical Society.* 82, 421-439 (1956)

- [14] Eckstein, J.: Splitting methods for monotone operators with applications to parallel optimization. Ph.D Thesis, Operations Research Center, MIT. (1989)
- [15] Eckstein, J. and Bertsekas, D.P.: On the Douglas-Rachford splitting method and the proximal point algorithm for maximal monotone operators. *Mathematical Programming.*55, 293-318 (1992)
- [16] Eckstein, J. and Svaiter, B.F.: General projective splitting methods for sums of maximal monotone operators. *SIAM Journal on Control and Optimization.* 48, 787–811 (2010)
- [17] Gabay, D.: Application of the method of multipliers to variational inequalities, In: Fortin, M., Glowinski, R., eds., *Augmented Lagrangian methods: Application to the numerical solution of Boundary-Value Problem*, North-Holland, Amsterdam, The Netherlands. 299-331 (1983)
- [18] Gabay, D. and Mercier, B.: A dual algorithm for the solution of nonlinear variational problems via finite-element approximations. *Computer and Mathematics with Applications.* 2, 17-40 (1976)
- [19] Glowinski, R.: *Numerical methods for nonlinear variational problems.* Springer-Verlag, New York, Berlin, Heidelberg, Tokyo (1984)
- [20] Glowinski, R. and Le Tallec, P.: *Augmented Lagrangian and operator splitting methods in nonlinear mechanics.* SIAM Studies in Applied Mathematics, Philadelphia, PA. (1989)
- [21] Goldfarb, D. and Ma, S.: Fast multiple splitting algorithms for convex optimization. Arxiv preprint arXiv:0912.4570 (2009)
- [22] Goldfarb, D., Ma, S. and Scheinberg, K.: Fast alternating linearization methods for minimizing the sum of two convex functions. Arxiv preprint arXiv:0912.4571 (2009)
- [23] Goldstein, A. A.: Convex programming in hilbert space. *Bull. Am. Math. Soc.* 70, 709–710 (1964)
- [24] Goldstein, T., O’Donoghue, B. and Setzer, S.: Fast alternating direction optimization methods. CAM report 12-35, UCLA, (2012).
- [25] He, B. S., Tao, M. and Yuan, X.M.: Alternating direction method with gaussian back substitution for separable convex programming. *SIAM J. Optim.* 22, 313-340 (2012)
- [26] He, B. S. and Yuan, X. M.: On the  $O(1/n)$  convergence rate of the Douglas-Rachford alternating direction method. *SIAM J. Numer. Anal.* 50, 700-709 (2012)

- [27] Herman, G. T.: Image Reconstruction from Projection: The Fundamentals of Computerized Tomography. Academic Press, New York. (1980)
- [28] Hoffman, A. J.: On Approximate Solutions of Systems of Linear Inequalities. J. Res. Nat. Bur. Standards. 49, 263-265 (1952)
- [29] Iusem, A. N.: On dual convergence and the rate of primal convergence of bregman's convex programming method. SIAM J. Control Optim. 1, 401-423 (1991)
- [30] Kontogiorgis, S. and Meyer, R. R.: A variable-penalty alternating directions method for convex optimization, Mathematical Programming. 83, 29-53 (1998)
- [31] Lamond, B., and Stewart, N. F.: Bregman's balancing method. Transportation Res. 15B, 239-248 (1981)
- [32] Levitin, E. S. and Poljak, B. T.: Constrained minimization methods. Z. Vycisl. Mat. i Mat. Fiz. 6, 787-823 (1965). English translation in USSR Comput. Math. Phys. 6, 1-50 (1965)
- [33] Lions, P. L., and Mercier., B.: Splitting algorithms for the sum of two nonlinear operators. SIAM Journal on Numerical Analysis, 16, 964-979, (1979)
- [34] Lin, Y. Y., and Pang, J.-S.: Iterative methods for large convex quadratic programs: a survey. SIAM J. Control Optim. 18, 383-411 (1987)
- [35] Luo, Z.-Q., and Tseng, P.: On the convergence of the coordinate descent method for convex differentiable minimization. J. Optim. Theory and Appl. 72, 7-35 (1992)
- [36] Luo, Z.-Q., and Tseng, P.: On the Linear convergence of descent methods for convex essentially smooth minimization. SIAM J. Control Optim. 30, 408-425 (1992)
- [37] Luo, Z.-Q. and Tseng, P.: On the convergence rate of dual ascent methods for strictly convex minimization. Mathematics of Operations Research 18, 846-867 (1993)
- [38] Ma, S.: Alternating proximal gradient method for convex minimization. Preprint (2012)
- [39] Mangasarian, O. L., and Shiau, T.-H.: Lipschitz Continuity of solutions of linear inequalities, programs and complementarity problems. SIAM J. Control Optim. 25 583-595 (1987)
- [40] Ohuchi, A., and Kaji, I.: Lagrangian dual coordinatwise maximization algorithm for network transportation problems with quadratic costs. Networks 14, 515-530 (1984)
- [41] Ortega, J. M., and Rheinboldt, W. C.: Iterative Solution of Nonlinear Equations in Several Variables. Academic Press, New York, New York. (1970)

- [42] Pang, J.-S.: On the convergence of dual ascent methods for large scale linearly constrained optimization problems. The University of Texas, School of Management, Dallas, Texas. (1984)
- [43] Pang, J.-S.: A posteriori error bounds for the linearly-constrained variational inequality problem. *Math. Oper. Res.* 12, 474–484 (1987)
- [44] Rockafellar, R. T.: *Convex Analysis*, Princeton University Press, Princeton, New Jersey. (1970)
- [45] Schneider, M. H., and Zenios, S. A.: A comparative study of algorithms for matrix balancing. *Oper. Res.* 38, 439–455 (1990)
- [46] Stephenson, D.: *Pipeline Analysis*, Elsevier, Amsterdam. (1984)
- [47] Tao, M. and Yuan, X.M.: Recovering low-rank and sparse components of matrices from incomplete and noisy observations. *SIAM J. Optim.* 21, 57-81 (2011)
- [48] Tibshirani, R.: Regression shrinkage and selection via the Lasso. *Journal of the Royal Statistical Society Ser. B.* 58, 267-288 (1996)
- [49] Tseng, P.: Dual ascent methods for problems with strictly convex costs and linear constraints: a unified approach. *SIAM J. Control Optim.* 28, 214-242 (1990)
- [50] Tseng, P.: Approximation accuracy, gradient methods, and error bound for structured convex optimization. Technical report. (2009)
- [51] Tseng, P., and Bertsekas D. P.: Relaxation methods for problems with strictly convex separable costs and linear constraints. *Math. Prog.* 38, 303–321 (1987)
- [52] Tseng, P., and Bertsekas D. P.: Relaxation methods for problems with strictly convex costs and linear constraints. *Math. Oper. Res.* 16, 462–481 (1991)
- [53] Ventura, J. A., and Hearn, D. W.: Computational development of a Lagrangian dual approach for quadratic networks. University of Florida, Industrial and Systems Engineering Department Report 88–14 (1988)
- [54] Wang, X. F., and Yuan, X. M.: The linearized alternating direction method of multipliers for dantzig selector. *SIAM Journal on Scientific Computing*, 34, 2792–2811, (2012).
- [55] Yang, J. F., and Zhang, Y.: Alternating direction algorithms for  $l_1$ -problems in compressive sensing. *SIAM Journal on Scientific Computing*, 33, 250–278 (2011).

- [56] Yuan, M. and Lin, Y.: Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. 68, 49-67 (2006)
- [57] Zhang, H., Jiang, J.J. and Luo, Z.-Q.: On the linear convergence of a proximal gradient method for a class of nonsmooth convex minimization problems. Preprint. (2012)
- [58] Zenios, S. A., and Mulvey, J. M.: Relaxation techniques for strictly convex network problems. *Ann. Oper. Res.* 5, 517–538 (1986)
- [59] Zhou, Z., Li, X., Wright, J., Candes, E.J., and Ma, Y.: Stable principal component pursuit. *Proceedings of 2010 IEEE International Symposium on Information Theory* (2010)