

Sedentary Behavior and College Students: Why It Matters Now

Samantha Zust
Project Advisor: Dr. Laura Ellingson

IOWA STATE UNIVERSITY

Department of Kinesiology, Honors Capstone, Spring 2016

Rationale

- Sedentary time (time spent sitting) has been shown to have negative effects on physical and mental health¹.
- These ill health effects are beginning to become more prevalent in younger individuals².
- Sedentary time may be a contributing factor, but little is known about the accumulation of sedentary time in this population².
- **PRIMARY AIMS:**
 - Aim 1: Characterize sedentary behavior in college-aged men and women
 - Aim 2: Examine differences in accumulation of sedentary time based on gender, age, and physical activity level

Methods

PARTICIPANTS

- 72 students (36 males, 36 females) over the age of 18

PROCEDURES

- Participants given a demographic questionnaire and ActiGraph accelerometer to be worn for one week.
- ActiGraph data processed via Sojourns software along with in-house software to create data summaries³

MEASURES

- ActiGraph GT3X+ used to objectively measure sedentary time.
 - Worn on the right hip
 - Measures acceleration as activity counts
 - Records intensity of activity in 3 axes (lying down, sitting, standing)

ANALYSES

- Calculated means and standard deviations for all data
- Effect size (Cohen's *d*) used to analyze differences in gender, age, and physical activity level

References

1. Dunstan, D. W., Howard, B., Healy, G. N., & Owen, N. (2012). Too much sitting—a health hazard. *Diabetes Research and Clinical Practice*, 97(3), 368–376. <http://doi.org/10.1016/j.diabres.2012.05.020>; Owen, N., Sparling, P. B., Healy, G. N., Dunstan, D. W., & Matthews, C. E. (2010). Sedentary behavior: emerging evidence for a new health risk. *Mayo Clinic Proceedings*, 85(12), 1138–1141. <http://doi.org/10.4065/mcp.2010.0444>; Teychenne, M., Ball, K., & Salmon, J. (2010). Sedentary behavior and depression among adults: a review. *International Journal of Behavioral Medicine*, 17(4), 246–254. <http://doi.org/10.1007/s12529-010-9075-z>; Teychenne, M., Costigan, S. A., & Parker, K. (2015). The association between sedentary behaviour and risk of anxiety: a systematic review. *BMC Public Health*, 15, 513. <http://doi.org/10.1186/s12889-015-1843-x>
2. McGavock, J. M., Anderson, T. J., & Lewanczuk, R. Z. (2006). Sedentary Lifestyle and Antecedents of Cardiovascular Disease in Young Adults. *American Journal of Hypertension*, 19(7), 701–707. <http://doi.org/10.1016/j.amjhyper.2006.01.013>
3. Lyden, K., Keadle, S. K., Staudenmayer, J., & Freedson, P. S. (2014). A method to estimate free-living active and sedentary behavior from an accelerometer. *Medicine and Science in Sports and Exercise*, 46(2), 386–397. <http://doi.org/10.1249/MSS.0b013e3182a42a2d>



Key Findings

- On average, subjects spent 72% of their day (639.76 + 80.09 minutes/day) sedentary.
 - 52% in bouts of 30+ minutes or more
 - 25% in bouts of 60+ minutes or more
- Small difference between weekday and weekend sedentary time
- Most sedentary between 12:00pm and 5:00pm

Conclusions

- Young adults lead a primarily sedentary lifestyle, possibly due to the structure of university life.
- Future interventions should focus on reducing sedentary behavior in addition to promoting MVPA in college students. Potential strategies include technology-based interventions, more breaks in class & while studying, and increasing campus walkability.

Results 1: How sedentary are college students?



Results 2: What is the relationship between sedentary time, MVPA, and light PA?

