

1-2002

Manure Resources

Jeffery Lorimor
Iowa State University

Antonio Mallarino
Iowa State University, apmallar@iastate.edu

John Sawyer
Iowa State University, jsawyer@iastate.edu

John Creswell
Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/extension_pubs

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Bioresource and Agricultural Engineering Commons](#)

Recommended Citation

Lorimor, Jeffery; Mallarino, Antonio; Sawyer, John; and Creswell, John, "Manure Resources" (2002). *Extension and Outreach Publications*. 102.
http://lib.dr.iastate.edu/extension_pubs/102

Iowa State University Extension and Outreach publications in the Iowa State University Digital Repository are made available for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current publications and information from Iowa State University Extension and Outreach, please visit <http://www.extension.iastate.edu>.

Manure Resources

Using manure as a resource.

Use manure as a nutrient source.

Each year, Iowa livestock producers recover enough crop-available nitrogen (N), phosphorus (P) and potassium (K) in manure to supply approximately 13 pounds of N, 13 pounds of P and 21 pounds of K for every acre of corn grown in Iowa.

Most producers understand that manure is a resource rather than a waste product. Since efficiently recycling manure nutrients can reduce reliance on fertilizers, applying manure using scientifically sound methods is like putting money in your pocket.

To realize the full value of manure requires your taking a few moments and working through some simple steps.

- Test the nutrient concentration of manure.
- Soil test and apply manure where it is needed, based on crop N, P, or K needs.
- Calibrate the manure applicator for a known and even distribution of manure.

Analyze manure before applying and build an analysis database.

Have a commercial laboratory determine the concentration of nutrients in the manure. Manure samples should be taken before application to help guide application rates. Multiple samples also should be collected during application to build a manure nutrient database. These analyses help manage the manure with the same care as the care you use with fertilizer.

Key Points

- Use manure as a nutrient source.
- Analyze manure before applying and build an analysis database.
- Apply manure where it is needed, according to nitrogen (N), phosphorus (P), or potassium (K) crop needs, soil test data and your manure management plan.
- Calibrate manure applicators.
- Contact your Iowa State University county Extension office for more information.

Manure consists of both solids and liquid, so one sample may not represent the overall nutrient value. For example: pits become stratified, with solids concentrated at the bottom. To get a sample of the average nutrient content, probe the full depth of the pit several times and combine into a composite sample. Or, agitate the manure vigorously for several hours before sampling. Sample while the agitator is running or

Table 1. Interpretation of P and K soil test values for surface soil samples (6 to 7-inch depth).

Relative level*	Bray P _i or Mehlich-3 P			Olsen P			Ammonium Acetate or Mehlich-3 Extractable K	
	Wheat, Alfalfa	All crops except wheat, alfalfa		Wheat, Alfalfa	All crops except wheat, alfalfa		All Crops	
	Subsoil P			Subsoil P			Subsoil K	
	Low	High		Low	High		Low	High
	parts per million (ppm)							
Very Low (VL)	0-15	0-8	0-5	0-10	0-5	0-3	0-60	0-40
Low (L)	16-20	9-15	6-10	11-14	6-10	4-7	61-90	41-80
Optimum (Opt)	21-25	16-20	11-15	15-17	11-14	8-11	91-130	81-120
High (H)	26-30	21-30	16-20	18-20	15-20	12-15	131-170	121-160
Very High (VH)	31+	31+	21+	21+	21+	16+	171+	161+

From PM 1688, General Guide for Crop Nutrient Recommendations in Iowa.

when the spreader is being loaded.

If you feed on an open lot, scrape across the lot with a shovel to get a composite sample.

Getting a uniform sample is difficult, but doing the best job you can means using a consistent and meticulous approach to sampling.

If management and feed rations do not change, manure analysis should remain fairly stable. But samples should still be taken every year. Maintain these manure analysis records, and build an analytical nutrient management database. Document the date, location, soil condition, and source and amount of manure applied to fields.

Apply manure where it is needed, according to N, P, or K needs, soil test data and your manure management plan.

Use a current soil test analysis from a certified laboratory. Have

a certified crop advisor or independent crop consultant assist you in developing a nutrient management plan for each field. Table 1 will help you understand which fields or field areas to target for manure application.

The optimum soil test category is the most profitable to maintain. The very high soil test category indicates that the nutrient concentration exceeds crop needs, and further additions of that nutrient very seldom produce a profitable yield response.

Applying manure in fields where fertility levels are very high wastes a valuable resource, and repeated applications will result in a rapid increase in soil test P and potassium (K).

Consider the needs of the crops in rotation, and allocate manure to fields based on the N, P, or K requirements of the crop to be planted. A suggested manure allocation strategy is:

- First, fields testing very low, low, or optimum in P and K that will be planted to corn.
- Second, fields testing very low, low, or optimum in P and K that will be planted to soybeans.
- Third, fields testing high to very high in P and K that will be planted to corn.

Note: To slow nitrification, do not apply manure in the fall until the soil temperature is 50 degrees F at a depth of four inches and cooling. Injection of manure is preferred over broadcast application.

Calibrate manure applicators.

Since livestock manure is a resource and is applied to cropland for its nutrient value, getting an even spread of manure is important. Correct manure applicator calibration, operating speed and even application

throughout the field is as important as properly sampling the manure.

From the laboratory analysis, calculate the nutrients to be applied in pounds or gallons per acre.

Field-calibrate your applicator. Assistance may be obtained from a certified crop advisor, independent crop consultant, or an ISU Extension crop or agricultural engineering specialist.

Contact your ISU county Extension office for more information.

Publications available to help you manage manure include:

PM 1558, *Management Practices: How to Sample Manure for Nutrient Analysis*

PM 1811, *Managing Manure Nutrients for Crop Production*

Best Management Practices, or BMPs, utilize the most effective and practical means available to reduce or prevent water pollution from farm operations. BMPs are selected based on assessment, analysis of the impact of alternative practices and their economic considerations. They are implemented using current available technologies, management skills and available resources. BMP information sheets available from ISU Extension include:

NMEP 1, *Soil Testing*

NMEP 2, *Phosphorus Application*

NMEP 3, *Manure Resources*

NMEP 4, *Residue Management*

NMEP 5, *Crop Rotation*

NMEP 6, *Crop Yields*

NMEP 7, *Nitrogen Application*

NMEP 8, *Nutrient Management Plan*

NMEP 9, *Equipment Calibration*

NMEP 10, *Conservation Reserve Program*

NMEP 11, *Conservation Practices*

Prepared by Jeffery Lorimor, associate professor, department of agricultural and biosystems engineering, Iowa State University; Antonio Mallarino and John Sawyer, associate professors, department of agronomy, Iowa State University; and John Creswell, coordinator of the nutrient management education project.

This publication has been funded in part by the Iowa Department of Natural Resources through a grant from the U.S. Environmental Protection Agency under the Federal Nonpoint Source Management Program (Section 319 of the Clean Water Act).

IOWA STATE UNIVERSITY

University Extension

Helping Iowans become their best.



... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Many materials can be made available in alternative formats for ADA clients. To file a complaint of discrimination, write USDA, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Stanley R. Johnson, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.