

2-11-2016

Asymptotic density and the coarse computability bound

Dennis R. Hirschfeldt

University of Chicago

Carl G. Jockusch

University of Illinois at Urbana-Champaign

Timothy H. McNicholl

Iowa State University, mcnichol@iastate.edu

Paul E. Schupp

University of Illinois at Urbana-Champaign

Follow this and additional works at: http://lib.dr.iastate.edu/math_pubs



Part of the [Mathematics Commons](#), and the [Numerical Analysis and Computation Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/math_pubs/110. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Mathematics at Iowa State University Digital Repository. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

ASYMPTOTIC DENSITY AND THE COARSE COMPUTABILITY BOUND

DENIS R. HIRSCHFELDT, CARL G. JOCKUSCH, JR., TIMOTHY H. MCNICHOLL,
AND PAUL E. SCHUPP

ABSTRACT. For $r \in [0, 1]$ we say that a set $A \subseteq \omega$ is *coarsely computable at density r* if there is a computable set C such that $\{n : C(n) = A(n)\}$ has lower density at least r . Let $\gamma(A) = \sup\{r : A \text{ is coarsely computable at density } r\}$. We study the interactions of these concepts with Turing reducibility. For example, we show that if $r \in (0, 1]$ there are sets A_0, A_1 such that $\gamma(A_0) = \gamma(A_1) = r$ where A_0 is coarsely computable at density r while A_1 is not coarsely computable at density r . We show that a real $r \in [0, 1]$ is equal to $\gamma(A)$ for some c.e. set A if and only if r is left- Σ_3^0 . A surprising result is that if G is a Δ_2^0 1-generic set, and $A \leq_T G$ with $\gamma(A) = 1$, then A is coarsely computable at density 1.

1. INTRODUCTION

There are two natural models of “imperfect computability” defined in terms of the standard notion of asymptotic density, which we now review. For $A \subseteq \omega$ and $n \in \omega \setminus \{0\}$, define $\rho_n(A)$, the density of A below n , by $\rho_n(A) = \frac{|A \upharpoonright n|}{n}$, where $A \upharpoonright n = A \cap \{0, 1, \dots, n-1\}$. Then

$$\underline{\rho}(A) = \liminf_n \rho_n(A) \quad \text{and} \quad \bar{\rho}(A) = \limsup_n \rho_n(A)$$

are respectively the *lower density* of A and the *upper density* of A . The (*asymptotic density*) of A is $\rho(A) = \lim_n \rho_n(A)$ provided the limit exists.

The idea of generic computability was introduced and studied in connection with group theory in [10] and then studied in connection with arbitrary subsets of ω in [9]. In generic computability we have a partial algorithm that is always correct when it gives an answer but may fail to answer on a set of density 0. The paper [5] began studying computability at densities less than 1 and introduced the following definitions.

Definition 1.1 ([5, Definition 5.9]). Let A be a set of natural numbers and let r be a real number in the unit interval $[0, 1]$. The set A is *partially computable at density r* if there is a partial computable function φ such that $\varphi(n) = A(n)$ for all n in the domain of φ and the domain of φ has lower density at least r .

Thus A is *generically computable* if and only if A is partially computable at density 1.

1991 *Mathematics Subject Classification*. Primary 03D28; Secondary 03D25.

Key words and phrases. Asymptotic density, Coarse computability, Turing degrees.

Hirschfeldt was partially supported by grant DMS-1101458 from the National Science Foundation of the United States.

McNicholl was partially supported by a Simons Foundation Collaboration Grant for Mathematicians.

Definition 1.2 ([5, Definition 6.9]). If $A \subseteq \omega$, the *partial computability bound* of A is

$$\alpha(A) = \sup\{r : A \text{ is partially computable at density } r\}.$$

In the paper [5] the term “partially computable at density r ” was simply called “computable at density r ” and the “partial computability bound” was called the “asymptotic computability bound”. That paper considered only partial computability at densities less than 1, but since we are here comparing the partial computability concepts with their coarse analogs, the present terminology is more exact.

If A is generically computable, then $\alpha(A) = 1$. The converse fails by [5, Observation 5.10]. There are sets that are partially computable at every density less than 1 but are not generically computable.

Definition 1.3. If $A, B \subseteq \mathbb{N}$, then A and B are *coarsely similar*, written $A \sim_c B$, if the density of the symmetric difference of A and B is 0, that is, $\rho(A \Delta B) = 0$. Given A , any set B such that $B \sim_c A$ is called a *coarse description* of A .

It is easy to check that coarse similarity is indeed an equivalence relation. Coarse similarity was called *generic similarity* in [9], but the current terminology seems better.

Coarse computability considers algorithms that always give an answer, but may give an incorrect answer on a set of density 0. We have the following definition.

Definition 1.4 ([9, Definition 2.13]). The set A is *coarsely computable* if there is a computable set C such that the density of $\{n : A(n) = C(n)\}$ is 1. That is, A is coarsely computable if it has a computable coarse description C .

The following definitions are similar to those for partial computability.

Definition 1.5. If $A \subseteq \omega$ and $r \in [0, 1]$, an *r -description* of A is any set B such that the lower density of $\{n : A(n) = B(n)\}$ is at least r . A set A is *coarsely computable at density r* if there is a computable r -description B of A .

Note that A is coarsely computable if and only if A is coarsely computable at density 1.

Definition 1.6. If $A \subseteq \omega$, the *coarse computability bound* of A is

$$\gamma(A) = \sup\{r : A \text{ is coarsely computable at density } r\}.$$

If A is coarsely computable, then $\gamma(A) = 1$, but the next lemma implies that the converse fails.

It is shown in [9, Proposition 2.15 and Theorem 2.26] that neither of generic computability and coarse computability implies the other, even for c.e. sets. Nonetheless, the following lemma gives an inequality between α and γ .

Lemma 1.7. *For any $A \subseteq \omega$, $\alpha(A) \leq \gamma(A)$. In particular, if A is generically computable then $\gamma(A) = 1$.*

Proof. Fix $\epsilon > 0$. If $\alpha(A) = r$ then there is a partial algorithm φ for A such that the lower density of the c.e. set $D = \text{dom } \varphi$ is greater than or equal to $r - \epsilon$. Theorem 3.9 of [5] shows that if D is a c.e. set there is a computable set $C \subseteq D$ such that $\underline{\rho}(C) > \underline{\rho}(D) - \epsilon$. Let $C_1 = \{n \in C : \varphi(n) = 1\}$. Then C_1 is a computable set and $\{n : A(n) = C_1(n)\} \supseteq C$. It follows that $\underline{\rho}(\{n : A(n) = C_1(n)\}) \geq \underline{\rho}(C) > \underline{\rho}(D) - \epsilon \geq r - 2\epsilon$, and hence A is coarsely computable at density $r - 2\epsilon$. Since $\epsilon > 0$ was arbitrary, it follows that $\gamma(A) \geq r = \alpha(A)$. \square

One consequence of this lemma is that any set that is generically computable but not coarsely computable is an example of a set A such that $\gamma(A) = 1$ but A is not coarsely computable.

Definition 1.8. If $A, B \subseteq \mathbb{N}$, let $D(A, B) = \bar{\rho}(A \Delta B)$.

It is shown in [5, remarks after Proposition 3.2] that D is a pseudometric on subsets of ω and, since $D(A, B) = 0$ exactly when A and B are coarsely similar, D is actually a metric on the space of coarse similarity classes. Note that γ is an invariant of coarse similarity classes.

Although easy, the following is useful enough to be stated as a lemma.

Lemma 1.9. *If $A \subseteq \omega$ then $\rho(A) = 1 - \bar{\rho}(\bar{A})$.*

Proof. Note that $\rho_n(A) = 1 - \rho_n(\bar{A})$ for all $n \geq 1$. The lemma follows by taking the lim inf of both sides of this equation. \square

Since we have a pseudometric space, we can consider the distance from a single point to a subset of the space in the usual way.

Definition 1.10. If $A \subseteq \omega$ and $\mathcal{S} \subseteq \mathcal{P}(\mathbb{N})$, let

$$\delta(A, \mathcal{S}) = \inf\{D(A, S) : S \in \mathcal{S}\}.$$

The above lemma shows that

$$\gamma(A) = 1 - \delta(A, \mathcal{C}),$$

where \mathcal{C} is the class of computable sets. Thus $\gamma(A) = 1$ if and only if A is a limit of computable sets in the pseudometric. A set A is coarsely computable at density r if and only if $\delta(A, \mathcal{C}) \leq 1 - r$.

The symmetric difference $A \Delta B = \{n : A(n) \neq B(n)\}$ is the subset of ω where A and B disagree. There does not seem to be a standard notation for the complement of $A \Delta B$, which is $\{n : A(n) = B(n)\}$, the ‘‘symmetric agreement’’ of A and B . We find it useful to use $A \nabla B$ to denote $\{n : A(n) = B(n)\}$.

We assume that the reader is familiar with basic computability theory. See, for example, [13]. If S is a set of finite binary strings and $A \subseteq \omega$ we say that A *meets* S if A extends some string in S and that A *avoids* S if A extends a string that has no extension in S .

2. TURING DEGREES, COARSE COMPUTABILITY, AND γ

It is easily seen that every Turing degree contains a set that is both coarsely and generically computable and hence a set A with $\alpha(A) = \gamma(A) = 1$. In the other direction it is shown in Theorem 2.20 of [9] that every nonzero Turing degree contains a set that is neither generically computable nor coarsely computable. The same construction now yields a quantitative version of that result.

Theorem 2.1. *Every nonzero Turing degree contains a set whose partial computability bound is 0 but whose coarse computability bound is $1/2$.*

Proof. Let $I_n = [n!, (n+1)!)$. Suppose that A is not computable, and let $\mathcal{I}(A) = \bigcup_{n \in A} I_n$. It is clear that $\mathcal{I}(A)$ is Turing equivalent to A . We prove first that $\gamma(\mathcal{I}(A)) \leq \frac{1}{2}$. If there is a computable C with $\rho(\mathcal{I}(A) \nabla C) > \frac{1}{2}$ we can compute A by ‘‘majority vote’’. That is, for all sufficiently large n , we have that n is in A if and only if more than half of the elements of I_n are in C . (For any n for which

this equivalence fails, we have $\rho_{(n+1)!}(\mathcal{I}(A) \nabla C) \leq (1 + (n+1)^{-1})/2$.) It follows that A is computable, a contradiction. If C is the set of even numbers, then it is easily seen that $\rho(C \nabla \mathcal{I}(A)) = \frac{1}{2}$, so $\gamma(\mathcal{I}(A)) \geq \frac{1}{2}$. It follows that $\gamma(\mathcal{I}(A)) = \frac{1}{2}$. To see that $\alpha(\mathcal{I}(A)) = 0$, note that any set of positive lower density intersects I_n for all but finitely many n , and apply this observation to the domain of any partial computable function that agrees with $\mathcal{I}(A)$ on its domain. \square

We next observe that a large class of degrees contain sets A with $\gamma(A) = 0$.

Theorem 2.2. *Every hyperimmune degree contains a set whose coarse computability bound is 0.*

Proof. A set $S \subseteq 2^{<\omega}$ of finite binary strings is *dense* if every string has some extension in S . Stuart Kurtz [11] defined a set A to be *weakly 1-generic* if A meets every dense c.e. set S of finite binary strings and proved that the weakly 1-generic degrees coincide with the hyperimmune degrees. Hence, it suffices to show that every weakly 1-generic set A satisfies $\gamma(A) = 0$. Assume that A is weakly 1-generic.

If f is a computable function then, for each $n, j > 0$, define

$$S_{n,j} = \left\{ \sigma \in 2^{<\omega} : |\sigma| \geq j \ \& \ \rho_{|\sigma|}(\{k < |\sigma| : \sigma(k) = f(k)\}) < \frac{1}{n} \right\}.$$

Each set $S_{n,j}$ is computable and dense so A meets each $S_{n,j}$. Thus $\{k : f(k) = A(k)\}$ has lower density 0. \square

In view of the preceding result, it is natural to ask whether *every* nonzero degree contains a set A such that $\gamma(A) = 0$. This question is answered in the negative in [1] where it is shown that every computably traceable set is coarsely computable at density $\frac{1}{2}$, and also that every set computable from a 1-random set of hyperimmune-free degree is coarsely computable at density $\frac{1}{2}$. Each of these results implies that there is a nonzero degree $\mathbf{a} \leq \mathbf{0}''$ such that every \mathbf{a} -computable set is coarsely computable at density $\frac{1}{2}$. Here it is not possible to replace $\frac{1}{2}$ by any larger number, by Theorem 2.1. In [1], the following definition is made for Turing degrees \mathbf{a} :

$$\Gamma(\mathbf{a}) = \inf\{\gamma(A) : A \text{ is } \mathbf{a}\text{-computable}\}.$$

By the above, Γ takes on the values 0 and $\frac{1}{2}$, and of course $\Gamma(\mathbf{0}) = 1$. By Theorem 2.1, Γ does not take on any values in the open interval $(\frac{1}{2}, 1)$. An open question posed in [1] is whether Γ takes on any values other than $0, \frac{1}{2}$, and 1.

3. COARSE COMPUTABILITY AT DENSITY $\gamma(A)$

If A is any set, it follows from the definition of $\gamma(A)$ that A is coarsely computable at every density less than $\gamma(A)$ and at no density greater than $\gamma(A)$. What happens at $\gamma(A)$? Let us say that A is *extremal for coarse computability* if it is coarsely computable at density $\gamma(A)$. In this section, we show that extremal and non-extremal sets exist. Moreover, we also show that every real in $(0, 1]$ is the coarse computability bound of an extremal set and of a non-extremal set. We also explore the distribution of these cases in the Turing degrees. Roughly speaking, we show that hyperimmune degrees yield extremal sets and high degrees yield non-extremal sets.

Theorem 3.1. *Every real in $[0, 1]$ is the coarse computability bound of a set that is extremal for coarse computability.*

Proof. Suppose $0 \leq r \leq 1$. By Corollary 2.9 of [9] there is a set A_1 such that $\rho(A_1) = r$. Let Z be a set with $\gamma(Z) = 0$, which exists by Theorem 2.2, and let $A = A_1 \cup Z$. Note first that that A is coarsely computable at density r via the computable set ω since

$$\underline{\rho}(A \nabla \omega) = \underline{\rho}(A) \geq \underline{\rho}(A_1) = r.$$

It follows that $\gamma(A) \geq r$, so it remains only to show that $\gamma(A) \leq r$.

Suppose for a contradiction that $\gamma(A) > r$, so A is coarsely computable at some density $r' > r$. Let C be a computable set such that $\underline{\rho}(A \nabla C) \geq r'$. Let:

$$\begin{aligned} S_1 &= A_1 \cap C \\ S_2 &= (Z \setminus A_1) \cap C \\ S_3 &= \bar{A} \cap \bar{C}. \end{aligned}$$

Note that $A \nabla C$ is the disjoint union of S_1 , S_2 , and S_3 so

$$\rho_n(A \nabla C) = \rho_n(S_1) + \rho_n(S_2) + \rho_n(S_3)$$

for all n .

Let $\epsilon = r' - r$. For all sufficiently large n we have $\rho_n(A \nabla C) > r + \frac{\epsilon}{2}$. Since $S_1 \subseteq A_1$ and $\rho_n(A_1) < r + \frac{\epsilon}{3}$ for all sufficiently large n , we have $\rho_n(S_2) + \rho_n(S_3) > \frac{\epsilon}{6}$ for all sufficiently large n . Hence $\underline{\rho}(S_2 \cup S_3) > 0$. But $S_2 \cup S_3 \subseteq C \nabla Z$ so $\underline{\rho}(C \nabla Z) > 0$, contradicting $\gamma(Z) = 0$. This contradiction shows that $\gamma(A) \leq r$, and the proof is complete. \square

Corollary 3.2 (to proof). *Suppose \mathbf{a} is a hyperimmune degree. Then, every Δ_2^0 real in $[0, 1]$ is the coarse computability bound of a set in \mathbf{a} that is extremal for coarse computability.*

Proof. Just note that the proof of the theorem can be carried out effectively in \mathbf{a} . In more detail, by Theorem 2.21 of [9] there is a computable set A_1 of density r . Further, by Theorem 2.2 there is an \mathbf{a} -computable set Z such that $\gamma(Z) = 0$. Then $A = A_1 \cup Z$ satisfies the theorem and is \mathbf{a} -computable. We can ensure that $A \in \mathbf{a}$ by coding a set in \mathbf{a} into A on a set of density 0. \square

We now consider sets that are not extremal for coarse computability. We first consider the degrees of the sets A such that $\gamma(A) = 1$ but A is not coarsely computable.

Define

$$R_n = \{k : 2^n \mid k \ \& \ 2^{n+1} \nmid k\}.$$

The sets R_n were heavily used in [9] and [5]. Note that they are uniformly computable and pairwise disjoint, and $\rho(R_n) = 2^{-(n+1)}$. As in [9] and [5], define

$$\mathcal{R}(A) = \bigcup_{n \in A} R_n.$$

Note that, for all A , we have that $A \equiv_{\mathcal{T}} \mathcal{R}(A)$ and $\alpha(\mathcal{R}(A)) = \gamma(\mathcal{R}(A)) = 1$. To see the latter (which was pointed out by Asher Kach), note that if $C_k = \bigcup\{R_n : n \in A \ \& \ n < k\}$, then C_k is computable and agrees with $\mathcal{R}(A)$ on $\bigcup_{n < k} R_n$, and the latter has density $1 - 2^{-k}$.

Theorem 3.3. (i) *If \mathbf{a} is a degree such that $\mathbf{a} \not\leq \mathbf{0}'$, then \mathbf{a} contains a set that is not coarsely computable but whose coarse computability bound is 1.*

- (ii) If \mathbf{a} is a nonzero c.e. degree, then \mathbf{a} contains a c.e. set that is not coarsely computable but whose coarse computability bound is 1.

Proof. It is shown in Theorem 2.19 of [9] that $\mathcal{R}(B)$ is coarsely computable if and only if B is Δ_2^0 . If $\mathbf{a} \not\leq \mathbf{0}'$ and B has degree \mathbf{a} , then $\mathcal{R}(B)$ is a set of degree \mathbf{a} that is not coarsely computable even though its coarse computability bound is 1. Part (i) follows.

Theorem 4.5 of [5] shows that every nonzero c.e. degree contains a c.e. set A that is generically computable but not coarsely computable. Then $\alpha(A) = 1$, so by Lemma 1.7, $\gamma(A) = 1$. This proves part (ii). \square

This result raises the natural question: Does *every* nonzero Turing degree contain a set A such that $\gamma(A) = 1$ but A is not coarsely computable? We will obtain a negative answer in Theorem 5.11 in the next section. In fact, we will show that if G is 1-generic and Δ_2^0 , and $A \leq_T G$ has $\gamma(A) = 1$, then A is coarsely computable.

We now consider the coarse computability bounds of non-extremal sets.

Theorem 3.4. *Every real in $(0, 1]$ is the coarse computability bound of a set that is not extremal for coarse computability.*

Proof. Suppose $0 < r \leq 1$. We construct a set A so that $\gamma(A) = r$ but A is not coarsely computable at density r . As an auxiliary for defining A , we first use the technique of Corollary 2.9 of [9] to define a set S of density r . To this end, we turn r into a set B in the natural way. That is, since $r > 0$, it has a non-terminating binary expansion $r = 0.b_0b_1\dots$. We then set $B = \{i : b_i = 1\}$. By restricted countable additivity (Lemma 2.6 of [9]), $\mathcal{R}(B)$ has density r . Set $S = \mathcal{R}(B)$.

We now divide S into “slices” S_0, S_1, \dots as follows. Let $c_0 < c_1 < \dots$ be the increasing enumeration of B . Set $S_e = R_{c_e}$. Note that the S_e ’s are pairwise disjoint and that $S = \bigcup_e S_e$. Note also that each S_e is computable (though not necessarily computable uniformly in e).

We now define A . We first choose a set Z so that $\gamma(Z) = 0$. Such a set exists by Theorem 2.2. Let C_0, C_1, \dots be an enumeration of the computable sets. We then set

$$A = (\overline{S} \cap Z) \cup \bigcup_e (S_e \cap \overline{C_e}).$$

We now claim that A is coarsely computable at density q whenever $0 \leq q < r$. For, suppose $0 \leq q < r$. Since the density of S is r , there is a number n so that $\rho(\bigcup_{e < n} S_e) \geq q$. Let $C = \bigcup_{e < n} (S_e \cap \overline{C_e})$. Then, C is a computable set. Also A and C agree on each S_e for $e < n$, so $\rho(A \nabla C) \geq \rho(\bigcup_{e < n} S_e) \geq q$. Hence, C witnesses that A is coarsely computable at density q .

To complete the proof, it suffices to show that A is not coarsely computable at density r . To this end, it suffices to show that the lower density of $A \nabla C_e$ is smaller than r for each e . Fix $e \in \mathbb{N}$. By construction, $(C_e \nabla A) \cap S$ is disjoint from S_e and so has *upper* density less than r . At the same time, note that $(A \nabla C_e) \cap \overline{S} \subseteq C_e \nabla Z$. Let $r_0 = \overline{\rho}((C_e \nabla A) \cap S)$, and let $\epsilon = r - r_0$. Then for infinitely many n we have

$$\rho_n(A \nabla C_e) = \rho_n((A \nabla C_e) \cap S) + \rho_n((A \nabla C_e) \cap \overline{S}) < \left(r_0 + \frac{\epsilon}{2}\right) + \frac{\epsilon}{3} < r.$$

It follows that $\rho(A \nabla C_e) < r$. Hence A is not coarsely computable at density r , which completes the proof. \square

We note that the proof of Theorem 3.4 shows that if A is any set so that $A \cap S_e = \overline{C_e} \cap S_e$ for all e , then A is computable at density q whenever $0 \leq q < r$. That is, the construction of $A \cap S$ ensures that $\gamma(A) \geq r$.

Corollary 3.5 (to proof). *Suppose \mathbf{a} is a high degree. Then, every computable real in $(0, 1]$ is the coarse computability bound of a set in \mathbf{a} that is not extremal for coarse computability.*

Proof. We just observe that the preceding proof can be carried out in an \mathbf{a} -computable fashion. By Theorem 1 of [8], there is a listing C_0, C_1, \dots of the computable sets that is uniformly \mathbf{a} -computable. Also, since r is computable, the sequence S_0, S_1, \dots in the proof of the theorem is also uniformly \mathbf{a} -computable. It does not affect the proof to modify each S_e so that it contains no numbers less than e , and then $S = \bigcup_e S_e$ is \mathbf{a} -computable. Finally, every high degree is hyperimmune by a result of D. A. Martin [12], and so every high degree computes a set Z with $\gamma(Z) = 0$ by Theorem 2.2. Hence the set A defined in the proof of the theorem can be chosen to be \mathbf{a} -computable. By coding a set in \mathbf{a} into A on a set of density 0 we can ensure that $A \in \mathbf{a}$. \square

By using suitable computable approximations, the previous corollary can be extended from computable reals to Δ_2^0 reals. We omit the details.

It was shown in Theorem 4.5 of [5] that every nonzero c.e. degree contains a c.e. set that is generically computable but not coarsely computable. It follows at once from Lemma 1.7 that every nonzero c.e. degree contains a c.e. set A such that $\gamma(A) = 1$ but A is not coarsely computable. We now use the method of Theorem 3.4 to extend this result to the case where $\gamma(A)$ is a given computable real.

Theorem 3.6. *Suppose \mathbf{a} is a nonzero c.e. degree. Then, every computable real in $(0, 1]$ is the coarse computability bound and the partial computability bound of a c.e. set in \mathbf{a} that is not extremal for coarse computability.*

Proof. Define the sets S, S_0, S_1, \dots as in the proof of Theorem 3.4 so that $S = \bigcup_e S_e$ and so that $\rho(S) = r$. Let B be a c.e. set of degree \mathbf{a} , and let $\{B_s\}$ be a computable enumeration of B . We construct the desired set $A \leq_T B$ using ordinary permitting; i.e. if $x \in A_{s+1} \setminus A_s$, then there exists $y \leq x$ such that $y \in B_{s+1} \setminus B_s$. To ensure that $B \leq_T A$, we code B into A on a set of density zero.

Let the requirement N_e assert that if Φ_e is total, then the lower density of the set on which it agrees with A is smaller than r . Thus, if N_e is met for every e , then A is not coarsely computable at density r . We meet N_e by appropriately defining A on S_e and on \overline{S} . If Φ_e is total, we meet N_e by making A completely disagree with Φ_e on infinitely many large finite sets $I \subseteq S_e \cup \overline{S}$. To this end, we effectively choose finite sets $I_{e,i}$ such that the following hold for all e, i, e' , and i' :

- (i) $I_{e,i} \subseteq (S_e \cup \overline{S})$.
- (ii) $\min I_{e,i+1} > \max I_{e,i}$.
- (iii) $\rho_m(I_{e,i}) \geq \frac{i}{i+1} \rho_m(S_e \cup \overline{S})$ where $m = \max I_{e,i} + 1$.
- (iv) If $(e, i) \neq (e', i')$, then $I_{e,i} \cap I_{e',i'} = \emptyset$.

The sets $I_{e,i}$ may be obtained by intersecting appropriately large intervals with $S_e \cup \overline{S}$ while preserving pairwise disjointness, and we will call the sets $I_{e,i}$ “intervals”. During the construction we will designate an interval $I_{e,i}$ as “successful” if we have ensured that Φ_e and A totally disagree on $I_{e,i}$. The construction is as follows:

Stage 0. Let $A_0 = \emptyset$.

Stage $s + 1$. For each $e, i \leq s$, declare $I_{e,i}$ to be successful if it has not yet been declared successful and if the following conditions are met.

- (1) $\Phi_{e,s}$ is defined on all elements of $I_{e,i}$.
- (2) $\min(I_{e,i})$ exceeds all elements of $A_s \cap S_e$.
- (3) At least one number in $B_{s+1} \setminus B_s$ is less than or equal to $\min(I_{e,i})$.

If $I_{e,i}$ is declared to be successful at stage $s + 1$, then enumerate into A all $x \in I_{e,i}$ with $\Phi_e(x) = 0$.

The set A is clearly c.e., and $A \leq_T B$ by ordinary permitting. If the interval $I_{e,i}$ is ever declared to be successful, then A and Φ_e totally disagree on $I_{e,i}$, by the action taken when it is declared successful and the disjointness condition (iv), which ensures that no elements of $I_{e,i}$ are enumerated into A except by this action.

Note that (2) ensures that $A \cap S_e$ is computable for each e . It follows that $\gamma(A) \geq \alpha(A) \geq r$ as in the proof of Theorem 3.4.

It remains to show that every requirement N_e is met. Suppose that Φ_e is total. We claim first that the interval $I_{e,i}$ is declared successful for infinitely many i . Suppose not. Then $A \cap S_e$ is finite. It follows that B is computable, since, for all sufficiently large i , if $s \geq i$ and $\Phi_{e,s}$ is defined on all elements of $I_{e,i}$, then no number less than $\min(I_{e,i})$ enters B after stage s . Since we assumed that B is noncomputable, the claim follows.

Suppose $I_{e,i}$ is successful. Set $I = I_{e,i}$. Then $A \Delta \Phi_e \supseteq I$, so

$$\rho_m(A \Delta \Phi_e) \geq \rho_m(I) \geq \frac{i}{i+1} \rho_m(S_e \cup \bar{S}),$$

where $m = \max I_{e,i} + 1$. There are infinitely many such i 's, and as i tends to infinity, the right hand side of the above inequality tends to $\rho(S_e) + \rho(\bar{S})$. It follows that $\bar{\rho}(A \Delta \Phi_e) \geq \rho(S_e) + (1 - r)$, and so by Lemma 1.9, $\underline{\rho}(A \nabla \Phi_e) \leq r - \rho(S_e) < r$, as needed to complete the proof. \square

4. COARSE COMPUTABILITY AND LOWNESS

We now consider the coarse computability properties of c.e. sets that have a density.

Proposition 4.1. *Every low c.e. set having a density is coarsely computable. Every c.e. set having a density has coarse computability bound 1.*

Proof. The first statement is Corollary 3.16 of [5]. Let A be a c.e. set that has a density and let $\epsilon > 0$. Theorem 3.9 of [5] shows that A has a computable subset C such that $\underline{\rho}(C) > \rho(A) - \epsilon$. Then $C \Delta A = A \setminus C$. Hence, by Lemma 1.9, $\underline{\rho}(A \nabla C) = 1 - \bar{\rho}(A \setminus C)$. But by Lemma 3.3 (iii) of [5],

$$\bar{\rho}(A \setminus C) \leq \rho(A) - \underline{\rho}(C) < \epsilon.$$

Hence $\underline{\rho}(A \nabla C) > 1 - \epsilon$. Since $\epsilon > 0$ was arbitrary, we conclude that $\gamma(A) = 1$. \square

The next result shows that the lowness assumption is strongly required in the first part of Proposition 4.1.

Theorem 4.2. *Every nonlow c.e. Turing degree \mathbf{a} contains a c.e. set of density $1/2$ that is not coarsely computable.*

Proof. The proof of the theorem is similar to the proof in Theorem 4.3 of [5] that every nonlow c.e. degree contains a c.e. set A such that $\rho(A) = 1$ but A has no computable subset of density 1. Hence we give only a sketch. Let C be a c.e. set of degree \mathbf{a} . We ensure that $A \leq_T C$ by a slight variation of ordinary permitting: If x enters A at stage s , then either some number $y \leq x$ enters C at s or $x = s$. This implies that $A \leq_T C$, and by coding C into A on a set of density 0 we can ensure that $A \equiv_T C$ without disturbing the other desired properties of A .

To ensure that $\rho(A) = \frac{1}{2}$, we arrange that $\rho(A \cap R_n) = \frac{\rho(R_n)}{2}$ for all n . Then by restricted countable additivity (Lemma 2.6 of [9]),

$$\rho(A) = \sum_n \rho(A \cap R_n) = \sum_n \frac{\rho(R_n)}{2} = \frac{\sum_n \rho(R_n)}{2} = \frac{1}{2}.$$

Let R_n be listed in increasing order as $r_{n,0}, r_{n,1}, \dots$. We require that, for all n and all sufficiently large k , exactly one of $r_{n,2k}$ and $r_{n,2k+1}$ is in A . This clearly implies that $\rho(A \cap R_n) = \frac{\rho(R_n)}{2}$.

Let N_e be the requirement that $\bar{\rho}(A \triangle \Phi_e) > 0$ if Φ_e is total. So, if N_e is met, then A is not coarsely computable via Φ_e . We will define a ternary computable function $g(e, i, s)$ to help us meet this requirement by “threatening” to witness that C is low. Let $N_{e,i}$ be the requirement that either N_e is met or $C'(i) = \lim_s g(e, i, s)$. Since C is not low, to meet N_e it suffices to meet all of its subrequirements $N_{e,i}$. Let $R_{e,i}$ denote $R_{\langle e,i \rangle}$. We use $R_{e,i}$ to meet $N_{e,i}$.

Fix e, i . Our module for satisfying $N_{e,i}$ proceeds as follows. Let s_0 be the least number so that $\Phi_{i,s_0}^{C_{s_0}}(i) \downarrow$; if there is no such number, then let $s_0 = \infty$. For each $s < s_0$, let $g(e, i, s) = 0$ and put s into A if s is of the form $r_{\langle e,i \rangle, 2k}$ for some k . If s_0 is infinite, that is if the search for s_0 fails, then no other work is done on $N_{e,i}$. (Note that in this case $\lim_s g(e, i, s) = 0 = C'(i)$, so $N_{e,i}$ is met.) Suppose s_0 is finite (that is, the search for s_0 succeeds). We choose an interval $I_0 \subseteq R_{e,i}$ as follows. Let I_0 be of the form $\{r_{\langle e,i \rangle, 2j}, \dots, r_{\langle e,i \rangle, 2k+1}\}$ so that $\min(I_0) > s_0$ and so that $\rho_m(I_0) \geq \rho_m(R_{e,i})/2$ where $m = r_{\langle e,i \rangle, 2k+1} + 1$. Let u_0 be the use of the computation $\Phi_{i,s_0}^{C_{s_0}}(i)$. Note that $u_0 < s_0$ by a standard convention and that no element of I_0 has been enumerated in A . We then restrain all elements of I_0 from entering A but continue putting alternate elements of $R_{e,i}$ above $\max I_0$ into A as before.

We then continue by searching for the least number $s_1 > s_0$ so that $\Phi_{e,s_1}(x) \downarrow$ for every $x \in I_0$ or some number less than u_0 is enumerated into C at stage s_1 . If no such number s_1 exists, then let $s_1 = \infty$. Set $g(e, i, s) = 0$ whenever $s_0 \leq s < s_1$. If s_1 is infinite, then no other work is done on $N_{e,i}$. (In this case, N_e is met because Φ_e is not total.) Suppose s_1 is finite (that is, this search succeeds). There are two cases. First, suppose some number less than u_0 is enumerated in C at stage s_1 . We then have permission from C to enumerate numbers in I_0 into A . Accordingly, we cancel the restraint on I_0 and put $r_{\langle e,i \rangle, 2j'}$ into A whenever $j \leq j' \leq k$. In this case the interval I_0 has become useless to us, and we go back to our first step but now starting at stage s_1 . If we find a stage $s_2 \geq s_1$ with $\Phi_{i,s_2}^{C_{s_2}}(i) \downarrow$, say with use u_1 , we choose a new interval I_1 of the same form as before, but now with $\min(I_1) > s_2$ and proceed as before with I_1 in place of I_0 , and setting $g(e, i, s) = 0$ for $s_1 \leq s < s_2$.

Now, suppose no number smaller than u_0 is enumerated into C at s_1 . Then, $\Phi_{e,s_1}(x) \downarrow$ for all $x \in I_0$. We are now in a position to make progress on N_e provided

that C later permits us to change A on I_0 . We then search for the least number $s_2 \geq s_1$ so that some number less than u_0 is enumerated in C at stage s_2 . If there is no such number then let $s_2 = \infty$. We set $g(e, i, s) = 1$ whenever $s_1 \leq s < s_2$ in order to force C to give us the desired permission. If s_2 is infinite, then no other work is done on $N_{e,i}$. (In this case, we have $\lim_s g(e, i, s) = 1 = C'(i)$.) Suppose s_2 is finite (that is, this search succeeds). We then declare the interval I_0 to be *successful* and cancel the restraint on I_0 . Since a number smaller than $u_0 < \min(I_0)$ has now entered C , we have permission to enumerate elements of I_0 into A . So, for each $j \leq j' \leq k$ put exactly one of $r_{\langle e,i \rangle, 2j'}$, $r_{\langle e,i \rangle, 2j'+1}$ into A so that A and Φ_e differ on at least one of these numbers. (This ensures that at least half of the elements of I_0 are in $A \Delta \Phi_e$ and hence that $\rho_m(A \Delta \Phi_e) > \frac{\rho_m(R_{e,i})}{4}$ where $m = \max I_0 + 1$.) We now restart our process as above. We continue in this fashion, defining a sequence of intervals. Note that, in general, $g(e, i, s) = 1$ if at stage s the most recently chosen interval has been declared successful and we are awaiting a C -change below it, and otherwise $g(e, i, s) = 0$.

This strategy clearly succeeds if any of its searches fail, by the parenthetical remarks in the construction. Also, if there are infinitely many successful intervals, it ensures that $\bar{\rho}(A \Delta \Phi_e) \geq \frac{\rho(R_{e,i})}{4} > 0$, so N_e is met. If all searches are successful but there are only finitely many successful intervals, then $C'(i) = 0 = \lim_s g(e, i, s)$ and $N_{e,i}$ is met. Only finitely many elements of $R_{e,i}$ are permanently restrained from entering A (namely the elements of the final interval, if any), so $\rho(A) = \frac{1}{2}$ for reasons already given. \square

We now obtain the following from Proposition 4.1 and Theorem 4.2.

Corollary 4.3. *If \mathbf{a} is a c.e. degree, then \mathbf{a} is low if and only if every c.e. set in \mathbf{a} that has a density is coarsely computable.*

For an application of this result to a degree structure arising from the notion of coarse computability, see Hirschfeldt, Jockusch, Kuyper, and Schupp [6].

5. DENSITY, 1-GENERICITY, AND RANDOMNESS

As we have already mentioned, it is easily seen that every degree contains a set that is both coarsely computable and generically computable, and every nonzero degree contains a set with neither of these properties. On the other hand, the next two results show that for “most” degrees \mathbf{a} , every \mathbf{a} -computable set that is generically computable is also coarsely computable. A set A is called *1-generic* if for every c.e. set S of binary strings, A either meets or avoids S .

Theorem 5.1. *Let A be a 1-generic set and let $r \in [0, 1]$. Suppose that $B \leq_T A$ and B is partially computable at density r . Then B is coarsely computable at density r .*

Proof. Fix a Turing functional Φ with $B = \Phi^A$ and a partial computable function φ such that $\varphi(n) = B(n)$ for all n in the domain of φ , and $\rho(\text{dom } \varphi) \geq r$. Let

$$S = \{\sigma \in 2^{<\omega} : \Phi^\sigma \text{ is incompatible with } \varphi\}.$$

Then S is a c.e. set of strings so A either meets or avoids S . If A meets S , then B disagrees with φ on some argument, a contradiction. Hence A avoids S . Fix a string $\gamma \prec A$ such that no string extending γ is in S . Now define a computable set C as follows. Given n , search for a string σ extending γ such that $\Phi^\sigma(n) \downarrow$ and put $C(n) = \Phi^\sigma(n)$ for the first such σ that is found. Then C is total because A

extends γ and Φ^A is total. Hence C is a computable set. Further, if $\varphi(n)\downarrow$ then $B(n) = C(n)$ since no extension of γ is in S . Hence $C \nabla B \supseteq \text{dom } \varphi$, so $\underline{\rho}(C \nabla B) \geq r$, and hence B is coarsely computable at density r . \square

Corollary 5.2. *If A is 1-generic and $B \leq_{\text{T}} A$ is generically computable, then B is coarsely computable.*

We do not need the definition of n -randomness here, but we simply point out the easy result that if A is 1-random, then $\gamma(A) = \frac{1}{2}$. A set A is called *weakly n -random* if A does not belong to any Π_n^0 class of measure 0.

Theorem 5.3. (i) *If A is weakly 1-random, $B \leq_{\text{tt}} A$, and B is partially computable at density r , then B is coarsely computable at density r .*
(ii) *If A is weakly 2-random, $B \leq_{\text{T}} A$, and B is partially computable at density r , then B is coarsely computable at density r .*

Proof. To prove (i), fix a Turing functional Φ such that $B = \Phi^A$ and Φ^X is total for all $X \subseteq \omega$. Let φ be a partial computable function that witnesses that B is partially computable at density r , and define

$$P = \{X : \Phi^X \text{ is compatible with } \varphi\}.$$

Then P is a Π_1^0 class and $A \in P$, so $\mu(P) > 0$, where μ is Lebesgue measure. By the Lebesgue density theorem, there is a string γ such that $\frac{\mu(P \cap [\gamma])}{\mu([\gamma])} > .6$, where $[\gamma] = \{X \in 2^\omega : \gamma \prec X\}$. Define

$$C = \left\{ n : \frac{\mu(\{Z \succ \gamma : \Phi^Z(n) = 1\})}{\mu([\gamma])} \geq .5 \right\}.$$

Then it is easily seen that C is a computable set and $C \nabla B$ contains the domain of φ , so B is coarsely computable at density r .

To prove (ii), fix a Turing functional Φ with $B = \Phi^A$ and fix a partial computable function φ that witnesses that B is partially computable at density r . Define

$$P = \{X : \Phi^X \text{ is total and compatible with } \varphi\}.$$

Then P is a Π_2^0 class and $A \in P$, so $\mu(P) > 0$. Then for notational convenience assume that $\mu(P) > .8$, applying the Lebesgue density theorem as in part (a). It follows that for every n there exists $i \leq 1$ such that $\mu(\{X : \Phi^X(n) = i\}) \geq .4$. Given n , one can compute such an i effectively, and then put n into C if and only if $i = 1$. One can easily check that C is computable and $C \nabla B \supseteq \text{dom } \varphi$, so $\underline{\rho}(C \nabla B) \geq \underline{\rho}(\text{dom } \varphi) \geq r$. Hence B is coarsely computable at density r . \square

Note that 1-randomness does not suffice in part (ii) of the above theorem, since every set is computable from some 1-random set.

Since the 1-generic sets are comeager and the weakly 2-generic sets have measure 1, it follows from the last two theorems that generic computability implies coarse computability below almost every set, both in the sense of Baire category and in the sense of measure. The next result contrasts with this fact.

Theorem 5.4. *If the degree \mathbf{a} is hyperimmune, there is a set $B \leq_{\text{T}} A$ such that B is bi-immune and of density 0.*

We omit the proof, which is an easy variation of Jockusch's proof in [7], Theorem 3, that every hyperimmune set computes a bi-immune set.

Bienvenu, Day, and Hölzl [2] proved the beautiful theorem that every nonzero Turing degree contains an absolutely undecidable set A ; that is, a set such that every partial computable function that agrees with A on its domain has a domain of density 0. We now consider the degrees of sets that are both absolutely undecidable and coarsely computable.

Corollary 5.5. *In the sense of Lebesgue measure, almost every set A computes a set B that is absolutely undecidable and coarsely computable.*

Proof. D. A. Martin (see [3, Theorem 8.21.1]) proved that almost every set has hyperimmune degree. It is obvious that every bi-immune set is absolutely undecidable. \square

On the other hand, Gregory Igusa has proved the following theorem using forcing with computable perfect trees.

Theorem 5.6 (Igusa, to appear). *There is a noncomputable set A such that no set $B \leq_T A$ is both coarsely computable and absolutely undecidable.*

We now turn to studying the degrees of sets A such that $\gamma(A) = 1$ but A is not coarsely computable. As shown in Theorem 3.3, if either $\mathbf{a} \not\leq \mathbf{0}'$ or \mathbf{a} is a nonzero c.e. degree, then \mathbf{a} contains such a set. This observation might lead one to conjecture that every nonzero degree computes such a set, but we shall prove the opposite for Δ_2^0 1-generic degrees. We will reach this result by first considering sets for which $\gamma(A) = 1$ is witnessed constructively.

Definition 5.7. We say that $\gamma(A) = 1$ *constructively* if there is a uniformly computable sequence of computable sets C_0, C_1, \dots such that $\bar{\rho}(A \Delta C_n) < 2^{-n}$ for all n .

Of course, if A is coarsely computable, then $\gamma(A) = 1$ constructively. Although the converse appears unlikely, it was proved by Joe Miller.

Theorem 5.8 (Joe Miller, private communication). *If $\gamma(A) = 1$ constructively, then A is coarsely computable.*

Proof. We present Miller's proof in essentially the form in which he gave it. Let I_k be the interval $[2^k - 1, 2^{k+1} - 1)$. For any set C , let $d_k(C)$ be the density of C on I_k , so $d_k(C) = \frac{|C \cap I_k|}{2^k}$. The following lemma, which will also be useful in the proof of Theorem 5.11, relates $\bar{\rho}(C)$ to $\bar{d}(C)$, where $\bar{d}(C) = \limsup_k d_k(C)$.

Lemma 5.9. *For every set C ,*

$$\frac{\bar{d}(C)}{2} \leq \bar{\rho}(C) \leq 2\bar{d}(C).$$

Proof. For all k ,

$$d_k(C) = \frac{|C \cap I_k|}{2^k} \leq \frac{|C \upharpoonright (2^{k+1} - 1)|}{2^k} \leq 2\rho_{2^{k+1}-1}(C).$$

Dividing both sides of this inequality by 2 and then taking the lim sup of both sides yields that $\frac{\bar{d}(C)}{2} \leq \bar{\rho}(C)$.

To prove that $\bar{\rho}(C) \leq 2\bar{d}(C)$, assume that $k - 1 \in I_n$, so $2^n \leq k < 2^{n+1}$. Then

$$\rho_k(C) = \frac{|C \upharpoonright k|}{k} \leq \frac{|C \upharpoonright (2^{n+1} - 1)|}{2^n} = \frac{\sum_{0 \leq i \leq n} 2^i d_i(C)}{2^n} < 2 \max_{i \leq n} d_i(C).$$

Let $\epsilon > 0$ be given. Then $d_i(C) < \bar{d}(C) + \epsilon$ for all sufficiently large i . Hence there is a finite set F such that $d_i(C \setminus F) < \bar{d}(C \setminus F) + \epsilon$ for all i . Then, by the above inequality applied to $C \setminus F$, we have $\rho_k(C \setminus F) < 2(\bar{d}(C \setminus F) + \epsilon)$ for all k , so $\bar{\rho}(C \setminus F) \leq 2\bar{d}(C \setminus F)$. As $\bar{\rho}$ and \bar{d} are invariant under finite changes of their arguments and $\epsilon > 0$ is arbitrary, it follows that $\bar{\rho}(C) \leq 2\bar{d}(C)$. \square

We now complete the proof of Theorem 5.8. Let the sequence C_n witness that $\gamma(A) = 1$ constructively, so $\{C_n\}$ is uniformly computable and $\bar{\rho}(A\Delta C_n) < 2^{-n}$ for all n . It follows from the lemma that $\bar{d}(A\Delta C_n) < 2^{-n+1}$. Hence, for each n , if k is sufficiently large, we have $d_k(A\Delta C_n) < 2^{-n+1}$.

For $m < n$, we say that C_m trusts C_n on I_k if $d_k(C_n\Delta C_m) < 2^{-m+2}$. We say that C_n is trusted on I_k if C_m trusts C_n for all $m < n$. Note that C_0 is trusted on every interval I_k . We now define a computable set C that will witness that A is coarsely computable. For each k , let $N \leq k$ be maximal such that C_N is trusted on I_k , and let $C \upharpoonright I_k = C_N \upharpoonright I_k$.

We claim that $\rho(A\Delta C) = 0$. Fix n . Let $k \geq n$ be large enough that $d_k(A\Delta C_m) < 2^{-m+1}$ for all $m \leq n$. Then $d_k(C_n\Delta C_m) \leq d_k(A\Delta C_n) + d_k(A\Delta C_m) < 2^{-n+1} + 2^{-m+1} < 2^{-m+2}$ for all $m < n$. Therefore, C_n is trusted on I_k . Hence $C \upharpoonright I_k = C_N \upharpoonright I_k$ for some $N \geq n$ such that C_N is trusted on I_k . Therefore, C_n trusts C_N on I_k , so $d_k(C_n\Delta C_N) < 2^{-n+2}$. It follows that $d_k(A\Delta C) = d_k(A\Delta C_N) \leq d_k(A\Delta C_n) + d_k(C_n\Delta C_N) < 2^{-n+1} + 2^{-n+2} < 2^{-n+3}$. Because this is true for every sufficiently large k , we have $\bar{d}(A\Delta C) \leq 2^{-n+3}$. Since n was arbitrary, it follows that $\bar{d}(A\Delta C) = 0$ and hence, by the lemma, $\rho(A\Delta C) = 0$. Thus A is coarsely computable. \square

Corollary 5.10. *Suppose there is a $0'$ -computable function f such that, for all e , we have that $\Phi_{f(e)}$ is total and $\{0, 1\}$ -valued, and $\bar{\rho}(A\Delta\Phi_{f(e)}) \leq 2^{-e}$. Then A is coarsely computable.*

Proof. By the theorem, it suffices to show that $\gamma(A) = 1$ constructively. Let g be a computable function such that $f(e) = \lim_s g(e, s)$. We now define a computable function h such that, for all e , we have that $\Phi_{h(e)}$ is total and differs on only finitely many arguments from $\Phi_{f(e)}$, so that $\Phi_{h(0)}, \Phi_{h(1)}, \dots$ witnesses that $\gamma(A) = 1$ constructively. To compute $\Phi_{h(e)}(n)$, search for $s \geq n$ such that $\Phi_{g(e,s)}(n)$ converges in at most s many steps, and let $\Phi_{h(e)}(n) = \Phi_{g(e,s)}(n)$. The s - m - n theorem gives us such an h , and clearly h has the desired properties. \square

We now have the tools to prove the following result, which we did not initially expect to be true.

Theorem 5.11. *Let G be a Δ_2^0 1-generic set, and suppose that $A \leq_T G$ and $\gamma(A) = 1$. Then A is coarsely computable.*

Proof. Fix Φ such that $A = \Phi^G$. As in the proof of Theorem 5.8 let I_k be the interval $[2^k - 1, 2^{k+1} - 1)$ and define $d_k(C) = \frac{|C \upharpoonright I_k|}{2^k}$ and $\bar{d}(C) = \limsup_k d_k(C)$.

Consider first the case that for some $\epsilon > 0$ and for every computable set C and every number k , we have that G meets the set $S_{\epsilon, C, k}$ of strings defined below:

$$S_{\epsilon, C, k} = \{\nu : (\exists l > k)[d_l(\Phi^\nu \Delta C) \geq \epsilon]\}.$$

Of course, ν must be such that $\Phi^\nu(j) \downarrow$ for all $j \in I_l$ for the above to make sense. We claim that $\gamma(A) < 1$ in this case, so that this case cannot arise. Let C be a

computable set and fix ϵ as in the case hypothesis. Then, for every k there exists $l > k$ such that $d_l(A\Delta C) \geq \epsilon$ by the choice of ϵ . It follows that $\bar{d}(A\Delta C) \geq \epsilon$, so $\bar{\rho}(A\Delta C) \geq \frac{\epsilon}{2}$ by Lemma 5.9. By Lemma 1.9 it follows that $\underline{\rho}(A\nabla C) \leq 1 - \frac{\epsilon}{2}$. Hence $\gamma(A) \leq 1 - \frac{\epsilon}{2} < 1$. Since $\gamma(A) = 1$ by assumption, this case cannot arise.

Since G is 1-generic, it follows that for every n there is a computable set C and a number k such that G avoids $S_{2^{-(n+2)}, C, k}$; i.e., there exists $\gamma \prec G$ such that γ has no extension in $S_{2^{-(n+2)}, C, k}$. Given $l \geq k$, let ν_0 and ν_1 be strings extending γ such that $\Phi^{\nu_i}(x) \downarrow$ for all $x \in I_l$ and $i \leq 1$. Then

$$d_l(\Phi^{\nu_0} \Delta \Phi^{\nu_1}) \leq d_l(\Phi^{\nu_0} \Delta C) + d_l(C \Delta \Phi^{\nu_1}) < 2^{-(n+2)} + 2^{-(n+2)} = 2^{-(n+1)}.$$

Since G is Δ_2^0 , using an oracle for $0'$ we can find γ_n and k_n such that for all ν_0, ν_1 extending γ_n and all $l \geq k_n$, if $\Phi^{\nu_i}(x) \downarrow$ for all $x \in I_l$ and $i \leq 1$ then $d_l(\Phi^{\nu_0} \Delta \Phi^{\nu_1}) \leq 2^{-(n+1)}$. Note that if we take $\nu_0 \prec G$ then $d_l(\Phi^{\nu_1} \Delta A) < 2^{-(n+1)}$.

For each n , define a computable set B_n as follows. On each interval I_k search for $\nu_1 \not\prec \gamma_n$ such that Φ^{ν_1} converges on I_k . Note that such a ν_1 exists because $\gamma_n \prec G$ and Φ^G is total. Let $B_n \upharpoonright I_k = \Phi^{\nu_1} \upharpoonright I_k$. Then B_n is a computable set, since the only non-effective part of its definition is the use of the *single* string γ_n . Furthermore, an index for B_n as a computable set can be effectively computed from γ_n and hence from $0'$.

We claim that $\bar{\rho}(B_n \Delta A) \leq 2^{-n}$. Fix n . By Lemma 5.9, it suffices to show that $\bar{d}(B_n \Delta A) \leq 2^{-(n+1)}$. For all k , we have that $d_k(B_n \Delta A) = d_k(\Phi^{\nu_1} \Delta A)$ for some string ν_1 extending γ_n . Hence, if k is sufficiently large, it follows that $d_k(B_n \Delta A) \leq 2^{-n+1}$, and hence $\bar{d}(B_n \Delta A) \leq 2^{-(n+1)}$, so $\bar{\rho}(B_n \Delta A) \leq 2^{-n}$. It now follows from Corollary 5.10 with $\Phi_{f(\epsilon)} = B_\epsilon$ that A is coarsely computable. \square

6. FURTHER RESULTS

In this section we investigate the complexity of $\gamma(A)$ as a real number when A is c.e. and look at the distribution of values of $\gamma(B)$ as B ranges over all sets computable from a given set A . A real is *left- Σ_3^0* if $\{q \in \mathbb{Q} : q < r\}$ is Σ_3^0 .

Proposition 6.1. *If A is a c.e. set, then $\gamma(A)$ is a left- Σ_3^0 real.*

Proof. Let A be a c.e. set, and let q be a rational number with $q \neq \gamma(A)$. Then the following two statements are equivalent:

- (i) $q < \gamma(A)$.
- (ii) There is a computable set C such that $\rho_n(A\nabla C) \geq q$ for all n .

It is immediate that (ii) implies (i) since (ii) implies that A is coarsely computable at density q and hence $q \leq \gamma(A)$.

Now assume (i) in order to prove (ii). Let r be a real number with $q < r < \gamma(A)$. Then A is coarsely computable at density r , so there is a computable set C such that $A\nabla C$ has lower density at least r . Since $q < r$, it follows that $\rho_n(A\nabla C) \geq r$ for all but finitely many n . By making a finite change in C , we can ensure that this inequality holds for all n .

Routine expansion shows that the set of rational numbers q satisfying (ii) is Σ_3^0 , so A is left- Σ_3^0 by definition.

Note: The formulation of (ii) was chosen in order to minimize the number of quantifiers when it is expanded. If we proceeded by simply using the fact that, for $q \neq \gamma(A)$, we have that $q < \gamma(A)$ if and only if A is coarsely computable at density

q and used a routine expansion of the latter, we could conclude only that $\gamma(A)$ is left- Σ_5^0 . \square

In the next result, we prove the converse and thus characterize the reals of the form $\gamma(A)$ for A c.e.

Theorem 6.2. *Suppose $0 \leq r \leq 1$. Then the following are equivalent:*

- (i) $r = \gamma(A)$ for some c.e. set A .
- (ii) r is left- Σ_3^0 .

Proof. It was shown in the previous proposition that (i) implies (ii), so it remains to be shown that (ii) implies (i). Let r be left- Σ_3^0 . Our proof is based on that of Theorem 5.7 of [5], which shows that r is the lower density of some c.e. set. That proof consists in taking a Δ_2^0 set B such that $\underline{\rho}(B) = r$ (which exists by the relativized form of Theorem 5.1 of [5]) and constructing a strictly increasing Δ_2^0 function t and a c.e. set A such that for each n ,

- (1) $\rho_{t(n)}(A) = \rho_n(B)$
- (2) $A \cap [t(n), t(n+1))$ is an initial segment of $[t(n), t(n+1))$.

It then follows easily that $\underline{\rho}(A) = \underline{\rho}(B) = r$.

Let S be the set of all pairs (k, e) such that $e \leq k$. Let f be a computable bijection between S and ω . We can easily adapt the proof of Theorem 5.7 of [5] to replace (1) by

- (1') $\rho_{t(f(k,e))}(A) = \rho_k(B)$ for each k and $e \leq k$,

while still having (2) hold for each n . Furthermore, we can also ensure that when a new approximation $t(n, s+1)$ to $t(n)$ is defined, it is chosen to be greater than both $2^{t(n-1, s+1)}$ and $2^{t(s, s)}$ (because for each instance of Lemma 5.8 of [5], there are infinitely many c witnessing the truth of the lemma).

We now define a c.e. set C . At stage s , proceed as follows for each pair (k, e) with $f(k, e) \leq s$. Let $n = f(k, e)$. If $\Phi_{e,s}(x) \downarrow$ for all $x \in [t(n-1, s), t(n, s))$, then for each such x for which $\Phi_e(x) = 0$, enumerate x into C (if x is not already in C). We say that x is put into C for the sake of (k, e) .

Let $D = A \cup C$. Then D is a c.e. set, and $\underline{\rho}(D) \geq \underline{\rho}(A) = r$. By Theorem 3.9 of [5], for each $\epsilon > 0$, there is a computable subset of D with lower density greater than $r - \epsilon$. It follows that $\gamma(D) \geq r$.

Now let e be such that Φ_e is total. Fix a k and let $n = f(k, e)$. Let s be least such that $t(n, s+1) = t(n)$. Every number put into C by the end of stage s is less than $t(s, s)$. Every number put into C after stage s for the sake of any pair other than (k, e) is either less than $t(n-1) = t(n-1, s+1)$ or greater than or equal to $t(n)$. By our assumption on the size of $t(n)$, it follows that $C(x) \neq \Phi_e(x)$ for every $x \in [\log_2 t(n), t(n))$, so $\rho_{t(n)}(C \nabla \Phi_e) \leq \frac{\log_2 t(n)}{t(n)}$, and hence

$$\begin{aligned} \rho_{t(n)}(D \nabla \Phi_e) &\leq \rho_{t(n)}(C \nabla \Phi_e) + \rho_{t(n)}(D \nabla C) \\ &\leq \frac{\log_2 t(n)}{t(n)} + \rho_{t(n)}(A) = \frac{\log_2 t(n)}{t(n)} + \rho_k(B). \end{aligned}$$

Since $\lim_n \frac{\log_2 t(n)}{t(n)} = 0$, we have $\underline{\rho}(D \nabla \Phi_e) \leq \underline{\rho}(B) = r$. Since e is arbitrary, $\gamma(D) \leq r$. \square

Definition 6.3. If $A \subseteq \mathbb{N}$ we call

$$S(A) = \{\gamma(B) : B \leq_T A\} \subseteq [0, 1]$$

the *coarse spectrum* of A .

Theorem 6.4. For any set A and any Δ_2^0 real $s \in [0, 1]$, we have that $s \cdot \gamma(A) + (1 - s) \in S(A)$. It follows that $S(A)$ is dense in the interval $[\gamma(A), 1]$.

Proof. We may assume that $s > 0$, since any computable $B \leq_T A$ witnesses the fact that $1 \in S(A)$. By Theorem 2.21 of [9] there is a computable set R of density s . Note that R is infinite. Let h be an increasing computable function with range R , and let $B = h(A)$. Then $B \leq_T A$, so it suffices to prove that $\gamma(B) = s \cdot \gamma(A) + (1 - s)$. For this, we need the following lemma, which relates the lower density of $h(X)$ to that of X . The corresponding lemma for density was proved as Lemma 3.4 of [4], and the proof here is almost the same.

Lemma 6.5. Let h be a strictly increasing function and let $X \subseteq \omega$. Then $\underline{\rho}(h(X)) = \underline{\rho}(\text{range}(h))\underline{\rho}(X)$, provided that the range of h has a density.

Proof. Let Y be the range of h , and for each u , let $g(u)$ be the least k such that $h(k) \geq u$. As shown in the proof of Lemma 3.4 of [4], $\rho_u(h(X)) = \rho_u(Y)\rho_{g(u)}(X)$ for all u , via bijections induced by h . Taking the lim inf of both sides and using the fact that $\rho(Y)$ exists, we see that

$$\underline{\rho}(h(X)) = \rho(Y)(\liminf \langle \rho_{g(0)}(X), \rho_{g(1)}(X), \dots \rangle).$$

It is easily seen that the function g is finite-one and $g(h(x)) = x$ for all x , and $g(u + 1) \leq g(u) + 1$ for all u . Hence the sequence on the right-hand side of the above equation can be obtained from the sequence $\rho_0(X), \rho_1(X), \dots$ by replacing each term by a finite, nonempty sequence of terms with the same value. Thus the two sequences have the same lim inf, and we obtain $\underline{\rho}(h(X)) = \rho(Y)\underline{\rho}(X)$, as needed to prove the lemma. \square

To prove that $\gamma(B) = s \cdot \gamma(A) + (1 - s)$, it suffices to show that for each $t \in [0, 1]$, A is coarsely computable at density t if and only if B is coarsely computable at density $st + 1 - s$. Suppose first that A is coarsely computable at density t , and let C be a computable set such that $\underline{\rho}(A \nabla C) \geq t$. Let $\widehat{C} = h(C) \cup \overline{R}$. Then \widehat{C} is a computable set and

$$\underline{\rho}(\widehat{C} \nabla B) = \underline{\rho}(h(C \nabla A) \cup \overline{R}) \geq \underline{\rho}(h(C \nabla A)) + \underline{\rho}(\overline{R}) = s\underline{\rho}(C \nabla A) + 1 - s \geq s \cdot t + (1 - s).$$

It follows that B is coarsely computable at density $st + (1 - s)$.

Conversely, if a computable set \widehat{C} witnesses that B is coarsely computable at density $st + (1 - s)$, let $C = h^{-1}(\widehat{C})$, and check by a similar argument that C witnesses that A is coarsely computable at density t since $s > 0$. \square

REFERENCES

- [1] U. Andrews, M. Cai, D. Diamondstone, C. Jockusch, and S. Lempp, *Asymptotic density, computable traceability, and 1-randomness*, in preparation.
- [2] L. Bienvenu, A. Day, and R. Hölzl, *From bi-immunity to absolute undecidability*, Journal of Symbolic Logic **78** (2013), 1218–1228.
- [3] R. G. Downey and D. R. Hirschfeldt, *Algorithmic Complexity and Randomness*, Theory and Applications of Computability, Springer, New York, 2010.
- [4] R. G. Downey, C. G. Jockusch, Jr., T. H. McNicholl, and P. E. Schupp, *Asymptotic density and the Ershov hierarchy*, Mathematical Logic Quarterly, to appear.

- [5] R. G. Downey, C. G. Jockusch, Jr., and P. E. Schupp, *Asymptotic density and computably enumerable sets*, Journal of Mathematical Logic **13** (2013), 1350005 (43 pages).
- [6] D. R. Hirschfeldt, C. G. Jockusch, Jr., R. Kuyper, and P. E. Schupp, Coarse reducibility and algorithmic randomness, in preparation.
- [7] C. G. Jockusch, Jr., *The degrees of bi-immune sets*, Zeitschrift für Mathematische Logik und Grundlagen der Mathematik **15** (1969), 135–140.
- [8] C. G. Jockusch, Jr., *Degrees in which the recursive sets are uniformly recursive*, Canadian Journal of Mathematics **24** (1972), 1092–1099.
- [9] C. G. Jockusch, Jr. and P. E. Schupp, *Generic computability, Turing degrees, and asymptotic density*, Journal of the London Mathematical Society, Second Series **85** (2012), 472–490.
- [10] I. Kapovich, A. Myasnikov, P. Schupp, and V. Shpilrain, *Generic-case complexity, decision problems in group theory, and random walks*, Journal of Algebra **264** (2003), 665–694.
- [11] S. A. Kurtz, *Notions of weak genericity*, Journal of Symbolic Logic **48** (1983), 764–770.
- [12] D. A. Martin, *Classes of recursively enumerable sets and degrees of unsolvability*, Zeitschrift für Mathematische Logik und Grundlagen der Mathematik **12** (1966), 295–310.
- [13] R. I. Soare, *Recursively Enumerable Sets and Degrees*, Perspectives in Mathematical Logic, Springer-Verlag, Berlin, 1987.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO

E-mail address: `drh@math.uchicago.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN

E-mail address: `jockusch@math.uiuc.edu`

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY

E-mail address: `mcnichol@iastate.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN

E-mail address: `schupp@illinois.edu`