

9-24-2012

## The Moist Soil Test for Potassium and Other Nutrients: What's It All About?

Antonio P. Mallarino

*Iowa State University*, [apmallar@iastate.edu](mailto:apmallar@iastate.edu)

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Soil Science Commons](#)

---

### Recommended Citation

Mallarino, Antonio P., "The Moist Soil Test for Potassium and Other Nutrients: What's It All About?" (2012). *Integrated Crop Management News*. 135.

<http://lib.dr.iastate.edu/cropnews/135>

**The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.**

---

# The Moist Soil Test for Potassium and Other Nutrients: What's It All About?

## **Abstract**

A new private soil-testing laboratory began operations this fall in Iowa (based in Ames), testing for most nutrients on non-dried soil samples. This has generated many questions concerning the procedure and interpretations of test results because the common lab procedure for most nutrients is to dry and grind the soil samples.

## **Keywords**

Agronomy

## **Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Soil Science

Subscribe to Crop News

#### Archives

[2014](#)

[2013](#)

[2012](#)

[2011](#)

[2010](#)

[2009](#)

[2008](#)

[Previous Years](#)

#### ISU Crop Resources

[Extension Field Agronomists](#)

[Crop & Soils Info](#)

[Pesticide Applicator Training](#)

[Agronomy Extension](#)

[Entomology Extension](#)

[Plant Pathology Extension](#)

[Ag and Biosystems Engineering Extension](#)

[Agribusiness Education Program](#)

[Iowa Grain Quality Initiative](#)

[College of Agriculture and Life Sciences](#)

[ISU Extension](#)

## Integrated Crop Management NEWS

-  [PRINT STORY](#)
-  [EMAIL STORY](#)
-  [ADD TO DELICIOUS](#)
-  [ATOM FEED](#)
-  [FOLLOW ON TWITTER](#)

### The Moist Soil Test for Potassium and Other Nutrients: What's It All About?

By Antonio P. Mallarino, Department of Agronomy

A new private soil-testing laboratory began operations this fall in Iowa (based in Ames), testing for most nutrients on non-dried soil samples. This has generated many questions concerning the procedure and interpretations of test results because the common lab procedure for most nutrients is to dry and grind the soil samples.

The idea for testing non-dried soil samples is not new. It has been known for decades that drying soil may affect the extraction and measurement of certain nutrients, especially potassium (K). However, drying soil is commonly done by labs because it used to be a more practical sample handling procedure, and it standardizes soil moisture across all conditions. Iowa State University (ISU) research during the 1960s and 1970s, mainly greenhouse trials but some field trials, had shown that testing non-dried (field-moist) soil samples provided a better estimate of K fertilizer needs than testing dried samples, but both procedures provided similar estimates for phosphorus (P). Therefore, testing field-moist samples was adopted by the ISU soil and plant analysis laboratory, and it was the standard procedure for P and K during the 1970s and 1980s.

Iowa State University discontinued field-moist soil testing in 1988 (I was a graduate student at the time); not because it was a bad procedure but because no other lab adopted it and only ISU soil test interpretations were based on moist testing. The ISU lab made that decision even though Iowa research had shown the moist test was better for K, and it was among the tests recommended for the North-Central region by the NCR-13 committee (North-Central Regional Committee for Soil Testing and Plant analysis). Since ISU discontinued moist soil testing in 1998, the NCR-13 committee dropped the procedure from its soil-test methods publication.

The general attitude about moist soil handling and testing changed considerably this year, when the private lab that began operations in Iowa developed a machine that easily handles moist samples and makes implementation of this procedure as practical as the common dry method (or even more practical because it avoids drying and grinding samples). This laboratory has been conducting soil testing research with Iowa soils since last year, some in collaboration with ISU.

Field research conducted during the 1990s showed much variability and uncertainty with K soil testing due to several reasons. So a portion of research my graduate students and I conducted since the early 2000s has focused on studying again if testing field-moist soil samples for K is better than testing dried samples.

The amount of extracted K is lower for the moist test than for the dry test at values usually optimum for crops or lower, but the difference decreases as levels increase. At extremely high levels, the moist test values can be higher than the dry test values. However, the moist and dry tests result in

approximately similar values for P (by the Bray-1, Olsen, or Mehlich-3 methods), calcium (Ca) and magnesium (Mg) (by the ammonium-acetate or Mehlich-3 methods), pH, and buffer pH (by the SMP or Sikora methods). Figure 1 shows, as an example, comparisons for K and P extracted by the Mehlich-3 method from moist and dry samples.




Figure 1. Comparison of amounts of soil K, P, and Mg extracted by the Mehlich-3 test from many soil samples taken from Iowa fields. The dotted diagonal line indicates an exact 1:1 ratio.

Results from more than 300 field-response trials for corn and soybean confirmed that the field-moist test for K is better. Figure 2 shows the relationship between soil K measured by moist or dry tests and the relative yield response of corn and soybean to K fertilization. The graph for the dry test shows the current ISU interpretations for K, and further information including recommended fertilization rates is available in Extension publication PM 1688, [A General Guide for Crop Nutrient and Limestone Recommendations in Iowa](#). The graph for the moist test shows the interpretations that ISU suggested for this test in the late 1980s. The fertilizer rate recommended for the Medium class used at the time was the amount to maintain soil-test values based on K removal, which in concept is similar to current Optimum class for the dry test. The old moist test classes relate to the recent yield responses almost exactly as they related to yield responses from research conducted during the 1980s (not shown).




Figure 2. Relationship between the relative yield response of corn and soybean and soil-test K measured from dried and moist soil samples (ammonium-acetate test, 6-inch sampling depth). VL, very low; L, low; O, optimum; M, medium; H, high; VH, very high.

Additional information and results will be presented at the North-Central Extension-Industry Soil Fertility Conference in Des Moines (November 14-15, 2012), the ISU Extension Integrated Crop Management Conference in Ames (November 28-29), and other ISU Extension conferences and workshops during the winter.

This fall the NCR-13 regional committee will publish an updated sample preparation chapter of the soil-test methods publication that includes the moist procedure. The ISU interpretations and fertilizer recommendations for the moist test will be developed in the future, as results for several ongoing field trials become available and can be merged with previous results. Interpretations for the moist test for P using Bray, Olsen, and Mehlich-3 (colorimetric or ICP procedures) should be similar to those for the dry test, since data already showed similar test results. The interpretations for the moist test for K likely will be similar to those suggested by ISU in the 1980s.

*Antonio Mallarino is a professor of agronomy with research and extension responsibilities in soil fertility and nutrient management. Mallarino can be reached at [apmallar@iastate.edu](mailto:apmallar@iastate.edu) or by calling (515) 294-6200.*

---

This article was published originally on 9/24/2012. The information contained within the article may or may not be up to date depending on when you are accessing the information.

---

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.

---

Copyright ©2014 [Iowa State University Extension](#) | [Iowa State University](#)  
[Contact us](#) | [For Staff](#) | [Nondiscrimination and Information Disclosures](#) | [CMS Admin](#)  
Last Updated 9/25/2012