

8-24-2012

Managing Western Corn Rootworm with Bt Corn

Aaron J. Gassmann

Iowa State University, aaronjg@iastate.edu

Erin W. Hodgson

Iowa State University, ewh@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Gassmann, Aaron J. and Hodgson, Erin W., "Managing Western Corn Rootworm with Bt Corn" (2012). *Integrated Crop Management News*. 124.

<http://lib.dr.iastate.edu/cropnews/124>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Managing Western Corn Rootworm with Bt Corn

Abstract

During the summer of 2011, the ISU Corn Entomology Lab visited several fields in Iowa in response to complaints for rootworm injury to Bt rootworm corn, and to follow-up on fields that were visited in 2010 in response to grower complaints. To date, resistance of western corn rootworm has been found only for Bt corn producing Cry3Bb1, which was originally released as YieldGard Rootworm and is now the rootworm trait found in VTtriple and VTtriple Pro. Cry3Bb1 also is present as a pyramid with Bt toxin Cry35/35Ab1 in SmartStax.

Keywords

Entomology

Disciplines

Agricultural Science | Agriculture | Entomology

search

Subscribe to Crop News

Archives

- [2014](#)
- [2013](#)
- [2012](#)
- [2011](#)
- [2010](#)
- [2009](#)
- [2008](#)

[Previous Years](#)

ISU Crop Resources

- [Extension Field Agronomists](#)
- [Crop & Soils Info](#)
- [Pesticide Applicator Training](#)
- [Agronomy Extension](#)
- [Entomology Extension](#)
- [Plant Pathology Extension](#)
- [Ag and Biosystems Engineering Extension](#)
- [Agribusiness Education Program](#)
- [Iowa Grain Quality Initiative](#)
- [College of Agriculture and Life Sciences](#)
- [ISU Extension](#)



- PRINT STORY
- EMAIL STORY
- ADD TO DELICIOUS
- ATOM FEED
- FOLLOW ON TWITTER

Managing Western Corn Rootworm with Bt Corn

By Aaron Gassmann and Erin Hodgson, Department of Entomology

During the summer of 2011, the ISU Corn Entomology Lab visited several fields in Iowa in response to complaints for rootworm injury to Bt rootworm corn, and to follow-up on fields that were visited in 2010 in response to grower complaints. To date, resistance of western corn rootworm has been found only for Bt corn producing Cry3Bb1, which was originally released as YieldGard Rootworm and is now the rootworm trait found in VTTriple and VTTriple Pro. Cry3Bb1 also is present as a pyramid with Bt toxin Cry35/35Ab1 in SmartStax.

Field visits in 2011 found some fields with severe root injury to Cry3Bb1 corn and to corn that produces Bt protein mCry3a, which is the Bt trait found in Agrisure (a Syngenta Product) and TRIsect (a Pioneer Product). We currently do not know the reason for the high level of rootworm injury observed for mCry3a rootworm corn, but this is an active area of research in the Gassmann Laboratory. So far in 2012, we have visited several fields with heavy rootworm feeding injury to Bt corn and patterns of injury are similar to 2011.

Table 1. Type of Bt toxin, level of root injury and number of fields visited in 2011.

Bt Toxin¹	Average Injury	Number of Fields Visited
Cry3Bb1	0 to 1 nodes	2
Cry3Bb1	1 to 2 nodes	3
Cry3Bb1	2 to 3 nodes	6
mCry3a	0 to 1 nodes	0
mCry3a	1 to 2 nodes	2
mCry3a	2 to 3 nodes	4
Cry34/35Ab1	0 to 1 nodes	2
Cry34/35Ab1	1 to 2 nodes	0
Cry34/35Ab1	2 to 3 nodes	0
Cry34/35Ab1 +Cry3Bb1	0 to 1 nodes	3
Cry34/35Ab1 +Cry3Bb1	1 to 2 nodes	0
Cry34/35Ab1 +Cry3Bb1	2 to 3 nodes	0

¹Cry3Bb1 is found in VTtriple and VTtriplePro (Monsanto); mCry3a is found in Agrisure (Syngenta) and TRIsect (Pioneer); Cry34/35Ab1 is found in Herculex (Dow and Pioneer); Cry34/35Ab1 + Cry3Bb1 is found in SmartStax (Dow and Monsanto).



Figure 1. A cornfield visited in Iowa during 2012 that was planted to Bt corn. The planting pattern in the photographed area alternated between 12 rows of Cry3Bb1 corn and 12 rows of mCry3a corn.

Data from field visits in 2011 and 2012 suggest that caution should be used when managing rootworm in continuous corn with either Cry3Bb1 traits or mCry3a traits. It is important to check continuous cornfields for rootworm injury. A high abundance of adult corn rootworm in a field is an indication of potential problems with larval management, but only digging and evaluating corn roots can confirm the presence of larval feeding injury. The relationship between root injury and yield is complicated and depends on several factors, such as soil moisture. A reasonable economic threshold for larval rootworm injury is 0.25 nodes, which means that ensuring less than 0.25 nodes of

injury on average is an economically sound management practice, but achieving further reductions in injury is not warranted.

As growers think about seed selection for 2013, they should be aware of current challenges encountered when managing corn rootworm with Bt corn, and use integrated pest management (IPM). Crop rotation, for example rotation to soybeans, is a highly effective way to manage western corn rootworm in Iowa. If farmers are concerned about extended diapause in northern corn rootworm, they can consider protecting first year corn with a soil-applied insecticide or Bt corn. In addition to crop rotation, growers should alternate their management tactics to control corn rootworm if they are growing continuous corn. Current options include rotation among single trait events, pyramided Bt events (corn that contains multiple traits targeting corn rootworm) and non-Bt corn with a soil-applied insecticide.

Aaron Gassmann is an assistant professor of entomology with research and teaching responsibilities in insect pest management. Erin Hodgson is an assistant professor of entomology with extension and research responsibilities; contact her at ewh@iastate.edu or phone 515-294-2847.

This article was published originally on 8/24/2012. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.