

2016

Enhanced destruction of *Salmonella enterica* in carrot and berry juices by a combination of cinnamaldehyde and high pressure processing

Aura Daraba
Iowa State University

Aubrey F. Mendonca
Iowa State University, amendon@iastate.edu

David Manu
Iowa State University

James S. Dickson
Iowa State University, jdickson@iastate.edu

Joseph G. Sebranek
Follow this and additional works at: https://lib.dr.iastate.edu/fshn_ag_pubs
Iowa State University, sebranek@iastate.edu

Part of the [Food Science Commons](#), [Human and Clinical Nutrition Commons](#), [Meat Science Commons](#), and the [Molecular, Genetic, and Biochemical Nutrition Commons](#)
See next page for additional authors

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/fshn_ag_pubs/166. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

Enhanced destruction of *Salmonella enterica* in carrot and berry juices by a combination of cinnamaldehyde and high pressure processing

Abstract

Statement of the Problem: In recent years several disease outbreaks were linked to unpasteurized juices contaminated with human enteric pathogens such as *Salmonella enterica* and *Escherichia coli* O157:H7 (CDC 2011; EFSA 2015). While heat pasteurization and canning can inactivate vegetative pathogens, such processes can destroy heat labile nutrients and negatively alter sensory characteristics of juices. Also growing consumer demands for foods which are nutritious, fresh-like, and devoid of synthetic preservatives, have forced juice manufacturers to explore non-thermal processes and natural antimicrobials for pathogen control in juices. The present study investigated the effect of low concentrations of cinnamaldehyde combined with high pressure processing (HPP) for killing *S. enterica* in carrot juice (CRJ) and a mixed berry juice (MBJ) at 4°C. **Methodology & Theoretical Orientation:** CRJ (pH 6.25) and MBJ (pH 3.59) with added cinnamaldehyde (0.10, 0.15 and 0.25 µl/ml) were inoculated with *S. enterica* (5-strain; final concentration ~10⁷ CFU/ml). Inoculated juices without added cinnamaldehyde served as control. Juices (4°C) were packaged in polyester pouches and pressurized (400 or 300 MPa) for 60, 90 and 120 seconds. The time between inoculation and HPP was approximately 1.5 hours. *Salmonella* survived for 42 days or more in control CRJ following HPP (400 MPa) for 30, 60, or 120 s. Addition of cinnamaldehyde to juices increased the sensitivity of *S. enterica* to HPP. Cinnamaldehyde (0.25 µl/ml) combined with 400 MPa (60 s) inactivated *S. enterica* by more than 5.5-log in CRJ. In MBJ, cinnamaldehyde (0.15 µl/ml) with a lower pressure (300 MPa for 120 s) resulted in complete inactivation (negative enrichment) and greater than a 5-log₁₀ CFU/ml reduction of *S. enterica*. **Conclusion & Significance:** The use of CA in conjunction with HPP has good potential to serve as an alternative process for heat pasteurization of juices and meet the 5-log reduction performance standard as stipulated in the juice HACCP regulations.

Disciplines

Food Science | Human and Clinical Nutrition | Meat Science | Molecular, Genetic, and Biochemical Nutrition

Comments

This abstract is published as Daraba[SLL[1], A., Mendonca, A., Manu, D., Dickson, J., Sebranek, J., Shaw, A., and A. DiSpirito. 2016. Enhanced destruction of *Salmonella enterica* in carrot and berry juices by a combination of cinnamaldehyde and high pressure processing. *Journal of Experimental Food Chemistry*: 2:4 (supplement). DOI: [10.4172/2472-0542.C1.006](https://doi.org/10.4172/2472-0542.C1.006).

Creative Commons License

Creative

Commons
Attribution-

Share

Authors

Anke

4.0

License

Aura Daraba, Aubrey F. Mendonca, David Manu, James S. Dickson, Joseph G. Sebranek, Angela M. Shaw, and Alan A. DiSpirito

10th Global Summit on

Food Safety, Processing & Technology

December 05-07, 2016 San Antonio, USA

Enhanced destruction of *Salmonella enterica* in carrot and berry juices by a combination of cinnamaldehyde and high pressure processing

Aura Daraba^{1,2}, Aubrey F Mendonca¹, David Manu¹, James S Dickson¹, Joseph Sebranek¹, Angela Shaw¹ and Alan DiSpirito¹¹Iowa State University, USA²University of Galati, Romania

Statement of the Problem: In recent years several disease outbreaks were linked to unpasteurized juices contaminated with human enteric pathogens such as *Salmonella enterica* and *Escherichia coli* O157:H7 (CDC 2011; EFSA 2015). While heat pasteurization and canning can inactivate vegetative pathogens, such processes can destroy heat labile nutrients and negatively alter sensory characteristics of juices. Also growing consumer demands for foods which are nutritious, fresh-like, and devoid of synthetic preservatives, have forced juice manufacturers to explore non-thermal processes and natural antimicrobials for pathogen control in juices. The present study investigated the effect of low concentrations of cinnamaldehyde combined with high pressure processing (HPP) for killing *S. enterica* in carrot juice (CRJ) and a mixed berry juice (MBJ) at 4°C.

Methodology & Theoretical Orientation: CRJ (pH 6.25) and MBJ (pH 3.59) with added cinnamaldehyde (0.10, 0.15 and 0.25 µl/ml) were inoculated with *S. enterica* (5-strain; final concentration ~107 CFU/ml). Inoculated juices without added cinnamaldehyde served as control. Juices (4°C) were packaged in polyester pouches and pressurized (400 or 300 MPa) for 60, 90 and 120 seconds. The time between inoculation and HPP was approximately 1.5 hours. *Salmonella* survived for 42 days or more in control CRJ following HPP (400 MPa) for 30, 60, or 120 s. Addition of cinnamaldehyde to juices increased the sensitivity of *S. enterica* to HPP. Cinnamaldehyde (0.25 µl/ml) combined with 400 MPa (60 s) inactivated *S. enterica* by more than 5.5-log in CRJ. In MBJ, cinnamaldehyde (0.15 µl/ml) with a lower pressure (300 MPa for 120 s) resulted in complete inactivation (negative enrichment) and greater than a 5-log₁₀ CFU/ml reduction of *S. enterica*.

Conclusion & Significance: The use of CA in conjunction with HPP has good potential to serve as an alternative process for heat pasteurization of juices and meet the 5-log reduction performance standard as stipulated in the juice HACCP regulations.

Biography

Aura Daraba has her expertise in Food Safety and Food Quality, use of natural antimicrobials to control pathogens, use of non-thermal food processing technologies, and implementation of HACCP in Food Industry and in Food Service Units. She has worked extensively along with Dr. Aubrey Mendonca and Dr. Angela Shaw in the use of High Pressure Processing and use of Natural Antimicrobials to control pathogens in foods.

aura.daraba@gmail.com

Notes: