

9-1997

Livestock Industry Facilities and Environment: Swine Breeding Systems—Effects of Heat Stress

Jay D. Harmon

Iowa State University, jharmon@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/extension_ag_pubs



Part of the [Agricultural Education Commons](#), and the [Bioresource and Agricultural Engineering Commons](#)

Recommended Citation

Harmon, Jay D., "Livestock Industry Facilities and Environment: Swine Breeding Systems—Effects of Heat Stress" (1997). *Agriculture and Environment Extension Publications*. 165.

http://lib.dr.iastate.edu/extension_ag_pubs/165

Iowa State University Extension and Outreach publications in the Iowa State University Digital Repository are made available for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current publications and information from Iowa State University Extension and Outreach, please visit <http://www.extension.iastate.edu>.

Swine Breeding Systems - *Effects of Heat Stress¹*



Livestock
Industry
Facilities &
Environment

The thermal environment in the breeding facility plays an important role in total herd management. This manifests itself in poor conception and farrowing rates that may not be detected until weeks or months past the occurrence of an improper thermal environment. Because of this, it is important to anticipate these potential lulls in breeding performance and counteract them by managing the breeding facility environment properly and adding additional open females to keep the facilities full.

High Temperature Effects

- Boar effects
 - Semen quality drops;
 - The effect begins 2 to 3 weeks after stress occurs and lasts until 8 or 9 weeks after the exposure.
 - Reduces libido.




Figure 1 Effect of Three Days of Heat Stress of Boars on Conception Rate, Roller, et al. (1973)

¹ Written by Jay D. Harmon, Ph.D., P.E., Extension Agricultural Engineer, Iowa State University, Ames, IA.

- Sow effects
 - Susceptible 2 or 3 weeks after breeding and just prior to farrowing;
 - Conception rate, litter size and litter weight may be affected.

Table 1 Effect of heat stress of sows on embryonic survival rate, (Tompkins, et al. 1967).

Day after mating heat stress began (5 day duration)	Control Sows	Stressed Sows
1	69%	39%
20	64%	77%

Table 2 Farrowing performance of gilts as affected by daily heat stress on days 102 through 110 of gestation, (Omtvedt et al. 1971).

Trait	Control Gilts	Stressed Gilts
No. Born Alive	10.4	6.0
No. Born Dead	0.4	5.2
Mean Birth Weight (kg)	1.4	1.2
Mean Litter Weight (kg)	13.6	8.6

Environmental Goal

- Average daily temperature of 80 F;
- Exposure above 84 F should be avoided.