

2018

The relationship between k -forcing and k -power domination

Daniela Ferrero
Texas State University

Leslie Hogben
Iowa State University, hogben@iastate.edu

Franklin H.J. Kenter
United States Naval Academy

Michael Young
Iowa State University, myoung@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/math_pubs



Part of the [Algebra Commons](#), and the [Discrete Mathematics and Combinatorics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/math_pubs/170. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Mathematics at Iowa State University Digital Repository. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

The relationship between k-forcing and k-power domination

Abstract

Zero forcing and power domination are iterative processes on graphs where an initial set of vertices are observed, and additional vertices become observed based on some rules. In both cases, the goal is to eventually observe the entire graph using the fewest number of initial vertices. The concept of k-power domination was introduced by Chang et al. (2012) as a generalization of power domination and standard graph domination. Independently, k-forcing was defined by Amos et al. (2015) to generalize zero forcing. In this paper, we combine the study of k-forcing and k-power domination, providing a new approach to analyze both processes. We give a relationship between the k-forcing and the k-power domination numbers of a graph that bounds one in terms of the other. We also obtain results using the contraction of subgraphs that allow the parallel computation of k-forcing and k-power dominating sets.

Keywords


k-power domination, k-forcing, Subgraph contraction, Sierpiński graphs

Disciplines

Algebra | Discrete Mathematics and Combinatorics

Comments

This is a manuscript of the article Ferrero, Daniela, Leslie Hogben, Franklin HJ Kenter, and Michael Young. "The relationship between k-forcing and k-power domination." *Discrete Mathematics* (2018): 1789-1797. DOI: [10.1016/j.disc.2017.10.031](https://doi.org/10.1016/j.disc.2017.10.031).



The relationship between k -forcing and k -power domination

Daniela Ferrero^{a,*}, Leslie Hogben^{b,c}, Franklin H.J. Kenter^d, Michael Young^b

^a Department of Mathematics, Texas State University, San Marcos, TX 78666, USA

^b Department of Mathematics, Iowa State University, Ames, IA 50011, USA

^c American Institute of Mathematics, 600 E. Brokaw Road, San Jose, CA 95112, USA

^d Department of Mathematics, United States Naval Academy, Annapolis, MD 21402, USA



ARTICLE INFO

Article history:

Received 21 March 2017

Received in revised form 19 October 2017

Accepted 28 October 2017

Available online 22 November 2017

Keywords:

k -power domination

k -forcing

Subgraph contraction

Sierpiński graphs

ABSTRACT

Zero forcing and power domination are iterative processes on graphs where an initial set of vertices are observed, and additional vertices become observed based on some rules. In both cases, the goal is to eventually observe the entire graph using the fewest number of initial vertices. The concept of k -power domination was introduced by Chang et al. (2012) as a generalization of power domination and standard graph domination. Independently, k -forcing was defined by Amos et al. (2015) to generalize zero forcing. In this paper, we combine the study of k -forcing and k -power domination, providing a new approach to analyze both processes. We give a relationship between the k -forcing and the k -power domination numbers of a graph that bounds one in terms of the other. We also obtain results using the contraction of subgraphs that allow the parallel computation of k -forcing and k -power dominating sets.

© 2017 Elsevier B.V. All rights reserved.

1. Introduction

Zero forcing was introduced as a process to obtain an upper bound for the maximum nullity of real symmetric matrices whose nonzero pattern of off-diagonal entries is described by a given graph [2]. The minimum rank problem was motivated by the inverse eigenvalue problem of a graph. Independently, zero forcing was introduced by mathematical physicists studying quantum systems [5]. Since its introduction, zero forcing has attracted the attention of a large number of researchers who find the concept useful to model processes in a broad range of disciplines. The need for a uniform framework for the analysis of the diverse processes where the notion of zero forcing appears led to the introduction of a generalization of zero forcing called k -forcing [3].

Amos et al. proposed k -forcing in [3] as the following graph coloring game. Assume the vertices of a graph are colored in two colors, say white and blue. Iteratively apply the following color change rule: if u is a blue vertex with at most k white neighbors, then change the color of all the neighbors of u to blue. Once this rule does not change the color of any vertex, if all vertices are blue, the original set of blue vertices is a k -forcing set of G . The original zero forcing is 1-forcing under this definition. Because the problem of deciding whether a graph admits a 1-forcing set of a given maximum size is NP-complete even if restricted to planar graphs [1, Theorem 2.3.1], the general problem of finding forcing sets cannot be solved algorithmically for large graphs without the development of further theoretical tools.

Power domination was introduced by Haynes et al. in [9] when using graph models to study the monitoring process of electrical power networks. When a power network is modeled by a graph, a power dominating set provides the locations

* Corresponding author.

E-mail addresses: dferrero@txstate.edu (D. Ferrero), hogben@iastate.edu, hogben@aimath.org (L. Hogben), kenter@usna.edu (F.H.J. Kenter), myoung@iastate.edu (M. Young).

where monitoring devices (Phase Measurement Units, or PMUs for short) can be placed in order to monitor the power network. Finding optimal PMU placements is an important practical problem in electrical engineering due to the cost of PMUs and network size. Although power domination is substantially different from standard graph domination, the notion of k -power domination was proposed as a generalization of both power domination ($k = 1$) and standard graph domination ($k = 0$) [6].

Chang et al. defined k -power domination in [6] using sets of *observed* vertices. Given a graph G and a set of vertices S , initially all vertices in S and their neighbors are observed; all other vertices are unobserved. Iteratively apply the following propagation rule: if there exists an observed vertex u that has k or fewer unobserved neighbors, then all the neighbors of u are observed. Once this rule does not produce any additional observed vertices, if all vertices of G are observed, S is a k -power dominating set of G . Many problems outside graph theory can be formulated in terms of minimum k -power dominating sets [6] so methods to obtain them are highly desired. An algorithmic approach has been attempted, but the problem of deciding if a graph admits a k -power dominating set of a given maximum size is NP-complete [6].

Although k -forcing and k -power domination have been studied independently, an in-depth analysis of k -power domination leads to the study of k -forcing. Indeed, after the initial step in which a set observes itself and its neighbors, the observation process in k -power domination proceeds exactly as the color changing process in k -forcing. The aim of this paper is to establish a precise connection between k -forcing and k -power domination to facilitate the transference of results, proofs, and methods between them, and ultimately to advance research on both problems.

Throughout this paper we work on k -forcing and k -power domination concurrently, using results in one process as stepping stones for results in the other one. In Section 2 we present the definitions and notation that we use in the rest of the paper. In Section 3 we give some core results and remarks that we use in the sections that follow.

In Section 4 we examine the effect of subgraph contraction in k -power domination and k -forcing. We obtain upper and lower bounds for the change in the k -power domination number produced by the contraction of a subgraph. Note that the contraction of a subgraph can increase or decrease its k -power domination number. In particular, we prove that the contraction of subgraphs of small degree can change the k -power domination number by at most one. In this section we also propose a way to decompose a graph in order to bound its k -power domination number in terms of that of smaller subgraphs. This can allow computation of k -power dominating sets to run in parallel. We also give the analogous results for k -forcing.

In Section 5 we present a lower bound for the k -power domination number of a graph in terms of its k -forcing number. This bound generalizes a known result for $k = 1$ that gives the only lower bound for the power domination number of an arbitrary graph available so far [4]. As an application, we find an upper bound for the k -forcing number of a graph in terms of its maximum degree.

2. Definitions and notation

A *graph* is an ordered pair $G = (V, E)$ where $V = V(G)$ is a finite nonempty set of *vertices* and $E = E(G)$ is a set of unordered pairs of distinct vertices called *edges*. All graphs considered in this paper are simple and undirected. The *order* of G is $|G| = |V(G)|$. Two vertices u and v are *adjacent* or *neighbors* in G if $\{u, v\} \in E(G)$. The *(open) neighborhood* of a vertex v is the set $N_G(v) = \{u \in V : \{u, v\} \in E\}$, and the *closed neighborhood* of v is the set $N_G[v] = N_G(v) \cup \{v\}$. Similarly, for any set of vertices S , $N_G(S) = \cup_{v \in S} N_G(v)$ and $N_G[S] = \cup_{v \in S} N_G[v]$. The *degree* of a vertex v is $\deg_G(v) = |N(v)|$. The *maximum* and *minimum degree* of G are $\Delta(G) = \max\{\deg_G(v) : v \in V\}$ and $\delta(G) = \min\{\deg_G(v) : v \in V\}$, respectively; a graph G is *regular* if $\delta(G) = \Delta(G)$. We will omit the subscript G when the graph G is clear from the context.

A path joining $u, v \in V$ is a sequence of distinct vertices $u = x_0, x_1, \dots, x_r = v$ such that $\{x_i, x_{i+1}\} \in E$ for each $i = 0, \dots, r - 1$. A graph G is *connected* if there is a path joining every pair of different vertices. If a graph is not connected, each maximal connected subgraph is a *component* of G . In this paper, $c(G)$ denotes the *number of components* of G and $G_1, \dots, G_{c(G)}$ denote the components of G . Most of the results in this work are given for connected graphs, since if a graph is not connected, we can apply the results to each component.

If X is a set of vertices of G , the *subgraph induced by X* (in G) is denoted as $G[X]$; it has vertex set X and edge set $\{\{u, v\} \in E : u, v \in X\}$. The graph $G - X$ is defined as $G[V \setminus X]$. The *contraction of X in G* is the graph G/X obtained by adding a vertex v_X to $G - X$ with $N_{G/X}(v_X) = N_G[X] \setminus X$. Note that G/X does not require $G[X]$ to be connected whereas the standard use of graph contraction does.

In a graph $G = (V, E)$, consider an arbitrary coloring of its vertices in two colors, say blue and white, and let T denote the set of blue vertices. For a positive integer k , the color changing process in k -forcing can be formally described by associating to T the family of sets $(\mathcal{F}_{G,k}^i(T))_{i \geq 0}$ recursively defined by the following rules.

1. $\mathcal{F}_{G,k}^0(T) = T$,
2. $\mathcal{F}_{G,k}^{i+1}(T) = \mathcal{F}_{G,k}^i(T) \cup \{N_G(v) : v \in \mathcal{F}_{G,k}^i(T) \text{ and } 1 \leq |N_G(v) \setminus \mathcal{F}_{G,k}^i(T)| \leq k\}$, for $i \geq 0$.

A set $T \subseteq V$ is a *k -forcing set* of G if there is an integer t such that $\mathcal{F}_{G,k}^t(T) = V$. A *minimum k -forcing set* is a k -forcing set of minimum cardinality. The *k -forcing number* of G is the cardinality of a minimum k -forcing set and is denoted by $Z_k(G)$. If $v \in \mathcal{F}_{G,k}^i(T)$ and $1 \leq |N_G(v) \setminus \mathcal{F}_{G,k}^i(T)| \leq k$ then v is said to *k -force* (or simply *force* if k is clear from the context) every vertex in $N_G(v) \setminus \mathcal{F}_{G,k}^i(T)$.

Let k be a positive integer. The definition of k -power domination on a graph G will be given in terms of a family of sets, $(\mathcal{P}_{G,k}^i(S))_{i \geq 0}$, associated to each set of vertices S in G .

1. $\mathcal{P}_{G,k}^0(S) = N_G[S]$,
2. $\mathcal{P}_{G,k}^{i+1}(S) = \mathcal{P}_{G,k}^i(S) \cup \{N_G(v) : v \in \mathcal{P}_{G,k}^i(S) \text{ and } 1 \leq |N_G(v) \setminus \mathcal{P}_{G,k}^i(S)| \leq k\}$, for $i \geq 0$.

A set $S \subseteq V$ is a *k-power dominating set* of G if there is an integer ℓ such that $\mathcal{P}_{G,k}^\ell(S) = V$. A *minimum k-power dominating set* is a *k-power dominating set* of minimum cardinality. The *k-power domination number* of G is the cardinality of a minimum *k-power dominating set* and is denoted by $\gamma_{p,k}(G)$.

Next we recall the definition of standard graph domination. A vertex v *dominates* all vertices in $N_G[v]$. A set $S \subseteq V$ is a *dominating set* of G if $N_G[S] = V$. The minimum cardinality of a dominating set is the *domination number* of G , denoted by $\gamma(G)$.

Note that 1-forcing coincides with zero forcing [3], while 1-power domination is exactly power domination and 0-power domination coincides with domination [6].

3. Preliminaries

The following observations follow directly from the definitions of *k-power domination* and *k-forcing*, and provide the initial connection between both concepts.

Observation 3.1. *In any graph G , if T is a k -forcing set, all sets $(\mathcal{F}_{G,k}^i(T))_{i \geq 0}$ are k -forcing sets of G ; if S is a k -power dominating set of G , the sets $(\mathcal{P}_{G,k}^i(S))_{i \geq 0}$ are also k -forcing sets of G .*

Observation 3.2. *In any graph G , if T is a k -forcing set of G then T is also a k -power dominating set. The converse is not necessarily true, but S is a k -power dominating set if and only if $N[S]$ is a k -forcing set. As a consequence, $\gamma_{p,k}(G) \leq Z_k(G) \leq \gamma_{p,k}(G)(\Delta(G)+1)$.*

Observation 3.3. *In a graph G , $S \subsetneq V(G)$ is a k -power dominating set of G if and only if $N[S] \setminus S$ is a k -forcing set of $G - S$.*

Note that given a graph $G = (V, E)$ and $S \subseteq X \subseteq V$, it is possible that for some $x \in X$, $\deg_{G[X]}(x) < \deg_G(x)$. Therefore, the *k-power domination* process starting with S in G is different from the one starting with S in $G[X]$. As a consequence, S being a *k-power dominating set* of $G[X]$ does not imply that S can *k-observe* all vertices in X when propagating in G . Analogously, if $T \subseteq X \subseteq V$ then T being a *k-forcing set* of $G[X]$ does not imply that T can *k-force* X in G . This observation motivates the following definitions.

Definition 3.4. Let $G = (V, E)$ be a graph and let $A \subseteq X \subseteq V$. We say that A is a *k-forcing set of X in G* if there exists a nonnegative integer t such that $X \subseteq \mathcal{F}_{G,k}^t(A)$.

Definition 3.5. Let $G = (V, E)$ be a graph and let $A \subseteq X \subseteq V$. We say that A is a *k-power dominating set of X in G* if there exists a nonnegative integer ℓ such that $X \subseteq \mathcal{P}_{G,k}^\ell(A)$.

The proofs of the next results are straightforward, and are omitted.

Lemma 3.6. *Let T be a k -forcing set of a graph G . Let $A \subseteq T$.*

- (1) *If A is k -forcing set of T in G , then A is a k -forcing set of G ;*
- (2) *If A is k -power dominating set of T in G , then A is a k -power dominating set of G .*

Lemma 3.7. *Let S be a k -power dominating set of a graph G . Let $A \subseteq S$.*

- (1) *If A is k -forcing set of $N[S]$ in G , then A is a k -forcing set of G ;*
- (2) *If A is k -power dominating set of $N[S]$ in G , then A is a k -power dominating set of G .*

Lemma 3.8. *Let $G = (V, E)$ be a graph and $X \subseteq V$ such that $G[X]$ is connected and $\deg_G(x) \leq k + 1$ for every $x \in X$. Let u be an arbitrary vertex in X . Then $\{u\}$ is a (minimum) k -power dominating set of $N_G[X]$ in G . In addition, if $\deg_G(u) \leq k$, then $\{u\}$ is also a (minimum) k -forcing set of $N_G[X]$ in G .*

Proof. If $S = \{u\}$, then $\mathcal{P}_{G,k}^0(S) = N_G[u]$ and $\mathcal{F}_{G,k}^0(S) = \{u\}$. Let $x \in X$. Since $\deg_G(x) \leq k + 1$, $x \neq u$ has at most k unobserved neighbors when x is observed. Thus, $\mathcal{P}_{G,k}^i(S) \supseteq N_G[\mathcal{P}_{G,k}^{i-1}(S) \cap X]$ for every integer $i \geq 1$. Since $G[X]$ is connected, there exists an integer $r \geq 1$ such that $X \subseteq \mathcal{P}_{G,k}^r(S)$. Once all vertices in X are observed, each of them can have at most k unobserved neighbors, so such a vertex can observe any unobserved neighbors. Thus, $N_G[X] \subseteq \mathcal{P}_{G,k}^{r+1}(S)$ and S is a *k-power dominating set* of $N_G[X]$ in G . Now suppose $\deg_G(u) \leq k$. Then $\mathcal{F}_{G,k}^1(S) = N_G[S]$ and the argument proceeds as before. \square

The following result follows immediately from Lemma 3.8, but is already known for *k-power domination* [6, Lemma 7]; a slightly weaker version for *k-forcing* is given in [3, Proposition 2.3].

Corollary 3.9. *Let G be a connected graph. If $\Delta(G) \leq k + 1$, then $\gamma_{p,k}(G) = 1$; if in addition $\delta(G) \leq k$, then $Z_k(G) = 1$.*

When $G[X]$ is not connected, we apply [Lemma 3.8](#) in each of its components and obtain the following result.

Corollary 3.10. *Let $G = (V, E)$ be a connected graph, $X \subseteq V$, and $u_j \in V(G[X]_j)$ for every $j = 1, \dots, c(G[X])$. Let $S = \{u_1, \dots, u_{c(G[X])}\}$. If $\deg_G(x) \leq k + 1$ for every $x \in X$, then S is a k -power dominating set of $N[X]$ in G ; if in addition $\deg_G(u_j) \leq k$ for every $j = 1, \dots, c(G[X])$, then S is a k -forcing set of $N[X]$ in G .*

Proof. By [Lemma 3.8](#), for every $j = 1, \dots, c(G[X])$, $\{u_j\}$ is a k -power dominating set of $G[X]_j$ in G . Thus, S is a k -power dominating set of $N[X]$ in G . The argument for k -forcing is analogous. \square

Lemma 3.11. *Let $G = (V_G, E_G)$ and $H = (V_H, E_H)$ be two graphs and let $A \subseteq V_G$ and $B \subseteq V_H$. Suppose that there is a graph isomorphism φ from $G - A$ to $H - B$ such that $\varphi(N_G[A] \setminus A) = N_H[B] \setminus B$. Then*

- (1) A is a k -power dominating set of G if and only if B is a k -power dominating set of H ;
- (2) $N_G[A]$ is a k -forcing set of G if and only if $N_H[B]$ is a k -forcing set of H .

Proof. (1) If A is a k -power dominating set of G , then $N_G[A] \setminus A$ is a k -forcing set of $G - A$ by [Observation 3.3](#). Since $\varphi(G - A) = H - B$ and $\varphi(N_G[A] \setminus A) = N_H[B] \setminus B$, we use φ to map the forcing process. That is, if $u \rightarrow v$ in $G - A$, then $\varphi(u) \rightarrow \varphi(v)$ in $H - B$, and obtain that $N_H[B] \setminus B$ is a k -forcing set of $H - B$. By [Observation 3.3](#), B is a k -power dominating set of H .

(2) If $N_G[A]$ is a k -forcing set of G , then A is a k -power dominating set of G by [Observation 3.2](#). Using (1) we conclude that B is a k -power dominating set of H , and by [Observation 3.2](#) we conclude that $N_H[B]$ is a k -forcing set of H . \square

Corollary 3.12. *Let $G = (V_G, E_G)$ and $H = (V_H, E_H)$ be two graphs and let $A \subseteq V_G$ and $B \subseteq V_H$. Suppose that there is a graph isomorphism φ from $G - A$ to $H - B$ such that $\varphi(N_G[A] \setminus A) = N_H[B] \setminus B$. Let $P \subseteq V_G \setminus A$. Then*

- (1) $A \cup P$ is a k -power dominating set of G if and only if $B \cup \varphi(P)$ is a k -power dominating set of H ;
- (2) $N_G[A] \cup P$ is a k -forcing set of G if and only if $N_G[B] \cup \varphi(P)$ is a k -forcing set of H .

Proof. Define $A' = A \cup P$ and $B' = B \cup \varphi(P)$. Then φ is an isomorphism from $G - A'$ to $H - B'$ and $\varphi(N_G[A'] \setminus A') = N_H[B'] \setminus B'$. So we can apply [Lemma 3.11](#) with G, H, A' and B' . \square

While all the previous results include analogous statements for k -forcing and a k -power domination, the following lemma does not have a k -forcing analog.

Lemma 3.13 [[6, Lemma 9](#)]. *If G is connected and $\Delta(G) \geq k + 2$, then there exists a minimum k -power dominating set S such that $\deg(v) \geq k + 2$ for all $v \in S$.*

To see that there is no k -forcing analog to [Lemma 3.13](#) it is sufficient to consider $K_{1,n}$, the complete bipartite graph with one vertex in one part and n vertices in the other. As shown in [[3](#)], if $n > k$ every minimum k -forcing set contains at least one vertex of degree 1.

4. Graph contraction

Definition 4.1. Let G be a graph and let $X \subseteq V(G)$. Define \widehat{X} to be the graph obtained from $G[X]$, by attaching to each vertex $x \in X$ exactly $|N_G(x) \setminus X|$ pendent vertices.

Lemma 4.2. *Let G be a connected graph and let $X \subseteq V(G)$. There exists $S \subseteq X$ such that S is a minimum k -power dominating set of \widehat{X} .*

Proof. Let $Y \subseteq X$ be a set of vertices that induces a component of $G[X]$. Suppose first that $\Delta(\widehat{Y}) \leq k + 1$. Then by [Lemma 3.8](#) any one vertex of Y is a k -power dominating set for $N_{\widehat{Y}}[Y] = V(\widehat{Y})$; a one vertex k -power dominating set is necessarily minimum. Now assume $\Delta(\widehat{Y}) \geq k + 2$. By definition of \widehat{Y} , $\deg_{\widehat{Y}}(u) = 1$ for every $u \in V(\widehat{Y}) \setminus Y$. Since $\Delta(\widehat{Y}) \geq k + 2$, by [Lemma 3.13](#) there exists a minimum k -power dominating set S of \widehat{Y} that contains only vertices in Y . When $G[X]$ has more than one component, we take the union across the components of the power dominating sets just found. \square

For the same reasons why there is no k -forcing analog to [Lemma 3.13](#), there is no k -forcing analog to [Lemma 4.2](#). Indeed, if $x \in V(G)$ and $\deg_G(x) \geq k + 1$, a minimum k -forcing set of $\{\widehat{x}\}$ must contain a vertex of degree 1.

Lemma 4.3. *Let G be a connected graph and let $X \subseteq V(G)$. If $S \subseteq X$ is a minimum k -power dominating set of \widehat{X} , then S is a k -power dominating set of $N_G[X]$ in G .*

Proof. Each vertex in $V(\widehat{X}) \setminus X$ arises from a vertex $y \notin X$ that is a neighbor of a vertex $x \in X$. For every $x \in X$, let N_x denote the (possibly empty) set of neighbors of x in $V(\widehat{X}) \setminus X$ i.e., $N_x = N_{\widehat{X}}(x) \setminus X$ and let $N'_x = N_G(x) \setminus X$. Since $S \subseteq X$ and $\deg_{\widehat{X}}(u) = 1$




Fig. 1. The graphs U_2, \widehat{X} , and U_2/X defined in Theorem 4.4 are shown. In each case, a minimum 2-power dominating set is indicated by coloring. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

for every $u \in V(\widehat{X}) \setminus X$, none of the vertices in N_x can be observed before x is observed, and moreover, all vertices in N_x are observed simultaneously. Since for every $x \in X$, $\deg_{\widehat{X}}(x) = \deg_G(x)$, the only difference between the k -power domination process starting with S in \widehat{X} and the one starting with S in G is that when the vertices in N_x are observed in \widehat{X} , the unobserved vertices in N'_x become observed in G . The reason why some vertices in N'_x could have been observed earlier is that a vertex in $G - X$ could have more than one neighbor in X so $(N'_x)_{x \in X}$ are not necessarily disjoint. Since for every $w \in N_G[X] \setminus X$ there exists $x \in X$ such that $w \in N'_x$, all vertices in $N_G[X]$ are observed. \square

Theorem 4.4. Let $G = (V, E)$ be a connected graph. If $X \subseteq V$,

$$\gamma_{p,k}(G/X) - 1 \leq \gamma_{p,k}(G) \leq \gamma_{p,k}(G/X) + \gamma_{p,k}(\widehat{X})$$

and both bounds are tight.

Proof. Let $H = G/X$. By Lemma 4.2 there exists $T \subseteq X$ such that T is a minimum k -power dominating set of \widehat{X} and by Lemma 4.3, T is also a k -power dominating set of $N_G[X]$ in G .

To prove the upper bound we show that if P is a k -power dominating set of H , then $(P \setminus \{v_X\}) \cup T$ is a k -power dominating set of G .¹ Since T is a k -power dominating set of $N_G[X]$ in G , clearly $(P \setminus \{v_X\}) \cup T$ is a k -power dominating set of $N_G[P \setminus \{v_X\}] \cup N_G[X] = N_G[P \setminus \{v_X\} \cup X]$ in G . We will prove that $(P \setminus \{v_X\}) \cup X$ is a k -power dominating set of G , which by Lemma 3.7 suffices to conclude that $(P \setminus \{v_X\}) \cup T$ is a k -power dominating set of G . Let $A = X$ and $B = \{v_X\}$. Since $H = G/X$, $G - A = H - B$ and $N_G[A] \setminus A = N_H[B] \setminus B$, we apply Corollary 3.12 with φ being the identity function and conclude that $(P \setminus \{v_X\}) \cup A$ is a k -power dominating set of G if and only if $(P \setminus \{v_X\}) \cup B$ is a k -power dominating set of H . Since $B = \{v_X\}$, $(P \setminus \{v_X\}) \cup B = P$ and P is a k -power dominating set of H , $(P \setminus \{v_X\}) \cup A$ is a k -power dominating set of G .

To prove the lower bound, we show that if S is a minimum k -power dominating set of G , then $(S \setminus X) \cup \{v_X\}$ is a k -power dominating set of H . As above, let $A = X$ and $B = \{v_X\}$ so $G - A = H - B$ and $N_G[A] \setminus A = N_H[B] \setminus B$. Then we apply Corollary 3.12 to conclude that $(S \setminus X) \cup A$ is a k -power dominating set of G if and only if $(S \setminus X) \cup B$ is a k -power dominating set of H . Since $X = A$, then $(S \setminus X) \cup A = S$ and it is a k -power dominating set of G . Then $(S \setminus X) \cup B = (S \setminus X) \cup \{v_X\}$ is a k -power dominating set of H . Thus, $\gamma_{p,k}(G/X) \leq |(S \setminus X) \cup \{v_X\}| \leq |S| + 1 = \gamma_{p,k}(G) + 1$.

To prove the upper bound is tight, for each integer $q \geq k$ we define a graph U_q and a set $X \subseteq V(U_q)$ such that $\gamma_{p,k}(U_q) = \gamma_{p,k}(U_q/X) + \gamma_{p,k}(\widehat{X})$ (see Fig. 1). Consider two disjoint copies of K_{q+2} , say G and G' , and vertices $x \in V(G)$ and $y \in V(G')$. Construct U_q by adding the edge $e = \{x, y\}$ and define $X = V(G') \setminus \{y\}$. Then $\gamma_{p,k}(U_q) = 2$, $\gamma_{p,k}(\widehat{X}) = 1$, and $\gamma_{p,k}(U_q/X) = 1$.

To show the lower bound is tight, for each integer $q \geq k$ we define a graph L_q and a set $X \subseteq V(L_q)$ such that $\gamma_{p,k}(L_q/X) - 1 = \gamma_{p,k}(L_q)$ (see Fig. 2). Assume first that $k \geq 2$. Construct L_q starting with a cycle of length $2q$ with vertices v_1, \dots, v_{2q} . Attach a pendent vertex to each vertex v_i , for $i = 1, \dots, 2q$. Then attach $q + 1$ pendent vertices to the pendent neighbor of v_1 , so $\gamma_{p,k}(L_q) = 1$. For $X = \{v_1, \dots, v_{2p}\}$, $\gamma_{p,k}(L_q/X) = 2$. Now suppose $k = 1$, and begin with a path of length 6 with vertices v_0, \dots, v_6 . Construct L_q by attaching q pendent vertices to v_0 . If $X = \{v_1, v_3, v_5\}$, then $\gamma_{p,1}(L_q) = 1$ and $\gamma_{p,1}(L_q/X) = 2$. \square

The next example shows that it is possible to find a graph G and a subgraph X for which the gap between $\gamma_{p,k}(G)$ and $\gamma_{p,k}(G/X)$ is arbitrarily large.

Example 4.5. Given a positive integer c , define $T_{k,c}$ as the tree obtained by adding $k + 2$ leaves to each leaf of $K_{1,c}$. If X is the set of all vertices of degree greater than one in $T_{k,c}$, then $\gamma_{p,k}(T_{k,c}) = \gamma_{p,k}(\widehat{X}) = c$ and $\gamma_{p,k}(T_{k,c}/X) = 1$.

Corollary 4.6. Let $G = (V, E)$ be a connected graph. Let $X \subseteq V$ such that $\deg_G(x) \leq k + 1$ for every $x \in X$. Then

$$\gamma_{p,k}(G/X) - 1 \leq \gamma_{p,k}(G) \leq \gamma_{p,k}(G/X) + c(G[X]).$$

¹ Note that whether $v_X \notin P$ or $v_X \in P$ does not affect the conclusion, since in any case $|(P \setminus \{v_X\}) \cup T| \leq |P| + |T| = \gamma_{p,k}(H) + \gamma_{p,k}(\widehat{X})$; we only exclude v_X from P to guarantee $(P \setminus \{v_X\}) \cup T \subseteq V(G)$.




Fig. 2. The graphs L_2 and L_2/X defined in [Theorem 4.4](#) are shown. In each case, a minimum 2-power dominating set is indicated by coloring. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Proof. It is sufficient to show that $\gamma_{P,k}(\widehat{X}) \leq c(G[X])$. Observe that $\deg_G(x) \leq k + 1$ for every $x \in X$ implies that $\deg_{\widehat{X}}(x) \leq k + 1$ for every $x \in V(\widehat{X})$. By [Corollary 3.10](#) there exists a k -power dominating set of $N_{\widehat{X}}[X] = V(\widehat{X})$ in G with cardinality $c(\widehat{X}) = c(G[X])$, so $\gamma_{P,k}(\widehat{X}) \leq c(G[X])$. \square

Corollary 4.7. Let $G = (V, E)$ be a connected graph. Let $X \subseteq V$ such that $G[X]$ is connected and $\deg_G(x) \leq k + 1$ for every $x \in X$. Then

$$\gamma_{P,k}(G/X) - 1 \leq \gamma_{P,k}(G) \leq \gamma_{P,k}(G/X) + 1.$$

Proposition 4.8. Let $G = (V, E)$ be a connected graph. Let $X \subseteq V$ such that $G[X]$ is connected and $\deg_G(x) \leq 2$ for every $x \in X$. Then $\gamma_{P,1}(G/X) \leq \gamma_{P,1}(G)$ and this bound is tight.

Proof. Let $H = G/X$. Since $\Delta(G) \leq 2$ implies that G itself is a path or a cycle, without loss of generality we can assume $\Delta(G) \geq 3$. By [Lemma 3.13](#), there exists a minimum 1-power dominating set S of G such that $\deg_G(u) \geq 3$ for every $u \in S$, so $S \subseteq V \setminus X$. We define H to be G/X and prove that S is also a 1-power dominating set of H .

As in the proof of [Theorem 4.4](#), $S \cup \{v_X\}$ is a 1-power dominating set of H . Then by [Observation 3.2](#), $N_H[S \cup \{v_X\}] = N_H[S] \cup N_H[v_X]$ is a 1-forcing set of H .

Note that $S \subseteq V \setminus X$ implies $\mathcal{P}_{G,1}^0(S) \setminus X = \mathcal{P}_{H,1}^0(S) \setminus \{v_X\}$ and as long as $\mathcal{P}_{G,1}^i(S) \subseteq V \setminus X$, $\mathcal{P}_{G,1}^i(S) = \mathcal{P}_{H,1}^i(S)$. Let x be a vertex of X that is observed first (meaning that no vertex of X has been observed earlier), and let y be the vertex in $G - X$ that dominates or forces x at time t ($x \in \mathcal{P}_{G,1}^0(S)$ for $t = 0$ or $x \in \mathcal{P}_{G,1}^t(S) \setminus \mathcal{P}_{G,1}^{t-1}(S)$ for $t \geq 1$). Since $\deg_G(y) \geq \deg_H(y)$, y can also dominate or force in H . Thus $v_X \in \mathcal{P}_{H,1}^t(S)$. Since $\deg_H(v_X) \leq 2$, it takes at most one additional application of 1-forcing to observe all vertices in $N_H[v_X]$, so $N_H[v_X] \subseteq \mathcal{P}_{H,1}^{t+1}(S)$.

Since $N_H[S] = \mathcal{P}_{H,1}^0(S) \subseteq \mathcal{P}_{H,1}^{t+1}(S)$ and $N_H[v_X] \subseteq \mathcal{P}_{H,1}^{t+1}(S)$, then $N_H[S \cup \{v_X\}] = N_H[S] \cup N_H[v_X] \subseteq \mathcal{P}_{H,1}^{t+1}(S)$. Moreover, since $N_H[S \cup \{v_X\}]$ is a 1-forcing set of H , so is $\mathcal{P}_{H,1}^{t+1}(S)$ and therefore, S is a 1-power dominating set of H .

To prove the tightness, observe that for $n \geq 3$, contracting the set of all vertices of degree 2 in the path P_n of order n produces the path P_3 . Now, $\gamma_{P,1}(P_n) = \gamma_{P,1}(P_3) = 1$. \square

Due to the computational complexity of the k -power domination problem, efficient algorithms to approximate optimal k -power dominating sets are of practical importance. [Theorem 4.4](#) could help in the parallel search for k -power dominating sets. The following result provides a theoretical framework to study practical uses of graph decomposition as a tool for the parallel computation of k -power dominating sets.

Theorem 4.9. Let $G = (V, E)$ be a connected graph and let P_1, \dots, P_r be a partition of V . Then

$$\gamma_{P,k}(G) \leq \sum_{i=1}^r \gamma_{P,k}(\widehat{P}_i).$$

Proof. By [Lemma 4.2](#), for every $i = 1, \dots, r$ there exists $S_i \subseteq P_i$ such that S_i is a minimum k -power dominating set of \widehat{P}_i . By [Lemma 4.3](#), S_i is also a k -power dominating set of $N_G[P_i]$ in G , and as a consequence, $S = \cup_{i=1}^r S_i$ is a k -power dominating set of G . Then $\gamma_{P,k}(G) \leq |S| = \sum_{i=1}^r |S_i| = \sum_{i=1}^r \gamma_{P,k}(\widehat{P}_i)$. \square

To prove that the bound in [Theorem 4.9](#) is tight we will use the family of Sierpiński graphs whose definition we recall, using the notation in [\[8\]](#). Given two positive integers n and p the Sierpiński graph S_p^n has as vertices all n -tuples of integers in $\{0, 1, \dots, p - 1\}$ denoted as $s_n s_{n-1} \dots s_2 s_1$. Two vertices $s_n \dots s_1$ and $t_n \dots t_1$ are adjacent in S_p^n if and only if there exists an r with $1 \leq r \leq n$ such that

- (i) $s_i = t_i$ for every $i \in \{r + 1, \dots, n\}$,
- (ii) $s_r \neq t_r$, and
- (iii) $s_i = t_r$ and $t_i = s_r$ for every $i \in \{1, \dots, r - 1\}$.

The definition of Sierpiński graphs implies that $S_p^1 = K_p$ and if $n \geq 2$, S_p^n has p^{n-i} induced copies of S_p^i . Moreover, the vertices in each of those copies coincide in the $n-i$ leftmost digits $s_n \cdots s_{i+1}$ [8]. If s is a $(n-i)$ -tuple of integers in $\{0, \dots, p-1\}$, let sS_p^i denote the set of vertices of S_p^n whose leftmost $n-i$ digits coincide with s . For simplicity, we use $S_p^n[s]$ to denote $S_p^n[sS_p^i]$ (the subgraph induced by sS_p^i in S_p^n), so $S_p^n[s]$ is isomorphic to S_p^i .

Lemma 4.10. *Given integers $n \geq 4$, $k \geq 1$ and $p \geq k + 2$, let s be a $(n - 3)$ -tuple of integers in $\{0, \dots, p - 1\}$. Then $\gamma_{p,k}(\widehat{sS_p^3}) = \gamma_{p,k}(S_p^3)$.*

Proof. Fix $k \geq 1$, $p \geq k + 2$, and $n \geq 4$. We begin by determining the degree of a vertex $sxyz$ in S_p^n and in $S_p^n[s] \cong S_p^3$. The definition of the Sierpiński graph S_p^n implies that vertices of the form a^n have degree $p - 1$ and all the other vertices have degree p in S_p^n . If xyz is nonconstant, then vertex $sxyz$ has degree p in both S_p^n and $S_p^n[s]$, so $sxyz$ does not have any pendent vertices added in $\widehat{sS_p^3}$. Now consider the constant sequence aaa . If $s \neq a^{n-3}$, then vertex $saaa$ has degree p in S_p^n but degree $p - 1$ in $S_p^n[s]$, so one leaf is added to $saaa$ in $\widehat{sS_p^3}$. If $s = a^{n-3}$, then vertex $saaa = a^n$ has degree $p - 1$ in both S_p^n and $S_p^n[s]$, so $saaa$ is unchanged in $\widehat{sS_p^3}$.

If $s \neq a^{n-3}$ for any $a \in \{0, \dots, p - 1\}$, then $\widehat{sS_p^3}$ is obtained from $S_p^n[s]$ by attaching one pendent vertex to each of the p vertices of the form $saaa$ for $a \in \{0, \dots, p - 1\}$, so that every vertex of $\widehat{sS_p^3}$ has degree p ; denote this graph by G_1 . If $s = a^{n-3}$ for some $a \in \{0, \dots, p - 1\}$, then $\widehat{sS_p^3}$ is obtained from $S_p^n[s]$ by attaching one pendent vertex to each of the $p - 1$ vertices of the form $sbbb$ for $b \neq a$, so that every vertex of $\widehat{sS_p^3}$ except a^n has degree p ; denote this graph by G_2 . Observe that the only difference between G_1 and G_2 is that G_2 is missing one leaf.

To show that $\gamma_{p,k}(G_i) = \gamma_{p,k}(S_p^3)$ for $i = 1, 2$, we first we prove $\gamma_{p,k}(G_i) \leq \gamma_{p,k}(S_p^3)$ by showing that if P is a k -power dominating set of S_p^3 , then P is also a k -power dominating set of G_i . For each pendent vertex x in G_i , let u_x denote its only neighbor (in G_i). Then u_x is a vertex of degree $p - 1$ in S_p^3 and therefore, in S_p^3 it is labeled as a^3 for some $a \in \{0, \dots, p - 1\}$. If $u_x \in P$, then $x \in N_{G_i}[P] = \mathcal{P}_{G_i,k}^0(P)$. If $u_x \notin P$, then $u_x \in \mathcal{P}_{S_p^3,k}^t(P)$ for some integer $t > 0$. Note that in S_p^3 , u_x cannot be observed until one of its neighbors is. Since $u_x = a^3$, its $p - 1$ neighbors in S_p^3 have labels in the form aab for $b = 0, \dots, p - 1$, $b \neq a$. Therefore, $N_{S_p^3}[u_x]$ induces a p -clique in S_p^3 . This means that when a neighbor forces u_x , it also forces all the vertices in the p -clique induced by $N_{S_p^3}[u_x]$. When this happens in G_i instead of in S_p^3 , u_x has exactly one unobserved neighbor (x), so $x \in \mathcal{P}_{S_p^3,k}^{t+1}(P)$.

Finally we prove $\gamma_{p,k}(G_i) \geq \gamma_{p,k}(S_p^3)$ by showing that there exists a minimum k -power dominating set Q of G_i that is also a k -power dominating set of S_p^3 . By Lemma 3.13 there exists a minimum k -power dominating set Q of G_i that does not contain vertices of degree 1, so $Q \subseteq V(S_p^3)$. Therefore, in the k -power domination process starting with Q in G_i , a vertex of degree 1 in G_i cannot be observed until its one neighbor in S_p^3 is. Then Q is a k -power dominating set of S_p^3 . We conclude that $\gamma_{p,k}(\widehat{sS_p^3}) = \gamma_{p,k}(S_p^3)$. □

It is known that if $n \geq 3$, $k \geq 1$, and $p \geq k + 2$, then $\gamma_{p,k}(S_p^n) = p^{n-2}(p - k - 1)$ [8] so we immediately obtain the following result.

Corollary 4.11. *Given integers $n \geq 4$, $k \geq 1$, and $p \geq k + 2$, let s be a $(n - 3)$ -tuple of integers in $\{0, \dots, p - 1\}$. Then $\gamma_{p,k}(\widehat{sS_p^3}) = p(p - k - 1)$.*

Lemma 4.12. *Given integers $n \geq 4$, $k \geq 1$, and $p \geq k + 2$, let T denote the set of all $(n - 3)$ -tuples of integers in $\{0, \dots, p - 1\}$. Then $\gamma_{p,k}(S_p^n) = \sum_{t \in T} \gamma_{p,k}(tS_p^3)$.*

Proof. By Corollary 4.11 $\gamma_{p,k}(tS_p^3) = \gamma_{p,k}(S_p^3) = p(p - k - 1)$ for every $t \in T$. There are p^{n-3} tuples in T , so $\sum_{t \in T} \gamma_{p,k}(tS_p^3) = p^{n-3}p(p - k - 1) = p^{n-2}(p - k - 1) = \gamma_{p,k}(S_p^n) = p^{n-2}(p - k - 1)$. □

The bound in Theorem 4.9 is $\gamma_{p,k}(S_p^n) \leq \sum_{t \in T} \gamma_{p,k}(tS_p^3)$, so the following result is an immediate consequence of Lemma 4.12.

Corollary 4.13. *The bound in Theorem 4.9 is tight.*

Next we present the equivalent results for k -forcing, taking into consideration the following differences between k -power domination and k -forcing.

1. For the lower bound, note that if $\deg_H(v_X) > k$, $\{v_X\}$ does not force $N_H(v_X) = N_G[X] \setminus X$. In that case, to obtain a k -forcing set of H from a k -forcing set of G it might be necessary to add at most $|N_G[X] \setminus X| - k$ vertices.
2. For the upper bound, since there is no k -forcing equivalent to Lemma 4.2, it could happen that every minimum k -forcing set of \widehat{X} contains a vertex $x \in N[X] \setminus X$ for which $\deg_{\widehat{X}}(x) = 1$ but $\deg_G(x) > k$. Thus, x forces its neighbors in \widehat{X} but not in G , and a k -forcing set of \widehat{X} might not force X in G .

The proofs are omitted because each proof is analogous to the proof of the corresponding result for power domination: [Lemma 4.14](#) corresponds to [Lemma 4.3](#), [Proposition 4.15](#) to [Theorem 4.4](#), [Proposition 4.16](#) to [Corollary 4.6](#), and [Theorem 4.17](#) to [Theorem 4.9](#).

Lemma 4.14. *Let G be a connected graph and let $X \subseteq V(G)$. If there exists $S \subseteq X$ minimum k -forcing set of \widehat{X} , then S is a k -forcing of $N_G[X]$ in G .*

Proposition 4.15. *Let $G = (V, E)$ be a connected graph. Let $X \subseteq V(G)$. If there exists a minimum k -forcing set of \widehat{X} that contains only vertices in X , then*

$$Z_k(G/X) + Z_k(\widehat{X}) \geq Z_k(G) \geq \begin{cases} Z_k(G/X) - 1 & \text{if } |N[X] \setminus X| \leq k, \\ Z_k(G/X) - |N[X] \setminus X| + k & \text{if } |N[X] \setminus X| > k. \end{cases}$$

Proposition 4.16. *Let $G = (V, E)$ be a connected graph. Let $X \subseteq V$ such that $\deg_G(x) \leq k$ for $x \in X$. If there exists a minimum k -forcing set of \widehat{X} that contains only vertices in X , then*

$$Z_k(G/X) + c(G[X]) \geq Z_k(G) \geq \begin{cases} Z_k(G/X) - 1 & \text{if } |N[X] \setminus X| \leq k, \\ Z_k(G/X) - |N[X] \setminus X| + k & \text{if } |N[X] \setminus X| > k. \end{cases}$$

Theorem 4.17. *Let $G = (V, E)$ be a connected graph and let P_1, \dots, P_r be a partition of V . If \widehat{P}_i has a minimum k -forcing set in P_i for every $i = 1, \dots, r$, then*

$$Z_k(G) \leq \sum_{i=1}^r Z_k(\widehat{P}_i).$$

[Theorems 4.9](#) and [4.17](#) provide upper bounds for the k -power domination and the k -forcing number of a graph in terms of the k -power domination and the k -forcing number of P_1, \dots, P_r , which can be computed in parallel. In particular, the importance of [Theorems 4.9](#) and [4.17](#) resides in the fact that \widehat{P}_i might have properties that do not hold for G . For example, suppose G is not a tree, but there is a linear algorithm to partition $V(G)$ into sets P_1, \dots, P_r such that $\widehat{P}_1, \dots, \widehat{P}_r$ are trees. Then using the linear algorithm for trees provided in [\[7\]](#), $\gamma_{p,k}(\widehat{X})$ can be computed in linear time. The exploration of possible uses of our results in algorithms to find k -power dominating or k -forcing sets a graph requires a detailed and careful analysis that is outside the scope of this paper.

5. k -power domination and k -forcing numbers

By [Observation 3.2](#), $\gamma_{p,k}(G) \leq Z_k(G) \leq \gamma_{p,k}(G)(\Delta(G) + 1)$. In this section we improve the upper bound in the previous inequality by generalizing a result by Benson et al. [\[4, Theorem 3.2\]](#) for 1-power domination. An important concept in this work is that of *private neighborhood*, which we recall. Suppose $v \in S \subseteq V$. A S -private neighbor of v is a vertex $x \in N(v)$ such that $x \notin N(S \setminus \{v\})$. Moreover, we say that x is an *external S -private neighbor* if $x \notin S$.

Lemma 5.1 ([\[6, Lemma 10\]](#)). *In every connected graph G with $\Delta(G) \geq k + 2$ there exists a minimum k -power dominating set S in which every vertex has at least $k + 1$ S -private neighbors.*

We strengthen [Lemma 5.1](#) by extending it to external private neighbors.

Lemma 5.2. *In every connected graph G with $\Delta(G) \geq k + 2$ there exists a minimum k -power dominating set S in which every vertex has at least $k + 1$ external S -private neighbors.*

Proof. By [Lemma 5.1](#) there exists a minimum k -power dominating set S in which every vertex has at least $k + 1$ S -private neighbors. Suppose that there exists $u \in S$ that has at most k external private neighbors. We prove that $S' = S \setminus \{u\}$ is k -power dominating set, which contradicts the minimality of S . Since u has at least $k + 1$ neighbors and at most k of them are outside S , there exists $y \in S$ such that u and y are neighbors. This implies that $u \in \mathcal{P}_{G,k}^0(S')$. Moreover, all non-external neighbors of u are in S and thus in $\mathcal{P}_{G,k}^0(S')$ so u has at most k unobserved neighbors. Thus, $N_G[u] \subseteq \mathcal{P}_{G,k}^1(S')$, so S' is a k -power dominating set. \square

Lemma 5.3. *If G is a connected graph with $\Delta(G) \geq k + 2$ and $S = \{u_1, \dots, u_t\}$ is a minimum k -power dominating set of G in which every vertex has at least $k + 1$ external S -private neighbors, then*

$$Z_k(G) \leq \sum_{i=1}^t (\deg u_i + 1 - k).$$

Proof. By hypothesis, for each $i = 1, \dots, t$ there exists a set $\{x_1^{(i)}, \dots, x_k^{(i)}\}$ of external S -private neighbors of u_i . We prove that $B = \bigcup_{i=1}^t (N[u_i] \setminus \{x_1^{(i)}, \dots, x_k^{(i)}\})$ is a k -forcing set of G . Since $x_1^{(i)}, \dots, x_k^{(i)}$ are external S -private neighbors of u_i , then

$\{x_1^{(i)}, \dots, x_k^{(i)}\} \cap S = \emptyset$, which implies $u_i \in B$, for every $i = 1, \dots, t$. In the first step of the k -forcing process each vertex u_i forces $x_1^{(i)}, \dots, x_k^{(i)}$ so B is a k -forcing set of $N[S]$ in G . Since S is a k -power dominating set of G , by **Observation 3.2** $N[S]$ is a k -forcing set of G . Then by **Lemma 3.6**, B is a k -forcing set of G so, $Z_k(G) \leq |B| \leq \sum_{i=1}^t |N[u_i] \setminus \{x_1^{(i)}, \dots, x_k^{(i)}\}| \leq \sum_{i=1}^t (\deg u_i + 1 - k)$. \square

Theorem 5.4. *In every connected graph G with $\Delta(G) \geq k + 2$,*

$$Z_k(G) \leq \gamma_{p,k}(G)(\Delta(G) + 1 - k), \text{ or equivalently, } \left\lceil \frac{Z_k(G)}{\Delta(G) + 1 - k} \right\rceil \leq \gamma_{p,k}(G)$$

and this lower bound for $\gamma_{p,k}(G)$ is tight.

Proof. By **Lemma 5.2** there exists a minimum k -power dominating set $S = \{u_1, \dots, u_{\gamma_{p,k}(G)}\}$ of G in which each vertex has at least $k + 1$ external S -private neighbors. By **Lemma 5.3**, $Z_k(G) \leq \sum_{i=1}^{\gamma_{p,k}(G)} (\deg u_i + 1 - k) \leq \gamma_{p,k}(G)(\Delta(G) + 1 - k)$, and as a consequence, $\left\lceil \frac{Z_k(G)}{\Delta(G) + 1 - k} \right\rceil \leq \gamma_{p,k}(G)$.

To prove that the bound is tight, let $r \geq 2$ and $p > 3r + k - 3$. Construct the graph $G_{p,r}$ by adding p pendent vertices to each vertex of a path of order r . Then $\Delta(G_{p,r}) = p + 2$ and since $p \geq k + 1$, $\gamma_{p,k}(G_{p,r}) = r$, $Z_k(G_{p,r}) = r(p - k)$, and $\left\lceil \frac{r(p-k)}{p+3-k} \right\rceil = r$. \square

Next we apply **Theorem 5.4** to obtain lower bounds for the k -forcing number of graphs from upper bounds for the k -power domination number of an arbitrary graph presented in [6] and improved in [7] for $(k + 2)$ -regular graphs.

Theorem 5.5 ([6, Theorem 11]). *Let G be a connected graph with $|G| \geq k + 2$. Then $\gamma_{p,k}(G) \leq \frac{|G|}{k+2}$.*

Corollary 5.6. *In a connected graph G with $\Delta(G) \geq k + 2$,*

$$Z_k(G) \leq \left\lfloor \frac{|G|}{k+2} (\Delta(G) + 1 - k) \right\rfloor$$

and this bound is tight.

Proof. Since $\Delta(G) \geq k + 2$ implies $|G| \geq k + 2$ we apply **Theorem 5.5** and obtain $\gamma_{p,k}(G) \leq \frac{|G|}{k+2}$. By **Theorem 5.4** we know $Z_k(G) \leq \gamma_{p,k}(G)(\Delta(G) + 1 - k)$ and combining both inequalities we conclude $Z_k(G) \leq \left\lfloor \frac{|G|}{k+2} (\Delta(G) + 1 - k) \right\rfloor$.

To show this bound is tight, observe that $Z_k(K_{k+3}) = 3$, and the upper bound in this case is $\left\lfloor \frac{k+3}{k+2} (k + 2 + 1 - k) \right\rfloor = 3$ for $k \geq 2$. \square

Theorem 5.7 ([6, Theorem 2.1]). *Let G be a connected $(k + 2)$ -regular graph. If $G \neq K_{k+2,k+2}$, then $\gamma_{p,k}(G) \leq \frac{|G|}{k+3}$.*

Corollary 5.8. *Let G be a connected $(k + 2)$ -regular graph. If $G \neq K_{k+2,k+2}$, then $Z_k(G) \leq \frac{3|G|}{k+3}$.*

Proof. Since G is $(k + 2)$ -regular, $\Delta(G) = k + 2$ so we apply **Theorem 5.4** and obtain $Z_k(G) \leq \frac{|G|}{k+3} (k + 2 + 1 - k) = \frac{3|G|}{k+3}$. To see that the bound is best possible it suffices to consider K_{k+3} which is $(k + 2)$ -regular and $Z_k(K_{k+3}) = 3 = \frac{3(k+3)}{k+3}$. \square

Acknowledgments

We thank the referees for their careful reading of the paper and many helpful comments, which have improved the paper.

References

- [1] A. Aazami, Hardness Results and Approximation Algorithms for Some Problems on Graphs, University of Waterloo, 2008. Available at <https://uwaterloo.ca/handle/10012/4147?show=full>.
- [2] AIM Minimum Rank–Special Graphs Work Group, F. Barioli, W. Barrett, S. Butler, S.M. Cioabă, D. Cvetković, S.M. Fallat, C. Godsil, W. Haemers, L. Hogben, R. Mikkelsen, S. Narayan, O. Pryporova, I. Sciriha, W. So, D. Stevanović, H. van der Holst, K. Vander Meulen, A. Wangsness, Zero forcing sets and the minimum rank of graphs, *Linear Algebra Appl.* 428 (2008) 1628–1648.
- [3] D. Amos, Y. Caro, R. Davila, R. Pepper, Upper bounds on the k -forcing number of a graph, *Discrete Appl. Math.* 181 (2015) 1–10.
- [4] K.F. Benson, D. Ferrero, M. Flagg, V. Furst, L. Hogben, V. Vasilevska, B. Wissman, Power domination and zero forcing. Available at <http://arxiv.org/abs/1510.02421>.
- [5] D. Burgarth, V. Giovannetti, Full control by locally induced relaxation, *Phys. Rev. Lett.* 99 (2007) 100501.
- [6] G.J. Chang, P. Dorbec, M. Montassier, A. Raspaud, Generalized power domination in graphs, *Discrete Appl. Math.* 160 (2012) 1691–1698.
- [7] P. Dorbec, M.A. Henning, C. Lowenstein, M. Montassier, A. Raspaud, Generalized power domination in regular graphs, *SIAM J. Discrete Math.* 27 (2013) 1559–1574.
- [8] P. Dorbec, S. Klavžar, Generalized power domination in graphs: propagation radius and Sierpiński graphs, *Acta Appl. Math.* 134 (2014) 75–86.
- [9] T.W. Haynes, S.M. Hedetniemi, S.T. Hedetniemi, M.A. Henning, Domination in graphs applied to electric power networks, *SIAM J. Discrete Math.* 15 (2002) 519–529.