

12-2018

Pentagons in triangle-free graphs

Bernard Lidicky

Iowa State University, lidicky@iastate.edu

Florian Pfender

University of Colorado, Denver

Follow this and additional works at: https://lib.dr.iastate.edu/math_pubs



Part of the [Discrete Mathematics and Combinatorics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/math_pubs/175. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Mathematics at Iowa State University Digital Repository. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Pentagons in triangle-free graphs

Abstract

For all $n \geq 9$, we show that the only triangle-free graphs on n vertices maximizing the number 5-cycles are balanced blow-ups of a 5-cycle. This completely resolves a conjecture by Erdős, and extends results by Grzesik and Hatami, Hladký, Kral, Norin and Razborov, where they independently showed this same result for large n and for all n divisible by 5.

Disciplines

Discrete Mathematics and Combinatorics | Mathematics

Comments

This is a manuscript of an article published as Lidický, Bernard, and Florian Pfender. "Pentagons in triangle-free graphs." *European Journal of Combinatorics* 74 (2018): 85-89. doi: [10.1016/j.ejc.2018.07.008](https://doi.org/10.1016/j.ejc.2018.07.008). Posted with permission.

Pentagons in triangle-free graphs

Bernard Lidický* Florian Pfender†

December 27, 2017

Abstract

For all $n \geq 9$, we show that the only triangle-free graphs on n vertices maximizing the number 5-cycles are balanced blow-ups of a 5-cycle. This completely resolves a conjecture by Erdős, and extends results by Grzesik and Hatami, Hladký, Král', Norin, and Razborov, where they independently showed this same result for large n and for all n divisible by 5.

1 Introduction

In 1984, Erdős [1] conjectured that for all $n \geq 5$, the balanced blow-up of a 5-cycle maximizes the number of 5-cycles in the class of triangle-free graphs on n vertices. *Balanced blow-up* means in this context that every vertex of the 5-cycle is replaced by an independent set of size $\lfloor \frac{n}{5} \rfloor$ or $\lceil \frac{n}{5} \rceil$, and edges are replaced by complete bipartite graphs between the sets. This conjecture, in a way, supports the Meta-Theorem saying that among all triangle-free graphs, this blow-up of C_5 is the "least" bipartite graph.

In 2012, Grzesik [2], and independently in 2013, Hatami, Hladký, Král', Norin, and Razborov [3] settled this conjecture asymptotically by showing that any n -vertex triangle-free graph has at most $\frac{5!}{5^5} \binom{n}{5} (1 + o(1)) = 0.0384 \binom{n}{5} (1 + o(1))$ 5-cycles. Furthermore, the second group of authors showed uniqueness of the extremal graphs for most values of n .

Theorem 1 (Hatami et. al. [3]). *Let n be either divisible by 5, or large enough. The maximum number of copies of a 5-cycle in triangle-free graphs on n vertices is*

$$\prod_{i=0}^4 \left\lfloor \frac{n+i}{5} \right\rfloor.$$

Moreover, the only triangle-free graphs on n vertices maximizing the number 5-cycles are balanced blow-ups of a 5-cycle.

*Department of Mathematics, Iowa State University, Ames, IA, E-mail: lidicky@iastate.edu. Research of this author is supported in part by NSF grant DMS-1600390.

†Department of Mathematical and Statistical Sciences, University of Colorado Denver, E-mail: Florian.Pfender@ucdenver.edu. Research is partially supported in part by NSF grant DMS-1600483.

Both papers [2] and [3] use the theory of flag algebras, a theory developed by Razbarov [7], which can find inequalities of subgraph densities in graph limits with the help of semi-definite programming.

In 2011, Michael [6] observed that the Möbius ladder on 8 vertices (i.e. the 8-cycle with all diagonals added) contains the same number of 5-cycles as the balanced blow-up of C_5 . In this note we resolve the remaining cases of the conjecture. We also employ flag algebra techniques to find helpful inequalities, and then we use stability results. Finally, enumeration is used to deal with $n \leq 9$. Note that our analysis to show uniqueness is significantly simpler than the analysis in [3].

Theorem 2. *For all n , the maximum number of copies of a 5-cycle in any triangle-free graph on n vertices is*

$$\prod_{i=0}^4 \left\lfloor \frac{n+i}{5} \right\rfloor.$$

Moreover, the only triangle-free graphs on $n \geq 5$ vertices maximizing the number 5-cycles are balanced blow-ups of a 5-cycle, and the Möbius ladder ML_8 for the special case of $n = 8$.

For the ease of presentation, we will use a simplified language of graph limits. No understanding past the following definitions is required to follow this note. A *graphon* B of a graph G with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$ is a symmetric function $b : [0, 1]^2 \rightarrow \{0, 1\}$ with

$$b(x, y) = \begin{cases} 1, & \text{if } \frac{i-1}{n} \leq x < \frac{i}{n}, \frac{j-1}{n} \leq y < \frac{j}{n}, v_i v_j \in E(G), \\ 0, & \text{if } \frac{i-1}{n} \leq x < \frac{i}{n}, \frac{j-1}{n} \leq y < \frac{j}{n}, v_i v_j \notin E(G). \end{cases}$$

For a set $X = \{x_1, x_2, \dots, x_k\}$ of real numbers in $[0, 1]$, $B(X)$ is the graph with vertex set X and edge set $\{\{x_i, x_j\} : B(x_i, x_j) = 1\}$. For a graph H , the *induced density* of H in B is

$$d(H) = d_B(H) = \mathbb{P}(B(X) \cong H),$$

where X is chosen uniformly at random from $[0, 1]^{|H|}$. Note that this quantity equals the limit of the densities of H in balanced blow-ups of G on N vertices, where $N \rightarrow \infty$. This notion of a graphon agrees with the theory of dense graph limits developed by Lovász and Szegedy ([5], see the book by Lovász [4] for a thorough introduction to the theory). We use a different parametrization to follow the theory of flag algebras by Razborov.

In the following, we assume some familiarity with flag algebras, and we provide only a very brief reminder of the definitions. For a proper introduction to flag algebras, see [7].

We will need the notion of a flag. A *flag* is a graph F on n vertices, in which some vertices, say the first k vertices $X = \{x_1, \dots, x_k\} \subseteq V(F)$, are labeled with distinct labels. The *type* σ of F is the labeled graph induced on X . Two flags are isomorphic if there exists an isomorphism of the underlying graphs which induces an isomorphism of the labeled vertices. The graphon B of the flag F is the graphon of the unlabeled graph underlying F , together

with the set $Y = \{\frac{0}{n}, \frac{1}{n}, \dots, \frac{k-1}{n}\} \in [0, 1]^{|X|}$. Note that $B(Y) = \sigma$. For any flag H of type σ , we then have

$$d(H) = d_B(H) = \mathbb{P}(B(X \cup Y) \cong H),$$

where X is chosen uniformly at random from $[0, 1]^{|H|-|Y|}$.

A *Flag Algebra* \mathcal{F}^σ consists of formal linear combinations of flags of the type σ , with the canonical definitions for addition and scalar multiplication. A multiplication of flags is also defined in a way that it naturally corresponds to the multiplication of flag densities in graphons of flags.

2 Proof of Theorem 2

The core of the proof of Theorem 2 is the following lemma.

Lemma 3. *If G is a triangle-free graph maximizing the number of 5-cycles on $n \geq 10$ vertices, then there exists a 5-cycle C in G such that every other vertex of G has exactly two neighbors in C .*

Before we prove the lemma, we first show how it implies Theorem 2.

Proof of Theorem 2. Let G be a triangle-free graph maximizing the number of C_5 s on n vertices. For $n \leq 9$, we use a computer to enumerate all 2480 triangle-free graphs on n vertices and check that the theorem is true. Hence we assume $n \geq 10$ and we can use Lemma 3.

Let $C = v_1v_2v_3v_4v_5v_1$ be the 5-cycle from Lemma 3. For $1 \leq i \leq 5$, let X_i be the set of vertices which have the same neighbors in C as v_i . Since G is triangle free, these five sets are independent sets and partition $V(G)$. In the following, all indices are calculated modulo 5. Vertices in X_i are not adjacent to vertices in X_{i+2} since v_{i+1} is their common neighbor, for all $1 \leq i \leq 5$. Therefore, for any $u_i \in X_i$ and $u_{i+1} \in X_{i+1}$, $N(u_i) \cap N(u_{i+1}) = \emptyset$.

If $u_i \in X_i$ and $u_{i+1} \in X_{i+1}$, then $u_iu_{i+1} \in E(G)$. Otherwise, we could add u_iu_{i+1} , creating a new C_5 in $u_iu_{i+1}v_{i+2}v_{i+3}v_{i+4}u_i$ without creating a triangle, contradicting the maximality of C_5 . Hence, $X_i \cup X_{i+1}$ induces a complete bipartite graph, and G is a blow-up of C_5 .

Denote $|X_i|$ by x_i for $1 \leq i \leq 5$, so $n = x_1 + x_2 + x_3 + x_4 + x_5$. Then the number of C_5 in G is $x_1x_2x_3x_4x_5$. This number is maximized if and only if $|x_i - x_j| \leq 1$ for all $0 \leq i \leq j \leq 5$. Therefore, G is a balanced blow-up of C_5 , proving Theorem 2. \square

Denote by C_5^+ the balanced blow-up of C_5 on 6 vertices, i.e., the graph obtained from C_5 by duplicating one vertex. The proof of Lemma 3 uses the following results that were obtained using flag algebras. For any triangle-free graphon B of a graph:

Claim 4 ([2, 3]). $d(C_5) \leq 0.0384$.

Claim 5. *If $d(C_5) \geq 0.034$, then $d(C_5^+) \geq 4.57771 \cdot (d(C_5) - 0.034) + 0.095058$.*

Claim 6. $d(C_5^+) \geq 6 \cdot (d(C_5) - 0.0384) + 0.1152$.

We defer the proofs of Claims 5 and 6 to the next section. With these claims, we are ready to prove the core lemma.

Proof of Lemma 3. Let G be a triangle-free graph on $n \geq 10$ vertices maximizing the number of C_5 s, and let B be its graphon. We want to show that there exists a 5-cycle $C = v_1v_2v_3v_4v_5v_1$ in G such that $V(G) = X_1 \cup \dots \cup X_5$, where $v \in X_i$ if v has the same neighbors in C as v_i for $1 \leq i \leq 5$. In B , this translates to

$$\exists x_1, x_2, x_3, x_4, x_5 \in [0, 1] : \mathbb{P}(B[\{x, x_1, x_2, x_3, x_4, x_5\}] \cong C_5^+) = 1$$

for $x \in [0, 1]$ picked uniformly at random.

In fact, by the integrality of the number of vertices of G , it is sufficient to show that

$$\exists x_1, x_2, x_3, x_4, x_5 \in [0, 1] : \mathbb{P}(B[\{x, x_1, x_2, x_3, x_4, x_5\}] \cong C_5^+) > 1 - \frac{1}{n}. \quad (1)$$

We may condition on $B[x_1, x_2, x_3, x_4, x_5] \cong C_5$, and then, picking $x, x_1, x_2, x_3, x_4, x_5 \in [0, 1]$ independently uniformly at random,

$$\begin{aligned} & \mathbb{P}(B[\{x, x_1, x_2, x_3, x_4, x_5\}] \cong C_5^+ \mid B[\{x_1, x_2, x_3, x_4, x_5\}] \cong C_5) \\ &= \frac{\mathbb{P}(B[\{x, x_1, x_2, x_3, x_4, x_5\}] \cong C_5^+ \wedge B[\{x_1, x_2, x_3, x_4, x_5\}] \cong C_5)}{\mathbb{P}(B[\{x_1, x_2, x_3, x_4, x_5\}] \cong C_5)} \\ &= \frac{\frac{2}{6}d(C_5^+)}{d(C_5)}. \end{aligned}$$

Let G_n be a balanced blow up of C_5 on n vertices, and let B_n be its graphon. Let $d_n = d_{B_n}(C_5) \leq d(C_5)$. For $10 \leq n < 100$, we use Claim 5, and get

$$\frac{d(C_5^+)}{3d(C_5)} \geq \frac{4.57771 \cdot (d(C_5) - 0.034) + 0.095058}{3d(C_5)} \geq \frac{4.57771 \cdot (d_n - 0.034) + 0.095058}{3d_n}.$$

We compute d_n explicitly for each n , and conclude that $\frac{d(C_5^+)}{3d(C_5)} > 1 - \frac{1}{n}$, implying (1).

For $n \geq 100$, we use Claim 6 and Claim 4, and get

$$\begin{aligned} \frac{d(C_5^+)}{3d(C_5)} &\geq \frac{6 \cdot (d(C_5) - 0.0384) + 0.1152}{3d(C_5)} \\ &\geq \frac{6 \cdot (d_n - 0.0384) + 0.1152}{0.1152}. \end{aligned} \quad (2)$$

For $n \equiv i \pmod{5}$, we have

$$d_n = \frac{5! \left(\frac{n+5-i}{5}\right)^i \left(\frac{n-i}{5}\right)^{5-i}}{n^5} > 0.0384 \left(1 - \frac{50}{n^2}\right),$$

where the last inequality can easily be checked for the five cases. Therefore, using (2),

$$\frac{d(C_5^+)}{3d(C_5)} > \frac{-6 \cdot 0.0384 \frac{50}{n^2} + 0.1152}{0.1152} = 1 - \frac{100}{n^2} \geq 1 - \frac{1}{n},$$

showing (1) and thus completing the proof. \square

3 Flag algebra calculations

In this section we prove Claims 5 and 6. Claim 4 was obtained in [2] and [3] by flag algebra computations which can easily be replicated by Flagmatic [8], including certificates.

Both Claims 5 and 6 are based on similar flag algebra calculations, but they exceed the current limitations of Flagmatic. We describe an outline here. The actual calculations are computer assisted. The source code of our software to recreate the calculations is available on arXiv and at <http://orion.math.iastate.edu/lidicky/pub/c5k3free/>, including all data files.

The following equations are valid for all graphons.

$$d(C_5^+) = \sum_{F \in \mathcal{F}_\ell} c_F d(F), \quad (3)$$

$$0 \geq \sum_{F \in \mathcal{F}_\ell} -\alpha_F d(F), \quad (4)$$

$$0 \geq -\sum_{\sigma} \llbracket x_{\sigma}^T M_{\sigma} x_{\sigma} \rrbracket_{\sigma}, \quad (5)$$

where ℓ is an integer at least 6, \mathcal{F}_ℓ is the set of all triangle-free graphs on ℓ vertices, c_F are suitably picked rational non-negative coefficients, α_F are any non-negative real numbers, x_{σ} are vectors of some flag densities of type σ , M_{σ} are any positive semidefinite matrices, and $\llbracket \cdot \rrbracket$ is the unlabeled operator.

In addition, if the graphon satisfies $d(C_5) \geq LB$ for some fixed number LB , then for any non-negative real number y holds

$$0 \geq -y \cdot (d(C_5) - LB). \quad (6)$$

Now we sum all the equations (3) to (6) and we get

$$d(C_5^+) \geq \left[\sum_{F \in \mathcal{F}_\ell} (c_F - \alpha_F) d(F) - \sum_{\sigma} \llbracket x_{\sigma}^T M_{\sigma} x_{\sigma} \rrbracket_{\sigma} - y \cdot d(C_5) \right] + y \cdot LB, \quad (7)$$

The maximum of the right-hand side can be obtained by solving a semidefinite program in variables α_F , M_{σ} , and y . Notice that for any assignment of the variables, (7) is valid for any graphon satisfying $d(C_5) \geq LB$. So for a fixed assignment of variables, we can change LB and obtain different lower bounds on $d(C_5^+)$, albeit not necessarily optimal bounds.

We optimize (7) for $LB \in \{0.034, 0.0384\}$ and obtain Claims 5 and 6 by rounding the solution to the corresponding semidefinite program. The rounding for Claim 5 is easy as the bound is not tight. The bound in Claim 6 is tight for $d(C_5) = 0.0384$, and the rounding of the semidefinite program required a bit more care.

References

- [1] P. Erdős. On some problems in graph theory, combinatorial analysis and combinatorial number theory. In *Graph theory and combinatorics (Cambridge, 1983)*, pages 1–17. Academic Press, London, 1984.
- [2] A. Grzesik. On the maximum number of five-cycles in a triangle-free graph. *Journal of Combinatorial Theory, Series B*, 102(5):1061–1066, 2012.
- [3] H. Hatami, J. Hladký, D. Král', S. Norin, and A. Razborov. On the number of pentagons in triangle-free graphs. *Journal of Combinatorial Theory, Series A*, 120:722–732, 2013.
- [4] L. Lovász. *Large Networks and Graph Limits*, volume 60 of *Colloquium Publications*. American Mathematical Society, 2012.
- [5] L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.
- [6] T. Michael. Cycles of length 5 in triangle-free graphs: A sporadic counterexample to a characterization of equality. manuscript, 2011.
- [7] A. A. Razborov. Flag algebras. *J. Symbolic Logic*, 72(4):1239–1282, 2007.
- [8] E. Vaughan. Flagmatic 2.0, 2012. <http://flagmatic.org>.