

4-2018

Polychromatic colorings of complete graphs with respect to 1-, 2-factors and Hamiltonian cycles

Maria Axenovich

Karlsruhe Institute of Technology

John Goldwasser

West Virginia University

Ryan Hansen

West Virginia University

Bernard Lidicky

Iowa State University, lidicky@iastate.edu

Ryan R. Martin

Iowa State University, rymartin@iastate.edu

See next page for additional authors

Follow this and additional works at: https://lib.dr.iastate.edu/math_pubs



Part of the [Mathematics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/math_pubs/188. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Mathematics at Iowa State University Digital Repository. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Polychromatic colorings of complete graphs with respect to 1-, 2-factors and Hamiltonian cycles

Abstract

If G is a graph and H is a set of subgraphs of G , then an edge-coloring of G is called H -polychromatic if every graph from H gets all colors present in G on its edges. The H -polychromatic number of G , denoted $\text{polyH}(G)$, is the largest number of colors in an H -polychromatic coloring. In this paper, $\text{polyH}(G)$ is determined exactly when G is a complete graph and H is the family of all 1-factors. In addition $\text{polyH}(G)$ is found up to an additive constant term when G is a complete graph and H is the family of all 2-factors, or the family of all Hamiltonian cycles.

Disciplines

Mathematics

Comments

This is the peer reviewed version of the following article: Axenovich M, Goldwasser J, Hansen R, Lidický B, Martin RR, Offner D, Talbot J, Young M. Polychromatic colorings of complete graphs with respect to 1-, 2-factors and Hamiltonian cycles. *J Graph Theory*. 2018;87:660–671, which has been published in final form at doi: [10.1002/jgt.22180](https://doi.org/10.1002/jgt.22180). This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Authors

Maria Axenovich, John Goldwasser, Ryan Hansen, Bernard Lidicky, Ryan R. Martin, David Offner, John Talbot, and Michael Young

Polychromatic colorings of complete graphs with respect to 1-, 2-factors and Hamiltonian cycles

Maria Axenovich* John Goldwasser† Ryan Hansen‡
Bernard Lidický§ Ryan R. Martin¶ David Offner|| John Talbot**
Michael Young††

July 7, 2017

Abstract

If G is a graph and \mathcal{H} is a set of subgraphs of G , then an edge-coloring of G is called \mathcal{H} -polychromatic if every graph from \mathcal{H} gets all colors present in G . The \mathcal{H} -polychromatic number of G , denoted $\text{poly}_{\mathcal{H}}(G)$, is the largest number of colors such that G has an \mathcal{H} -polychromatic coloring. In this paper, $\text{poly}_{\mathcal{H}}(G)$ is determined exactly when G is a complete graph and \mathcal{H} is the family of all 1-factors. In addition $\text{poly}_{\mathcal{H}}(G)$ is found up to an additive constant term when G is a complete graph and \mathcal{H} is the family of all 2-factors, or the family of all Hamiltonian cycles.

1 Introduction

If G is a graph and \mathcal{H} is a set of subgraphs of G , we say that an edge-coloring of G is \mathcal{H} -polychromatic if every graph from \mathcal{H} gets all colors present in G . The \mathcal{H} -polychromatic number of G , denoted $\text{poly}_{\mathcal{H}}(G)$, is the largest number of colors in an \mathcal{H} -polychromatic coloring. If an \mathcal{H} -polychromatic coloring of G uses $\text{poly}_{\mathcal{H}}(G)$ colors, it is called an optimal \mathcal{H} -polychromatic coloring of G .

*Karlsruhe Institute of Technology, Karlsruhe, Germany, maria.aksenovich@kit.edu.

†West Virginia University, Morgantown, WV, USA, jgoldwas@math.wvu.edu.

‡West Virginia University, Morgantown, WV, USA, rhansen@mail.wvu.edu

§Iowa State University, Ames, IA, USA, lidicky@iastate.edu. Supported by NSF grant DMS-1600390.

¶Iowa State University, Ames, IA, USA, rymartin@iastate.edu. Research supported in part by Simons Foundation Collaboration Grant (#353292, to R.R. Martin).

||Westminster College, New Wilmington, PA, USA, offnerde@westminster.edu.

**University College London, London, UK, j.talbot@ucl.ac.uk.

††Iowa State University, Ames, IA, USA, myoung@iastate.edu.

1.1 Background

Let Q_n denote the hypercube of dimension n . Let $G = Q_n$ and \mathcal{H} be the family of all subgraphs of G isomorphic to Q_d . If d is fixed and n is large, then Alon, Krech, and Szabó [3] showed that $\lfloor \frac{(d+1)^2}{4} \rfloor \leq \text{poly}_{\mathcal{H}}(Q_n) \leq \binom{d+1}{2}$. Offner [11] proved that the lower bound is tight for all sufficiently large values of n . Bialostocki [4] treated the special case when $d = 2$ and $n \geq 2$. Goldwasser *et al.* [9] considered the case where \mathcal{H} is the family of all subgraphs of Q_n isomorphic to a Q_d minus an edge or a Q_d minus a vertex.

If T is a tree and \mathcal{H} is the set of all paths of length at least r , then $\text{poly}_{\mathcal{H}}(T) = \lceil r/2 \rceil$, as was shown by Bollobás *et al.* [5]. When $G = K_n$ and \mathcal{H} is the set of all r -vertex cliques, $\text{poly}_{\mathcal{H}}(G)$ was considered by Erdős and Gyárfás [6, 10] with the respective colorings called balanced. When G is an arbitrary multigraph of minimum degree d , and \mathcal{H} is the set of all stars with center v and leaves $N(v)$, $v \in V(G)$, then it was shown by Alon *et al.* [2], that $\text{poly}_{\mathcal{H}}(G) \geq \lfloor (3d + 1)/4 \rfloor$. Goddard and Henning [7] considered vertex-colorings of graphs such that each open neighborhood gets all colors present in G .

Polychromatic colorings were also investigated for vertex-colored hypergraphs. These colorings are essential tools in studying covering problems which are of fundamental importance in general graph and hypergraph settings, especially in geometric hypergraphs, and they exhibit connections to VC-dimension, see [1, 2, 5, 12].

1.2 Main Results

In this paper, we consider the case where G is a complete graph and \mathcal{H} is a family of spanning subgraphs. Let $F_1 = F_1(n)$ be the family of all 1-factors of K_n , $F_2 = F_2(n)$ be the family of all 2-factors of K_n and $\text{HC} = \text{HC}(n)$ be the family of all Hamiltonian cycles of K_n . Our main results are as follows:

Theorem 1. *If n is an even positive integer, then $\text{poly}_{F_1}(K_n) = \lfloor \log_2 n \rfloor$.*

Theorem 2. *There exists a constant c such that $\lfloor \log_2 2(n + 1) \rfloor \leq \text{poly}_{F_2}(K_n) \leq \text{poly}_{\text{HC}}(K_n) \leq \lfloor \log_2 n \rfloor + c$. Moreover, $\lfloor \log_2 \frac{8(n-1)}{3} \rfloor \leq \text{poly}_{\text{HC}}(K_n)$.*

It is claimed in a follow-up paper [8], that in fact $\text{poly}_{F_2}(K_n) = \lfloor \log_2 2(n + 1) \rfloor$ and $\text{poly}_{\text{HC}}(K_n) = \lfloor \log_2 \frac{8(n-1)}{3} \rfloor$ for $n \geq 3$. However, the arguments there include more case analysis and greater detail than what is required for the small additive constant given in Theorem 2.

The paper is structured as follows. We start with basic definitions in Section 2. In Section 3, we give constructions of polychromatic colorings, which provide the lower bounds for Theorems 1 and 2. In Section 4, we prove Theorem 1. Section 5 contains the proof of Theorem 2.

2 Definitions

Let the vertices of K_n be denoted by v_1, v_2, \dots, v_n . An edge-coloring φ is *ordered at v_i* for $i \in [n]$ if there exists a color a , called the *main color at v_i* , such that $\varphi(v_i v_j) = a$ for all $j \in \{i + 1, \dots, n\}$. Notice that v_{n-1} and v_n are ordered with respect to any coloring. We define the main color of v_n to be the same as for v_{n-1} . A vertex v_i is *unitary* if there are colors $a \neq b$ such that v_i is incident with $n - 2$ edges colored a and one edge $v_i v_j$ colored b , where v_j is unitary with $n - 2$ incident edges colored b . For v_i unitary, we also call a the *main color* of v_i .

An edge-coloring is *ordered* if all vertices are ordered with respect to some ordering of vertices. See Figure 1 for an example of an ordered coloring. We call an edge-coloring *combed* if each vertex is either ordered or unitary.

Let φ be an ordered or combed coloring. The *inherited coloring* is the vertex-coloring φ' obtained by coloring each vertex with its main color. Its *inherited color class M_i of color i* is the set of all vertices v with $\varphi'(v) = i$. Let $M_t(j) = M_t \cap \{v_1, v_2, \dots, v_j\}$. In this paper, we shall always think of the ordered vertices as arranged on a horizontal line with v_i to the left of v_j if $i < j$. We say that an edge $v_i v_j$, $i < j$, goes from v_i to the right and from v_j to the left. If φ is an edge-coloring of a graph G , the *maximum monochromatic degree* of G is the largest integer d such that some vertex of G is incident with d edges of the same color. We say such a vertex is a *max-vertex*. If X is a subset of $V(K_n)$, we say that the edge-coloring φ of K_n is

- *X-constant* if for any $v \in X$, $\varphi(vu) = \varphi(vw)$ for all $u, w \in V \setminus X$,
- *X-ordered* if there is an ordering v_1, \dots, v_n of the vertices of K_n such that $X = \{v_1, \dots, v_m\}$ for some integer m and φ is ordered on vertices in X .

Notice that if a coloring φ is X -ordered, then it is also X -constant.

3 Constructions of Polychromatic Colorings

We construct three edge-colorings of K_n , and show that they are polychromatic for F_1 , F_2 , and HC, respectively.

3.1 F_1 -polychromatic Coloring φ_{F_1}

Let $n \geq 2$ be even, and let k be the largest positive integer such that $2^k \leq n$, i.e., $k = \lfloor \log_2 n \rfloor$. Let φ' be a vertex-coloring of K_n with vertex set $\{v_1, \dots, v_n\}$ and colors $1, \dots, k$, where for each $i \in [k]$, M_i is the color class of color i . Moreover, for any $1 \leq i < j \leq k$, every vertex in M_i precedes every vertex in M_j , and $|M_t| = 2^{t-1}$ for $t = 1, \dots, k - 1$. Hence the color




Figure 1: F_1 -polychromatic coloring φ_{F_1} .




Figure 2: F_2 -polychromatic coloring φ_{F_2} .

87 classes M_1, M_2, \dots, M_k have sizes $1, 2, 4, \dots, 2^{k-2}, n - 2^{k-1} + 1$, respectively. Let φ_{F_1} be the
 88 ordered coloring for which φ' is the inherited coloring.

89 Consider an arbitrary 1-factor F of K_n and $t \in [k]$. Consider the set F_t of all edges
 90 of F with at least one endpoint in M_t . Since $|M_1| + \dots + |M_i| = 2^i - 1$ and $|M_k| =$
 91 $n - |M_1| - \dots - |M_{k-1}| \geq 2^k - 2^{k-1} + 1 = 2^{k-1} + 1$, we have $\sum_{i < t} |M_i| < |M_t|$ for any $t \in [k]$.
 92 Thus at least one edge of F_t joins a vertex from M_t to a vertex to the right, so this edge is
 93 of color t . Therefore F has edges of each color. Hence φ_{F_1} is F_1 -polychromatic and it uses
 94 $\lceil \log_2 n \rceil$ colors.

95 3.2 F_2 -polychromatic Coloring φ_{F_2}

96 Let k be the largest positive integer such that $n \geq 2^{k-1} - 1$, i.e., $k = 1 + \lfloor \log_2(n + 1) \rfloor$. Let
 97 φ' be a vertex-coloring of K_n with vertex set $\{v_1, \dots, v_n\}$ and colors $1, \dots, k$, where for each
 98 $i \in [k]$, M_i is the color class of color i . Moreover, for any $1 \leq i < j \leq k$, every vertex in M_i
 99 precedes every vertex in M_j , and $|M_t| = 2^{t-2}$ for $t = 4, \dots, k-1$, and $|M_1| = |M_2| = |M_3| = 1$.
 100 Hence the color classes $M_1, M_2, \dots, M_{k-1}, M_k$ have sizes $1, 1, 1, 4, 8, \dots, 2^{k-3}, n - 2^{k-2} + 1,$
 101 respectively. Let φ_{F_2} be obtained by taking the ordered coloring for which φ' is the inherited
 102 coloring and then recoloring the edge v_1v_3 from color 1 to color 3. See Figure 2.




Figure 3: HC-polychromatic coloring φ_{HC} .

103 Observe that the inherited color classes M_1 , M_2 , and M_3 contain unitary vertices. More-
 104 over, $|M_1| + \dots + |M_t| = 2^{t-1} - 1$ for $3 \leq t \leq k - 1$, and $|M_k| = n - |M_1| - \dots - |M_{k-1}| \geq$
 105 $2^{k-1} - 1 - 2^{k-2} + 1 = 2^{k-2}$. So, $|M_t| > \sum_{i < t} |M_i|$ for any $t \geq 4$. Consider an arbitrary 2-factor
 106 F of K_n and $t \in [k]$. For $i \leq 3$, v_i is a unitary vertex with main color i , so F must have
 107 edges of colors 1, 2, and 3. For a color $t \geq 4$, consider the set F_t of edges of F with least one
 108 endpoint in M_t . Then F_t has an edge of color t unless F_t forms a bipartite graph G_t with one
 109 part M_t and another $M'_t = \bigcup_{i=1}^{t-1} M_i$. The degree of each vertex of G_t from M_t is two, and
 110 the degree of each vertex of G_t from M'_t is at most two. Thus $|M'_t| \geq |M_t|$, a contradiction.
 111 Thus, F_t , and therefore F , has at least one edge of color t . So, φ_{F_2} is F_2 -polychromatic and
 112 it uses $k = \lfloor \log_2 2(n+1) \rfloor$ colors.

113 3.3 HC-polychromatic Coloring φ_{HC}

114 Let k be the largest positive integer such that $n \geq 3 \cdot 2^{k-3} + 1$, i.e., $k = 3 + \lfloor \log_2(n-1)/3 \rfloor$.
 115 Let φ' be a vertex-coloring of K_n with vertex set $\{v_1, \dots, v_n\}$ and colors $1, \dots, k$, where
 116 for each $i \in [k]$, M_i is the color class of color i . Moreover, for any $1 \leq i < j \leq k$,
 117 every vertex in M_i precedes every vertex in M_j , and $|M_t| = 3 \cdot 2^{t-4}$ for $t = 4, \dots, k-1$,
 118 and $|M_1| = |M_2| = |M_3| = 1$. Hence the color classes $M_1, M_2, \dots, M_{k-1}, M_k$ have sizes
 119 $1, 1, 1, 3, 6, 12, \dots, 3 \cdot 2^{k-5}, n - 3 \cdot 2^{k-4}$, respectively. Let φ_{HC} be obtained by taking the
 120 ordered coloring for which φ' is the inherited coloring and then recoloring the edge $v_1 v_3$ from
 121 color 1 to color 3. See Figure 3.

122 We have that $|M_1| + \dots + |M_t| = 3 \cdot 2^{t-3}$ for $3 \leq t \leq k-1$. Moreover, $|M_k| =$
 123 $n - |M_1| - \dots - |M_{k-1}| \geq 3 \cdot 2^{k-3} + 1 - 3 \cdot 2^{k-4} = 3 \cdot 2^{k-4} + 1$. Thus $|M_k| > \sum_{i < k} |M_i|$ and
 124 $|M_t| \geq \sum_{i < t} |M_i|$ for all $t \geq 4$. Consider an arbitrary Hamiltonian cycle H of K_n . For $i \leq 3$,
 125 v_i is a unitary vertex with main color i , so H must have edges of colors 1, 2, and 3. For
 126 each color $t \geq 4$, let H_t be the set of edges of H with at least one endpoint in M_t . Then H_t
 127 has an edge of color t unless H_t forms a bipartite graph G_t with one part M_t and another

128 $M'_t = \bigcup_{i=1}^{t-1} M_i$. The degree of each vertex of G_t from M_t is two, and the degree of each
 129 vertex of G_t from M'_t is at most two. If $4 \leq t < k$, $|M_t| = |M'_t|$, the degree of each vertex of
 130 G_t from M'_t is also two. Hence G_t is a union of cycles, so it could not be a proper subgraph
 131 of a Hamiltonian cycle. If $t = k$, $|M_k| > |M'_k|$, and hence such a bipartite graph G_t could
 132 not exist. Thus H has an edge of color t for each $t = 1, \dots, k$, φ_{HC} is HC-polychromatic,
 133 and it uses $\left\lceil \log_2 \frac{8(n-1)}{3} \right\rceil$ colors.

134 4 Proof of Theorem 1

135 We prove Theorem 1 by first showing the existence of an optimal edge-coloring that is
 136 ordered. Then we use Lemma 3 below which states that, for every inherited color class M_t ,
 137 there exists j such that a majority of v_1, \dots, v_j is in M_t . This leads to a counting argument
 138 that gives the upper bound in Theorem 1. For the lower bound we use the coloring φ_{F_1} .

139 **Lemma 3.** *Let $\varphi : E(K_n) \rightarrow [k]$, where n is even, be an ordered coloring with inherited*
 140 *color classes M_1, \dots, M_k . If the coloring φ is F_1 -polychromatic, then $\forall t \in [k] \exists j \in [n-1]$*
 141 *such that $|M_t(j)| > j/2$.*

142 *Proof.* Assume there exists t such that for each $j \in [n-1]$, $|M_t(j)| \leq j/2$. We will show
 143 that there exists a 1-factor where each vertex x in M_t is joined to some vertex y to the left
 144 and outside of M_t . Let x_1, \dots, x_m be the vertices of M_t in order and let y_1, \dots, y_{n-m} be the
 145 other vertices of K_n in order. Let H consist of the edges $y_1x_1, y_2x_2, \dots, y_mx_m$ and a perfect
 146 matching on $\{y_{m+1}, \dots, y_{n-m}\}$ (if this set is non-empty). Since $|M_t(j)| \leq j/2$ for all j , the
 147 number of y 's that must precede x_i is at least i for each $i = 1, \dots, m$. Hence y_i is to the left
 148 of x_i and not in M_t for each $i = 1, \dots, m$. Therefore all edges in H incident with vertices in
 149 M_t go to the left and do not have color t . The edges of H that are not incident with vertices
 150 in M_t are also not of color t . Hence φ is not F_1 -polychromatic, a contradiction. \square

151 *Proof of Theorem 1.* By subsection 3.1, $\text{poly}_{F_1}(G)$ is well-defined. Let $k = \text{poly}_{F_1}(K_n)$ be the
 152 polychromatic number for 1-factors in $K_n = (V, E)$. Among all F_1 -polychromatic colorings
 153 of K_n with k colors we choose ones that are X -ordered for a subset X (possibly empty) of
 154 the largest size, and, of these, choose a coloring φ whose restriction to $V \setminus X$ has the largest
 155 maximum monochromatic degree. Suppose for contradiction that $V \neq X$.

156 Let $Z = V \setminus X$ and G be the subgraph of K_n induced by Z . Let v be a vertex of maximum
 157 monochromatic degree, d , in φ restricted to G , and let 1 be a color for which there are d
 158 edges incident with v in G with color 1. By the maximality of $|X|$, there is a vertex u in
 159 Z such that $\varphi(uv) \neq 1$. Assume $\varphi(uv) = 2$. If every 1-factor containing uv had another
 160 edge of color 2, then the color of uv could be changed to 1, resulting in an F_1 -polychromatic
 161 coloring where v has a larger maximum monochromatic degree in G , a contradiction. Hence,
 162 there is a 1-factor F in which uv is the only edge with color 2 in φ .

163 Let $\varphi(vy_i) = 1$, $y_i \in Z$, $i = 1, \dots, d$. For each $i \in [d]$, let y_iw_i be the edge of F containing
 164 y_i (perhaps $w_i = y_j$ for some $j \neq i$); see Figure 4. We can get a different 1-factor F_i by
 165 replacing the edges uv and y_iw_i in F with edges vy_i and uw_i . Since F_i must have edges of
 166 all colors, it has an edge of color 2. As $\varphi(vy_i) = 1$, we must have $\varphi(uw_i) = 2$ for each $i \in [d]$.




Figure 4: Maximum polychromatic degree in an F_1 -polychromatic coloring.

167 If $w_i \in X$ for some i then, since φ is X -constant, $\varphi(w_iy_i) = \varphi(w_iu) = 2$, so y_iw_i
 168 and uv are two edges of color 2 in F , a contradiction. So, $w_i \in Z$ for all $i \in [d]$. Thus
 169 $\varphi(uv) = \varphi(uw_1) = \dots = \varphi(uw_d) = 2$, and the monochromatic degree of u in G is at least
 170 $d + 1$, larger than that of v , a contradiction.

171 We conclude that $X = V$. Hence φ is an ordered F_1 -polychromatic coloring of K_n . By
 172 Lemma 3, for every $t \in [k]$ there exists j_t such that $|M_t(j_t)| > \frac{j_t}{2}$. By permuting the colors,
 173 we can assume $j_{t_1} < j_{t_2}$ whenever $t_1 < t_2$. This gives us an ordering of inherited color
 174 classes M_1, M_2, \dots, M_k . Since $|M_1| \geq 1$ and $|M_t(j_t)| > \frac{j_t}{2}$, we can use induction to show
 175 $|M_t| \geq |M_t(j_t)| \geq 2^{t-1}$ as follows

$$|M_t| \geq |M_t(j_t)| > \sum_{1 \leq i < t} |M_i(j_t)| \geq \sum_{1 \leq i < t} |M_i(j_i)| \geq \sum_{1 \leq i < t} 2^{i-1} = 2^{t-1} - 1.$$

176 The sum of the sizes of all inherited color classes is n , and we get

$$n = \sum_{t=1}^k |M_t| \geq \sum_{t=1}^k 2^{t-1} = 2^k - 1.$$

177 Since n is even, $n \geq 2^k$ and $\text{poly}_{F_1}(K_n) = k \leq \lfloor \log_2 n \rfloor$.

178 The fact that $\text{poly}_{F_1}(K_n) \geq \lfloor \log_2 n \rfloor$ follows from the coloring φ_{F_1} . This finishes the
 179 proof of Theorem 1. \square

180 5 Proof of Theorem 2

181 Recall that we call an edge-coloring φ *combed* if all vertices are either ordered or unitary.

182 We prove Theorem 2 by first showing the existence of an optimal edge-coloring that is
 183 combed. Then we use Lemma 4 below which states that, for every inherited color class M_t ,
 184 either there exists j such that at least half of v_1, \dots, v_j is in M_t or M_t contains a unitary
 185 vertex. This leads to a counting argument that finishes the proof of Theorem 2.

186 **Lemma 4.** *Let $\varphi : E(K_n) \rightarrow [k]$ be a combed coloring with inherited color classes M_1, \dots, M_k .
 187 If the coloring φ is F_2 -polychromatic, or HC-polychromatic, then $\forall t \in [k] \exists j \in [n-1]$ such
 188 that $|M_t(j)| \geq \frac{j}{2}$ or M_t contains a unitary vertex.*

189 *Proof.* Let $\mathcal{H} \in \{F_2, \text{HC}\}$. Let φ be a combed \mathcal{H} -polychromatic coloring with inherited color
 190 classes M_1, \dots, M_k . It is sufficient to consider an arbitrary color $t \in [k]$ and show that the
 191 condition on M_t is satisfied.

192 Let x_1, \dots, x_m be the vertices of M_t in order and let y_1, \dots, y_{n-m} be the other vertices
 193 of K_n in order. Suppose for contradiction that there exists t such that $|M_t(j)| < \frac{j}{2}$ for
 194 all $j \in [n-1]$ and M_t does not contain any unitary vertex. Thus φ is ordered at each
 195 $x_i \in M_t$ and so y_{i+1} is to the left of x_i for each $i \in [m]$. Consider a Hamiltonian cycle
 196 $H = y_1x_1y_2x_2 \cdots y_mx_my_{m+1} \cdots y_{n-m}y_1$.

197 Since $|M_t(j)| < j/2$ for all j , the number of y 's that must precede x_i is at least $i+1$ for
 198 each $i = 1, \dots, m$. Hence y_i and y_{i+1} are to the left of x_i for each $i = 1, \dots, m$. Therefore
 199 each edge in H incident with a vertex x_i in M_t goes to the left from the perspective of x_i .

200 Let yx be an edge of H , where $x \in M_t$. Since $y \notin M_t$, the majority color r of y is not
 201 t . Since φ is combed, either $\varphi(yx) = r$ or $\varphi(yx) \neq r$ and both y and x are unitary vertices.
 202 Recall M_t does not contain any unitary vertices. Hence no edge in H is colored by t . This
 203 contradicts the fact that φ is \mathcal{H} -polychromatic. \square

204 We say that a Hamiltonian cycle H' is obtained from another Hamiltonian cycle H by a
 205 *twist* of disjoint edges e_1 and e_2 of H if $E(H) \setminus \{e_1, e_2\} \subseteq E(H')$, i.e. we remove e_1, e_2 from
 206 H and introduce two new edges to make the resulting graph a Hamiltonian cycle. Note that
 207 the choice of these two edges to add is unique. The other choice of two edges to add does
 208 not preserve connectivity. Without the connectivity requirement, the operation is known as
 209 a *2-switch*.

210 Notice that any 2-switch could be applied to a 2-factor and the result will be again a
 211 2-factor. Here, it might be possible to add the two new edges in two different ways.

212 For $\mathcal{H} \in \{\text{HC}, F_2\}$, Lemma 5 can be used to show that there exists an optimal \mathcal{H} -
 213 polychromatic coloring that is combed.

214 **Lemma 5.** *Suppose $n \geq 3$ and $X \subset V(K_n)$. Let $\mathcal{H} \in \{\text{HC}, F_2\}$ and φ_1 be an optimal
 215 \mathcal{H} -polychromatic coloring of K_n that is X -constant. Then there exists an optimal \mathcal{H} poly-
 216 chromatic coloring φ of K_n that agrees with φ_1 on all edges with at least one endpoint in X
 217 such that*

- 218 (A) there exists a vertex $v \in V(K_n) \setminus X$ such that φ is $(X \cup \{v\})$ -constant; or
 219 (B) $X = \emptyset$ and there exist vertices x, y, z , such that these vertices are unitary under φ of
 220 distinct main colors. This implies φ is $\{x, y, z\}$ -constant and xyz is a rainbow triangle.

221 *Proof.* Let $\mathcal{H} \in \{F_2, \text{HC}\}$. Let $Z = V(K_n) \setminus X$ and G be the subgraph of K_n induced by Z .
 222 Let $|Z| = m$. If $m \leq 2$ then $X \neq \emptyset$ and (A) is trivially satisfied. Hence $m \geq 3$. If $X = \emptyset$ and
 223 there exists an optimal \mathcal{H} -polychromatic coloring φ with three unitary vertices x, y , and z
 224 of distinct main colors, then (B) is satisfied. Hence we assume there is no such edge-coloring
 225 φ .

226 Choose φ to be an optimal \mathcal{H} -polychromatic coloring such that it agrees with φ_1 on
 227 edges with at least one endpoint in X and subject to this, it maximizes the maximum
 228 monochromatic degree of G . Define d to be the maximum monochromatic degree of vertices
 229 in G with φ .

230 First suppose $d = m - 1$. Let v be a vertex of maximum monochromatic degree d in G .
 231 Then φ is $(X \cup \{v\})$ -constant and we have (A). Hence we assume $d \leq m - 2$.

232 Let $\ell = m - 1 - d$ and let φ use colors $1, \dots, k$. If color a appears d times in G at a
 233 vertex $v \in Z$, we say v is an a -max-vertex. If the ℓ edges incident with v in G which do not
 234 have color a all have color b , we call v an (a, b) -max-vertex with *minority color* b .

235 **Claim 1.** If $a, b \in [k]$ are two distinct colors, $v \in Z$ is an a -max-vertex and $\varphi(vu) = b$ for
 236 some other vertex $u \in Z$, then all of the following hold:

- 237 (1) u is a b -max-vertex,
 238 (2) v is an (a, b) -max-vertex,
 239 (3) either $X = \emptyset$ or $\mathcal{H} = F_2$, and
 240 (4) at least half of the edges between X and Z have color b .

241 *Proof.* For ease of notation, we assume that $a = 1$ and $b = 2$. Let v be a 1-max-vertex. Let
 242 $u \in Z$ be a vertex such that $\varphi(vu) = 2$. If every $H \in \mathcal{H}$ containing uv contains another edge
 243 of color 2, we could change the color of uv to color 1, giving an \mathcal{H} -polychromatic coloring
 244 where v has monochromatic degree $d + 1$, a contradiction. Hence, there must be $H \in \mathcal{H}$
 245 where uv is the only edge of color 2.

246 Cyclically orient the edges of each cycle in H such that uv is an arc, and denote the
 247 resulting directed graph \vec{H} . Let $\varphi(vy_j) = 1$, for $y_j \in Z$, $j = 1, 2, \dots, d$. For each $j \in [d]$,
 248 let w_j be the predecessor of y_j in \vec{H} , so $\vec{w_j y_j} \in \vec{H}$ for each j . We assume $w_j \neq v$ for
 249 $j = 2, 3, \dots, d$, but perhaps $w_1 = v$ and perhaps $w_j = y_i$ for some $i \neq j$. See Figure 5.

250 Now we shall prove (1). If $w_j \neq v$, twist the edges uv and $w_j y_j$ of H to get a new
 251 $H_j \in \mathcal{H}$ containing vy_j and uw_j . Since H_j must have an edge of color 2 and $\varphi(vy_j) = 1$, we




Figure 5: Situation in Claim 1.

252 must have $\varphi(uw_j) = 2$. Hence, $\varphi(uw) = \varphi(uw_2) = \varphi(uw_3) = \dots = \varphi(uw_d) = 2$. Note that
 253 $w_j \in Z$ for each $j \in [d]$. This is because if $w_j \in X$, then, since φ is X -constant and $y_j \in Z$,
 254 $\varphi(w_j y_j) = \varphi(w_j u) = 2$, so $w_j y_j$ is another edge in H with color 2, a contradiction. That
 255 gives us d edges of color 2 at u in G . Note that if $w_1 \neq v$, then uw_1 is another edge of color
 256 2 incident with u , so $w_1 = v$ and $\vec{v y_1}$ is an arc of \vec{H} . Therefore, u is a 2-max-vertex. This
 257 proves (1).

258 Next, we prove (2), i.e., that v is a $(1, 2)$ -max-vertex. Let $z \in Z$ be a vertex distinct from
 259 u and v such that $\varphi(vz) \neq 1$. Let w be the vertex such that $\vec{w z}$ is an arc in \vec{H} . We know
 260 that $w \notin \{v = w_1, w_2, \dots, w_d, u\}$, since $z \notin \{y_1, \dots, y_d, v\}$. Let $H_z \in \mathcal{H}$, containing vz and
 261 uw , be obtained from H by twisting uv and wz . Since uv was the unique edge of H colored
 262 by 2, either $\varphi(uw) = 2$ or $\varphi(vz) = 2$. Suppose $w \in Z$. Since the maximum monochromatic
 263 degree is d and $\varphi(uw_j) = 2$ for all $j \in [d]$, $\varphi(uw) \neq 2$, so $\varphi(vz) = 2$. Suppose $w \in X$.
 264 Since $\varphi(wz) \neq 2$ and φ is X -constant, $\varphi(wz) = \varphi(uw) \neq 2$, so $\varphi(vz) = 2$. In both cases,
 265 $\varphi(vz) = 2$. Therefore, v is a $(1, 2)$ -max-vertex and the proof of (2) is done.

266 If $X = \emptyset$ then both (3) and (4) hold. So, assume that $X \neq \emptyset$. Let $H \in \mathcal{H}$. Assume that
 267 there is an edge of H with one endpoint in X and another in Z . Then there exist $x \in X$ and
 268 $y \in Z$ such that $\vec{y x}$ is an arc in \vec{H} . We know $y \notin \{v = w_1, \dots, w_d, u\}$, because the successor
 269 of y in \vec{H} is in X . If we twist yx and uv we get an $H_x \in \mathcal{H}$ containing uy and vx , where
 270 one of these edges must have color 2. However, since $\varphi(xv) = \varphi(xy) \neq 2$, we must have
 271 $\varphi(yu) = 2$, and u has monochromatic degree $d + 1$ in G , a contradiction. Hence there is no
 272 edge in \vec{H} with one endpoint in X and another in Z , and thus X induces a 2-factor in H .
 273 In particular, since $Z \neq \emptyset$, H is not a Hamiltonian cycle, and $\mathcal{H} = \mathcal{F}_2$. Let $p, q \in X$ with
 274 $pq \in E(H)$. Since both $(H \setminus \{uv, pq\}) \cup \{pv, qu\}$ and $(H \setminus \{uv, pq\}) \cup \{pu, qv\}$ are 2-factors

275 in \mathcal{H} , and φ is X -constant, either $\varphi(pv) = \varphi(pu) = 2$ or $\varphi(qv) = \varphi(qu) = 2$. In fact, since φ
 276 is X -constant, for each edge pq of H , where $p, q \in X$, all the edges from either p or q into Z
 277 have color 2. Since $H[X]$ is a union of cycles, at least half the edges between X and Z have
 278 color 2. This proves (3) and (4) and finishes the proof of Claim 1. \square

279 **Claim 2.** The graph G does not contain a $(1, 2)$ -max-vertex, a $(2, 3)$ -max-vertex, and a
 280 $(3, 1)$ -max-vertex at the same time.

281 *Proof.* Let x, y , and z be a $(1, 2)$ -max-vertex, a $(2, 3)$ -max-vertex, and a $(3, 1)$ -max-vertex,
 282 respectively. Applying Claim 1 to $\{v, u\} = \{x, y\}$, then $\{v, u\} = \{y, z\}$, and then with
 283 $\{v, u\} = \{z, x\}$, the conclusion (4) gives that at least half of the edges between X and Z
 284 have color 2, 3, and 1, respectively. Since colors 1, 2, and 3 are distinct, we conclude $X = \emptyset$.
 285 Let $H \in \mathcal{H}$. Observe that x, y , and z could be incident only with edges of H with colors in
 286 $\{1, 2, 3\}$ in φ , so all other colors in H come from edges not incident with these vertices.

287 Let φ^* be obtained from φ by the following modification

$$\varphi^*(uv) = \begin{cases} 1 & u = x \text{ and } v \neq y, \\ 2 & u = y \text{ and } v \neq z, \\ 3 & u = z \text{ and } v \neq x, \\ \varphi(uv) & \text{otherwise.} \end{cases}$$

288 Observe that the union of edges of H with at least one endpoint in $\{x, y, z\}$ contains all
 289 colors $\{1, 2, 3\}$ in φ^* . Hence H is polychromatic in φ^* and φ^* is \mathcal{H} -polychromatic. Moreover,
 290 φ^* is $\{x, y, z\}$ -constant and all the other properties of (B) hold, which is a contradiction.
 291 This finishes the proof of Claim 2. \square

292 **Claim 3.** If v is a $(1, 2)$ -max-vertex and $u \in Z$ such that $\varphi(uv) = 2$, then u is a $(2, 1)$ -max-
 293 vertex.

294 *Proof.* Let v be a $(1, 2)$ -max-vertex and $u \in Z$ such that $\varphi(uv) = 2$. Claim 1 implies that
 295 u is a $(2, \star)$ -max-vertex. Suppose for contradiction that u is a $(2, 3)$ -max-vertex. Since the
 296 number of edges incident with v colored 2 is the same as the number of edges incident with
 297 u colored 3 and $\varphi(uv) = 2$, there is a vertex x such that $\varphi(ux) = 3$ and $\varphi(vx) = 1$. Again
 298 by Claim 1, x is a $(3, 1)$ -max-vertex, contradicting Claim 2. \square

299 **Claim 4.** If there is a $(1, 2)$ -max-vertex, then there is no $(1, b)$ -max-vertex for any $b \neq 2$.

300 *Proof.* By symmetry suppose for contradiction that $v \in Z$ is a $(1, 2)$ -max-vertex and $u \in Z$
 301 is a $(1, 3)$ -max-vertex. Let $x \in Z$ be a vertex with $\varphi(vx) = 2$. By Claim 3, x is a $(2, 1)$ -max-
 302 vertex. Notice $\varphi(ux) \in \{1, 2\} \cap \{1, 3\} = \{1\}$. Now Claim 3 applied to x and u gives that u
 303 is a $(1, 2)$ -max-vertex, which is a contradiction. \square




Figure 6: Final part of proof of Lemma 5 with only (1,2)- and (2,1)-max-vertices, solid edges correspond to color 1, dotted edges correspond to color 2.

304 Claims 3 and 4 imply that if there is an (a,b) -max-vertex, then $\{a,b\} = \{1,2\}$.

305 Let S be the set of all (1,2)-max-vertices, T be the set of all (2,1)-max-vertices, and
 306 $W = Z \setminus (S \cup T)$. By Claim 3, both S and T are not empty. Edges within S and from S
 307 to W must have color 1 (because any minority color edge at a max-vertex is incident with
 308 a max-vertex of that color), edges within T and from T to W must have color 2, and each
 309 edge between S and T must have color 1 or 2.

310 Suppose $X = \emptyset$. Let $s \in S$ and $t \in T$. Let φ^* be obtained from φ by recoloring all edges
 311 incident with s to 1 and by recoloring all edges incident with t and not incident with s to 2.
 312 Notice that φ^* is \mathcal{H} -polychromatic. This contradicts the maximality of the monochromatic
 313 degree in φ . Therefore, $X \neq \emptyset$.

314 By symmetry, we can assume $|S| \leq |T|$. Let $v \in S$ and $u \in T$ be vertices such that
 315 $\varphi(vu) = 2$. By the maximality of the monochromatic degree of v in Z , there exists $H \in \mathcal{H}$,
 316 where uv is the unique edge colored by 2. Recall $\ell = m - 1 - d$. Let $s_1, \dots, s_\ell \in S$, where
 317 $\varphi(us_i) = 1$ for all $i \in [\ell]$. Let \vec{H} be a directed cycle(s) obtained by orienting edges of H such
 318 that $\vec{uv} \in \vec{H}$. Let t_i be a vertex such that $\vec{s_i t_i} \in \vec{H}$ for all $i \in [\ell]$. See Figure 6.

319 If there is an $i \in [\ell]$ such that $\varphi(vt_i) \neq 2$, then twist of vu and $s_i t_i$ contains vt_i and us_i
 320 and leaves no edge colored 2, which is a contradiction with φ being \mathcal{H} -polychromatic. Hence
 321 $\varphi(vt_i) = 2$ and $t_i \in T$ for all $i \in [\ell]$. Notice that v has exactly ℓ incident edges colored 2 and
 322 the other ends of these edges must be t_1, \dots, t_ℓ . By symmetry, we assume $t_1 = u$.

323 Suppose there exists \vec{xz} in \vec{H} with $x \in X$ and $z \in Z$. Since uv is the unique edge of
 324 \vec{H} colored 2, $\varphi(xz)$ is not 2 and since φ is X -constant, $\varphi(xz) = \varphi(xu) \neq 2$. Notice that
 325 $z \notin \{u = t_1, \dots, t_\ell\}$ since for every $i \in [\ell]$, the predecessor of t_i in \vec{H} is s_i and $s_i \in S$. Hence
 326 $\varphi(vz) = 1$ and the twist of xz and uv contains xu and vz . Since $\varphi(xu) \neq 2$ and $\varphi(vz) \neq 2$,
 327 we get a contradiction to φ being \mathcal{H} -polychromatic. Therefore, there is no edge of H between

328 X and Z .

329 Since there is no edge of H between X and Z and $X \neq \emptyset$, H is not connected. Therefore,
 330 $\mathcal{H} = F_2$.

331 Recall that all edges between T and W have color 2, hence they are not in H . Since
 332 there are no edges of H between X and Z , and all edges within T have color 2, every vertex
 333 in T has both neighbors from H in S . On the other hand, every vertex in S has at most two
 334 neighbors from H in T . Thus $|S| \geq |T|$. Recall we assumed $|S| \leq |T|$. Hence $|S| = |T|$ and
 335 there are no edges of H between $S \cup T$ and W .

336 Let J be the bipartite subgraph of G induced on parts S and T . Since vertices in T are
 337 $(2,1)$ -max-vertices, each of them is incident in J with ℓ edges of color 1 and $|S| - \ell$ edges of
 338 color 2. Similarly, each vertex in S is incident in J with ℓ edges of color 2 and $|T| - \ell = |S| - \ell$
 339 edges of color 1. Hence $|S| = 2\ell$ and the subgraph B of J consisting of all edges in J of color
 340 1 is an ℓ -regular bipartite graph.

341 If $\ell \geq 2$ then there exists a 2-factor K in B . Let H^* be obtained from H by removing
 342 edges incident with vertices in $S \cup T$ and adding K . Since all edges of K have color 1 and
 343 uv was the unique edge of H colored 2, we conclude H^* has no edge colored 2, contradicting
 344 the assumption that φ is \mathcal{H} -polychromatic.

345 Finally, if $\ell = 1$, then B is a matching on four vertices and the other two edges between
 346 S and T must have color 2. Hence $S \cup T$ does not contain a 2-factor in which uv would be
 347 the unique edge colored 2. This contradicts the existence of H .

348 This finishes the proof of Lemma 5. □

349 *Proof of Theorem 2.* Let $\mathcal{H} \in \{F_2, HC\}$. Let φ_1 be an optimal \mathcal{H} -polychromatic coloring of
 350 $E(K_n)$ with k colors and $[k]$ be the set of colors. We choose $X = \emptyset$, then we repeatedly
 351 apply Lemma 5. In the first application, we may get Lemma 5(B) and get $X = \{x, y, z\}$
 352 that are unitary of distinct colors or Lemma 5(A) and $|X| = 1$. But after that Lemma 5(A)
 353 always applies. Note that there are no unitary vertices except possibly x, y , and z because
 354 each other vertex is incident with edges with distinct colors c_x, c_y, c_z , the main colors of x ,
 355 y , and z . This results in a combed edge-coloring φ with zero or three first unitary vertices
 356 and all others being ordered vertices.

357 Let M_i be the inherited color classes obtained from φ . Let M_1, \dots, M_{k-3} be the inherited
 358 color classes not containing x, y , or z . By Lemma 4, for each color class M_t there is j_t such
 359 that $|M_t(j_t)| \geq \frac{j_t}{2}$. By symmetry, assume $j_i < j_t$ for all $1 \leq i < t \leq k - 3$. This leads to

$$|M_t(j_t)| \geq \sum_{i < t} |M_i(j_t)| \geq \sum_{i < t} |M_i(j_i)|$$

360 for $t = 2, \dots, k - 3$ and $|M_1| \geq 1$. Hence by induction we get

$$|M_t(j_t)| \geq 1 + \sum_{2 \leq i < t} |M_i(j_i)| \geq 1 + \sum_{2 \leq i < t} 2^{i-2} = 2^{t-2}.$$

361 Therefore, $|M_t| \geq 2^{t-2}$ for $t \geq 2$ and

$$n \geq \sum_{1 \leq t \leq k-3} |M_t| \geq 1 + \sum_{2 \leq t \leq k-3} 2^{t-2} \geq 2^{k-4}.$$

362 Hence $k \leq \log_2 n + 4$. By splitting the cases to (A) and (B), we could show $k \leq \log_2 n + 2$.

363 The lower bounds in Theorem 2 follow from colorings φ_{F_2} and φ_{HC} . Since every Hamil-
364 tonian cycle is also a 2-factor, we obtain $\text{poly}_{F_2}(K_n) \leq \text{poly}_{HC}(K_n)$. \square

365 6 Closing Remarks

366 We show above that c from Theorem 2 is at most 4. It is possible to get a more precise
367 version of Lemma 4 and use it to get sharp bounds in Theorem 2. We do not provide the
368 details in order to keep the paper short and less technical. Details should be in the follow-up
369 paper [8] together with generalizations which allow \mathcal{H} to be the family of all 1-regular or
370 2-regular graphs that span all but a fixed number of vertices.

371 7 Acknowledgements

372 This research was initiated at a workshop made possible by the Alliance for Building Fac-
373 ulty Diversity in the Mathematical Sciences (DMS 0946431). We would like to thank to
374 anonymous referees for their comments.

375 References

- 376 [1] E. Ackerman, B. Keszegh, and M. Vizer. Coloring Points with Respect to Squares.
377 In S. Fekete and A. Lubiw, editors, *32nd International Symposium on Computational
378 Geometry (SoCG 2016)*, volume 51 of *Leibniz International Proceedings in Informatics
379 (LIPIcs)*, pages 5:1–5:16, Dagstuhl, Germany, 2016. Schloss Dagstuhl–Leibniz-Zentrum
380 fuer Informatik.
- 381 [2] N. Alon, R. Berke, K. Buchin, M. Buchin, P. Csorba, S. Shannigrahi, B. Speckmann,
382 and P. Zumstein. Polychromatic colorings of plane graphs. *Discrete Comput. Geom.*,
383 42(3):421–442, 2009.
- 384 [3] N. Alon, A. Krech, and T. Szabó. Turán’s theorem in the hypercube. *SIAM J. Discrete
385 Math.*, 21(1):66–72, 2007.
- 386 [4] A. Bialostocki. Some Ramsey type results regarding the graph of the n -cube. *Ars
387 Combin.*, 16(A):39–48, 1983.

- 388 [5] B. Bollobás, D. Pritchard, T. Rothvoss, and A. Scott. Cover-decomposition and poly-
389 chromatic numbers. *SIAM J. Discrete Math.*, 27(1):240–256, 2013.
- 390 [6] P. Erdős and A. Gyárfás. Split and balanced colorings of complete graphs. *Discrete*
391 *Math.*, 200(1-3):79–86, 1999. Paul Erdős memorial collection.
- 392 [7] W. Goddard and M. Henning. Thoroughly distributed colorings. *arXiv:1609.09684*.
- 393 [8] J. Goldwasser and R. Hansen. Polychromatic colorings of 1-regular and 2-regular sub-
394 graphs of complete graphs. In preparation, 2017.
- 395 [9] J. Goldwasser, B. Lidický, R. Martin, D. Offner, J. Talbot, and M. Young. Polychro-
396 matic colorings on the hypercube. *arXiv:1603.05865*, 2016.
- 397 [10] A. Gyárfás. Problems and memories. *arXiv: 1307.1768v1*. Paul Erdős memorial collec-
398 tion.
- 399 [11] D. Offner. Polychromatic colorings of subcubes of the hypercube. *SIAM J. Discrete*
400 *Math.*, 22(2):450–454, 2008.
- 401 [12] J. Pach and G. Tóth. Decomposition of multiple coverings into many parts. *Comput.*
402 *Geom.*, 42(2):127–133, 2009.