

IOWA STATE UNIVERSITY
Digital Repository
Farm Progress Reports

2016 Report
Issue 1 2016 *Farm Progress Reports*

Number RFR-A1635

2017

Fairway Height Bentgrass Cultivar Trial

Nick Christians
Iowa State University, nchris@iastate.edu

Adam Thoms
Iowa State University, athoms@iastate.edu

Isaac Mertz
Iowa State University, imertz@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/farmprogressreports>



Part of the [Agriculture Commons](#), and the [Horticulture Commons](#)

Recommended Citation

Christians, Nick; Thoms, Adam; and Mertz, Isaac (2017) "Fairway Height Bentgrass Cultivar Trial," *Farm Progress Reports*: Vol. 2016 : Iss. 1 , Article 32.

DOI: <https://doi.org/10.31274/farmprogressreports-180814-1604>

Available at: <https://lib.dr.iastate.edu/farmprogressreports/vol2016/iss1/32>

This Horticulture Station is brought to you for free and open access by the Extension and Experiment Station Publications at Iowa State University Digital Repository. It has been accepted for inclusion in Farm Progress Reports by an authorized editor of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Fairway Height Bentgrass Cultivar Trial

RFR-A1635

Nick Christians, university professor
Adam Thoms, assistant professor
Isaac Mertz, graduate research assistant
Department of Horticulture

Introduction

This is the second year of data collection from the fairway height bentgrass trial established in the fall of 2014.

Materials and Methods

The area was maintained at a 0.5 in. mowing height. This is a National Turfgrass Evaluation (NTEP) trial and is being conducted at several research stations in the United States. It contains 17 of the newest cultivars and a number of experimental cultivars. The cultivars are maintained with 4 lb of N/1,000 ft² per growing season. Fungicides are used as needed in a preventive program. Herbicides and insecticides also are applied as needed.

Results and Discussion

The visual quality was evaluated monthly in 2016 from May through October (Table 1). The values listed under each month in Table 1 are the averages of visual quality ratings made on three replicated plots. Visual quality was based on a scale of 9 to 1: 9 = best quality, 6 = lowest acceptable quality, and 1 = worst quality. The annual average of the monthly data for each cultivar was calculated and is listed in the last column (Mean). The last row states the LSD (least significant difference), which is a statistical measurement of how widely the datum in each column must vary before they are considered to be different from one another. Genetic color was rated using a 9 to 1 scale with 9 = dark and 1 = light green.

Acknowledgements

Appreciation is extended to the NTEP organization for funding to conduct this project.

Table 1. Data for the fairway height creeping bentgrass trial, 2016.

No.	Cultivars	Color ²	Quality ¹						
			May	June	July	Aug	Sept	Oct	Mean
1	DLFPS-AT/3026	8	5	6	6	6	6	7	6
2	7	7	6	6	6	6	6	7	6
3	Penncross	7	5	6	6	5	5	7	6
4	Crystal Blue Links	8	6	7	6	7	6	6	6
5	Greentime	8	5	6	6	6	6	6	6
6	L-93XD	6	6	7	7	6	7	7	7
7	Armor	8	6	5	6	6	7	7	6
8	Kingdom	8	6	7	7	6	6	7	7
9	Nightlife	8	7	6	6	7	7	7	7
10	V-8	7	5	6	6	6	7	7	6
11	DC-1	8	6	6	6	7	7	7	6
12	PPG-AT 104	7	5	6	7	6	6	6	6
13	H10G-OP	7	6	6	7	6	7	8	7
14	Shark	7	6	6	7	6	7	7	6
15	Barracuda	7	6	7	7	6	6	7	6
16	PST-ORBS	7	6	6	7	7	8	7	7
17	PST-DCV6	7	6	6	7	7	6	7	6
	LSD 0.05	1	1	NS	1	1	1	1	1

¹Quality rated on a 1 to 9 scale with 9 = best quality, 1 = worst quality, and 6 = lowest acceptable quality.

²Color rated on a 1 to 9 scale with 9 = dark, 1 = light green.