

2014

Long-term fate of nitrate fertilizer in agricultural soils is not necessarily related to nitrate leaching from agricultural soils

Michael J. Castellano
Iowa State University, castelmj@iastate.edu

Mark B. David
University of Illinois at Urbana-Champaign

Follow this and additional works at: http://lib.dr.iastate.edu/agron_pubs

 Part of the [Agriculture Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/agron_pubs/204. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Agronomy at Iowa State University Digital Repository. It has been accepted for inclusion in Agronomy Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Long-term fate of nitrate fertilizer in agricultural soils is not necessarily related to nitrate leaching from agricultural soils

Abstract

Accounting for the fate of inorganic N fertilizer in agricultural systems is critical to sustainable production. Sebilo et al. (1) provide a unique long-term record of $^{15}\text{NO}_3$ fertilizer fate that demonstrates N molecules from a discrete fertilizer application are transferred to soil organic matter (SOM) and subsequently mineralized over the course of ca. 100 years during which they contribute to NO_3 leaching. The authors conclude “attempts to reduce agricultural nitrate contamination of aquatic systems must consider the long-term legacy of past applications of synthetic fertilizers”. Further, they suggest a recent decrease in anthropogenic N inputs to the Mississippi River Basin, without a concomitant decrease in riverine NO_3 loads, is consistent with their conclusion

Disciplines

Agriculture

Comments

This is an article from Castellano, Michael J., and Mark B. David. "Long-term fate of nitrate fertilizer in agricultural soils is not necessarily related to nitrate leaching from agricultural soils." *Proceedings of the National Academy of Sciences* 111, no. 8 (2014): E766-E766. doi: [10.1073/pnas.1321350111](https://doi.org/10.1073/pnas.1321350111). Posted with permission.

The long-term fate of nitrate fertilizer in agricultural soils is not necessarily related to nitrate leaching from agricultural soils

Michael J. Castellano*, Department of Agronomy, Iowa State University, Ames, Iowa 50011

Mark B. David, Department of Natural Resources and Environmental Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801

*corresponding author: castellanomichaelj@gmail.com

Accounting for the fate of inorganic N fertilizer in agricultural systems is critical to sustainable production. Sebito et al. (1) provide a unique long-term record of $^{15}\text{NO}_3$ fertilizer fate that demonstrates N molecules from a discrete fertilizer application are transferred to soil organic matter (SOM) and subsequently mineralized over the course of ca. 100 years during which they contribute to NO_3 leaching. The authors conclude “attempts to reduce agricultural nitrate contamination of aquatic systems must consider the long-term legacy of past applications of synthetic fertilizers”. Further, they suggest a recent decrease in anthropogenic N inputs to the Mississippi River Basin, without a concomitant decrease in riverine NO_3 loads, is consistent with their conclusion.

In contrast to the authors’ interpretation, we highlight that short-term changes in agricultural management can lead to large short-term reductions in NO_3 leaching. We attribute the authors’ results to long-term soil organic matter (SOM) turnover and associated NO_3 losses that occur in the absence of high biological N demand. We suggest that changes in agricultural management that produce rapid reductions in total NO_3 leaching would produce proportional reductions in NO_3 leaching derived from historical fertilizer applications.

Diversified cropping systems, conversions to perennial biofuels, and cover crops all provide direct evidence of the short-term potential for decreases in NO_3 leaching. In maize-alfalfa cropping systems, NO_3 leaching under alfalfa is 75-80% lower than the preceding maize crop (2). Perennial biofuel crops can reduce NO_3 leaching by >90% within four years of establishment (3). Non-legume cover crops provide on average a 70% reduction in NO_3 leaching during 2-3 year experiments (4).

^{15}N labeling studies demonstrate the proportion of fertilizer transferred to SOM is rapidly distributed among pools with fast and slow turnover rates (*e.g.*, 5). Consistent with ecosystem models, a small portion of the ^{15}N transferred to slow-turnover SOM-N pools will mineralize every year. In annual cropping systems where plant uptake is not fully synchronized with N mineralization, a portion of the mineralized N will nitrify and leach. As such, we expect strategies that reduce total NO_3 leaching will provide proportionally equal reductions in all

sources of NO₃ leaching independent of whether the NO₃ is derived from fast- or slow-turnover SOM-N pools.

We also question the suggestion that net anthropogenic N inputs (NANI) to the Mississippi River basin are decreasing. None of the cited papers demonstrate a decrease in NANI. And recent analyses show NANI to be relatively static over the last 20 years.

With a goal to reduce N losses to the Gulf of Mexico by 45%, we are concerned that readers will misinterpret the authors' data to suggest short-term agricultural management strategies cannot provide large, rapid reductions in NO₃ leaching. Strategies that minimize bare fallows and maximize plant N demand are well known to provide rapid reductions in NO₃ leaching (2,3,4). Although mineralization of historical N fertilizer applications cannot be prevented, the mineralized N can be recycled into SOM – particularly with strategies that aim to restore depleted soil organic carbon stocks.

1. Sebilo M, Mayer B, Nicolardot B, Pinay G, Mariotti A (2013) Long-term fate of nitrate fertilizer in agricultural soils. *Proceedings of the National Academy of Sciences of the United States of America* doi/10.1073/pnas.1305372110.
2. Toth JD, Fox RH (1998) Nitrate Losses from a Corn-Alfalfa Rotation: Lysimeter Measurement of Nitrate Leaching. *Journal of Environmental Quality* 27:1027–1033.
3. Smith CM et al. (2013) Reduced nitrogen losses after conversion of row crop agriculture to perennial biofuel crops. *Journal of Environmental Quality* 42:219–28.
4. Tonitto C, David M, Drinkwater L (2006) Replacing bare fallows with cover crops in fertilizer-intensive cropping systems: A meta-analysis of crop yield and N dynamics. *Agriculture, Ecosystems & Environment* 112:58–72.
5. Kaye J, Barrett J, Burke I (2002) Stable Nitrogen and Carbon Pools in Grassland Soils of Variable Texture and Carbon Content. *Ecosystems* 5:461–471.