

5-2015

Differential Improvements in Student Fruit and Vegetable Selection and Consumption in Response to the New National School Lunch Program Regulations: A Pilot Study

Karen W. Cullen

United States Department of Agriculture

Tzu-An Chen

United States Department of Agriculture

Jayna M. Dave

United States Department of Agriculture

Helen Jensen

Iowa State University, hhjensen@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/econ_las_pubs

 Part of the [Agricultural and Resource Economics Commons](#), [Agricultural Economics Commons](#), and the [Health Economics Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/econ_las_pubs/217. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Economics at Iowa State University Digital Repository. It has been accepted for inclusion in Economics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Differential Improvements in Student Fruit and Vegetable Selection and Consumption in Response to the New National School Lunch Program Regulations: A Pilot Study



Karen W. Cullen, DrPH; Tzu-An Chen, PhD; Jayna M. Dave, PhD; Helen Jensen, PhD

ARTICLE INFORMATION

Article history:

Accepted 23 September 2014
Available online 2 January 2015

Keywords:

School lunch regulations
Children
Lunch consumption
Fruit
Vegetables

2212-2672/Copyright © 2015 by the Academy of
Nutrition and Dietetics.
<http://dx.doi.org/10.1016/j.jand.2014.10.021>

ABSTRACT

Objective To investigate changes in student food selection and consumption in response to the new National School Lunch Program meal patterns during fall 2011.

Design Eight elementary and four intermediate schools in one Houston area school district were matched on free/reduced-price meal eligibility and randomized into control or intervention conditions.

Intervention Both intervention and control school cafeterias served the same menu. The intervention school cafeterias posted the new meal pattern daily; students could select one fruit and two vegetable servings per reimbursable meal. Control school students could only select the previous meal pattern: a total of two fruit and vegetable servings per meal.

Main outcome measures Students were observed during lunch: student sex and foods selected/consumed were recorded. Diet analysis software was used to calculate energy/food groups selected/consumed.

Statistical analyses performed Cochran-Mantel-Haenszel χ^2 tests examined differences in the percent of students selecting each meal component by condition, controlling for sex, grade, and school free/reduced-price meal eligibility. Analysis of covariance assessed differences in amount of energy/food groups selected and consumed, and differences in percent of food groups consumed.

Results Observations were conducted for 1,149 elementary and 427 intermediate students. Compared with students in the control schools, significantly more intervention elementary and intermediate school students selected total ($P<0.001$, $P<0.05$) and starchy vegetables ($P<0.001$, $P<0.01$); more intervention intermediate school students selected fruit ($P<0.001$), legumes ($P<0.05$), and protein foods ($P<0.01$). There were significantly greater amounts of these foods selected and consumed, but no differences in the proportion of the foods consumed by condition. Fewer calories were consumed by elementary and intermediate school intervention students.

Conclusions More intervention students selected fruit and vegetables at lunch and consumed them compared with control condition students. Future studies with larger and more diverse student populations are warranted.

J Acad Nutr Diet. 2015;115:743-750.

THE NATIONAL SCHOOL LUNCH PROGRAM (NSLP) sponsored by the US Department of Agriculture (USDA) served more than 31 million lunches each day during fiscal year 2012.¹ About 68% of these meals were provided to students at a free or reduced price (FRP).² In 2008, the USDA commissioned the Institute of Medicine to provide new meal pattern recommendations to align the federal school meal programs with the Dietary Guidelines for Americans and Dietary Reference Intakes to ensure that the meals promoted health and reduced inadequate and excessive intakes.³ The 2010 Healthy, Hunger-Free Kids Act directed USDA to update the school meal patterns and nutrition standards based on these recommendations.^{4,5}

The new meal patterns were implemented at the beginning of the 2012-2013 school year. For the first time, both minimum and maximum calorie limits were set to ensure age-appropriate meals for children in three grade groupings: 550 to 650 kcal for kindergarten through grade five, 600 to 700 kcal for grades 6 through 8, and 750 to 850 kcal for grades 9 through 12.⁴ The amount of fruit and vegetables (F/V) in the school menu pattern increased to align to the amounts in the Dietary Guidelines for Americans.⁴ For lunch, the new meal pattern provides a minimum of two servings (up to $\frac{1}{2}$ cup each) of vegetables and one serving ($\frac{1}{2}$ cup) fruit per lunch meal, one serving more than the previous standard of two servings total of F/V per lunch meal. A

specific number of servings of dark green and red/orange vegetables and legumes must be offered each week.⁴ For the offer vs serve option (OVS) in the new rules, students have to select at least one serving of a fruit or vegetable for the meal to count as reimbursable.⁴ OVS is optional for elementary schools.⁴ The expectation was that the opportunity to select more F/V at lunch would increase student F/V consumption. Other meal improvements included the specification that all grains be whole-grain rich (ie, must contain at least 50% whole grains and the remaining grain, if any, must be enriched) by July 2014, and a gradual reduction in the sodium content of the meals over 10 years.⁴

We present the results of a pilot study that investigated changes in student food selection and consumption in response to partial implementation of the new NSLP meal patterns for F/V during the fall 2011 semester. The main hypotheses were that intervention school students would select more servings of F/V, resulting in greater amounts selected and consumed compared with students in schools without access to the new F/V meal patterns.

METHODS

This pilot study was conducted during the fall semester of 2011, before the final NSLP meal patterns were published. Participants included intermediate and elementary school students in one school district in the Houston, TX, area. The school district had 37,000 students (21% Hispanic, 10% African American, 10% Asian, 59% white, and 26% FRP), 26 elementary (kindergarten through grade 5), and 10 intermediate schools (grades 6 through 8). The district received a monetary reimbursement to cover potential increases in food cost due to the menu changes. The child nutrition director selected the schools based on eligibility for FRP meals: four low (49% to 79% FRP) and four middle-income elementary schools (7% to 18% FRP), and two low (~34% FRP) and two middle-income (~20% FRP) intermediate schools. The schools were matched on grade level and FRP, and randomized to intervention or control conditions by the study statistician using a random numbers generator (Excel, Microsoft Corp). The planned study sample size (540 elementary and 540 intermediate observations) provided 80% power to detect significant differences with $\alpha=.05$.

The study was approved by the Institutional Review Board at Baylor College of Medicine. Because the study data were collected using anonymous observations of student food selection and consumption in the school cafeterias, consent forms were not required.

Menu Changes

The school district used a 2-week menu cycle; new menu patterns were developed that met the new guidelines.⁶ The menu included almost all whole grains, and only low-fat white milk or nonfat flavored milk. A fresh fruit was available every day, plus a raw vegetable, canned fruit, and cooked vegetable. Intervention and control schools served the same menu. The intervention school students were allowed to select one fruit serving and two vegetable servings (three total), plus a protein food, two grain servings, and a milk for the reimbursable meal. Control school students could only select the current USDA meal pattern of a total of two servings of fruit and/or vegetables, but similar servings of grain,

protein, and milk as in the intervention schools. The district allowed the OVS option at all grade levels; students could select fewer meal components as long as the meal met a specified minimum.⁶ Because this study took place before the nationwide implementation of the new guidelines, the district was unable to implement the new OVS rule that students had to select at least one fruit or vegetable serving for the meal to qualify as a reimbursable meal.

In the six intervention schools, letters in English and Spanish that explained the new menu pattern were sent home to all parents/guardians, and the teachers received information to display in their classrooms. Each intervention cafeteria set up an easel at the entrance to the serving line and displayed color photographs of the foods being served that day showing the correct number of servings to select under each food group category. There was also a small sign placed on the serving line that identified that one fruit and two vegetable servings could be selected for each reimbursable lunch meal. No easel or signage was present in the control schools.

Cafeteria Observation Procedures

Student consumption data were collected by direct observation in the cafeterias during lunch periods. All foods provided on the menus and sold as à la carte were preprinted on an observation checklist. There were columns to check the foods the student selected in the cafeteria line, and identify source (eg, using codes for NSLP, home, à la carte, or a friend), and whether food was given away, spilled, or obtained (eg, purchase or trade). Extra lines were available to record other foods and the source. For each item, the amount eaten was recorded using the quarter waste method (ie, 0, $\frac{1}{4}$, $\frac{1}{2}$, $\frac{3}{4}$, or all), which has high interrater and intermethod reliability.⁷ Student sex and grade level were also recorded.

Seven research staff members (four registered dietitian nutritionists, two staff with nutrition degrees, and one college undergraduate) attended a 3-hour training. Each observer conducted two to four practice observations, with the research coordinator also recording consumption. Interrater reliability was assessed and practice continued until acceptable (>90%) agreement was obtained. One trained observer conducted quality control checks once a month.

Observers were assigned to specific schools and visited each school 1 day each week during the semester. The schools did not know the day of the visit in advance. Each observer obtained the cafeteria seating arrangements and established the weekly data collection rotation for each school. Elementary school classrooms were assigned a table and this information was used so that equal numbers of students in each grade were observed during the semester. The intermediate schools did not have grade-specific lunch periods. Therefore, only intermediate grade level was recorded on the intermediate school observation sheets. The observer developed a rotation plan so that all tables in the schools would be observed in a systematic method during the semester. Because names were not obtained, a student could have been observed more than once.

There was a continuous influx of elementary school students into the cafeteria lunch line; the intermediate schools had three lunch periods each day. Each data collector first checked the cafeteria line lunch items against the observation

checklist and menu for the day. Then the observer selected three to four students with a reimbursable NSLP meal, defined as a meal containing at least three of the five lunch components, who approached and then sat at the tables to be observed that day. The observations of these selected students were conducted unobtrusively from a distance.

Nutrient and Food Group Analyses

For each student lunch observation form, the foods selected and consumed were entered into separate Nutrition Data System for Research (NDSR) files (version 2011, Nutrition Coordinating Center, University of Minnesota) by trained registered dietitian nutritionists. The project manager and research dietitian nutritionists obtained the recipes and nutrient information for menu items from the food service department, and NDSR recipes were created for each item. Student intake of calories and food groups (ie, fruit, 100% fruit juice, vegetables [total; dark green; red-orange; starchy such as white potatoes, corn, or peas; other such as green beans or celery; legumes; and high-fat vegetables], grains, protein foods, snack chips, sugar-sweetened beverages, desserts, and milk) were obtained with the NDSR food group system.^{8,9} It is important to note that unlike the USDA MyPyramid coding system, the NDSR food group system does not count the F/V contributions from high-fat, high-sugar foods.¹⁰

Statistical Methods

Separate analyses were conducted for elementary and intermediate school student data. To test for the difference in the percentage of students selecting each food component by condition, Cochran-Mantel-Haenszel χ^2 tests were used, controlling for student sex and school FRP (low or middle income) for both elementary and middle schools, and also controlling for grade for elementary schools. Next, separate analyses of covariance were conducted to compare the mean amounts of calories and food groups selected and consumed, and the percentage of each food group consumed, with intervention or control school membership as the between-group factor, adjusting for student sex and school FRP for elementary and intermediate school student data, and for grade for elementary school data. The adjusted outcome means for each food component for intervention and control groups were reported. An alpha level of $P < 0.05$ was used to determine statistical significance. All the analyses were performed using SAS (version 9.3, 2011, SAS Institute Inc).

RESULTS

Observations of student food selection and consumption were obtained for 1,149 elementary school students (535 intervention and 614 control) and 427 intermediate

Table 1. Percent of students selecting each food component in schools testing the new National School Lunch Program meal patterns for fruit and vegetables during fall 2011^a

Food component	Elementary School		Intermediate School	
	Intervention n=535	Control n=614	Intervention n=212	Control n=215
	←----- % ----->			
Fruit ^{b***}	56	51	45	21
Juice ^{c***}	50	61	52	43
Fruit and juice ^{b***c*}	82	87	78	58
Total vegetables ^{b*c***}	63	52	52	41
Dark green vegetables	10	8	11	14
Red orange vegetables	19	17	20	20
Starchy vegetables ^{b*c***}	47	36	39	27
Other vegetables	13	11	17	16
Legumes ^{d*}	4	2	9	4
Grains	100	100	100	99
Whole grains ^{c*}	40	48	49	52
Protein foods ^{b**}	100	100	100	97
Milk	90	91	76	74

^aCochran-Mantel-Haenszel test controlling for sex and school socioeconomic status (and grade for elementary school data).

^bSignificant difference between intermediate intervention and control school students.

^cSignificant difference between elementary intervention and control school students.

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

school students (212 intervention and 215 control). Significantly more intervention elementary school students selected total vegetables ($P<0.001$) and starchy vegetables ($P<0.001$), but significantly fewer selected juice ($P<0.001$) and whole grains ($P<0.05$), compared with control school students (Table 1).

More than twice as many intervention intermediate school students selected fruit (45%) compared with control school students (21%) ($P<0.001$). More intervention intermediate school students also selected total vegetables ($P<0.05$), starchy vegetables ($P<0.01$), legumes ($P<0.05$), and protein foods ($P<0.01$) (Table 1).

As selected, elementary intervention school student lunch meals contained significantly greater quantities of total vegetables ($P<0.001$), dark green vegetables ($P<0.01$), and starchy vegetables ($P<0.001$), but less juice ($P<0.001$), whole grains ($P<0.01$), and protein foods ($P<0.001$) compared with control school lunches (Table 2). The amounts actually consumed had similar findings except for dark green vegetables for which there was no significant difference in consumption between groups (Table 2). Elementary intervention students also consumed significantly more other vegetables than control students ($P<0.05$). Overall, intervention elementary school students selected and consumed

significantly less energy ($P<0.05$ for both) than control school students.

Compared with the lunches of the intermediate control school students, intermediate school intervention lunch meals as selected contained significantly greater amounts of fruit ($P<0.001$), juice ($P<0.05$), total vegetables ($P<0.01$), starchy vegetables ($P<0.001$), and legumes ($P<0.05$); and significantly more of these food groups were actually consumed (Table 3). However, intermediate school intervention students selected lunch meals with significantly fewer whole grains ($P<0.05$) and they consumed less total grains ($P<0.05$) and whole grains ($P<0.01$) than control school students. Overall, intervention intermediate school students consumed significantly less energy ($P<0.01$) than control school students.

Intervention elementary school students consumed a greater proportion of other vegetables selected ($P<0.05$) and a lower proportion ($P<0.01$) of protein foods selected than control elementary school students, but there were no significant differences between intervention and control students in the proportion of any of the other types of vegetables or fruit they consumed (Table 4).

Intervention intermediate school students consumed a significantly lower proportion ($P<0.001$) of calories, total

Table 2. Mean amounts of calories and foods selected and consumed at lunch by all elementary school students observed in the four intervention and four control schools testing the new National School Lunch Program meal patterns for fruit and vegetables during fall 2011^a

Food	Food Selected		Food Consumed	
	Intervention (n = 535)	Control (n = 614)	Intervention (n = 535)	Control (n = 614) ^b
	←————— <i>mean ± standard error</i> —————→			
Calories ^{c,d*}	598±5.00	614±4.66	449±6.65	469±6.21
Fruit (c)	0.29±0.01	0.26±0.01	0.20±0.01	0.18±0.01
Juice (c) ^{c***d***}	0.26±0.01	0.32±0.01	0.18±0.01	0.23±0.01
Fruit and juice (c) ^{c*}	0.54±0.01	0.59±0.01	0.38±0.01	0.41±0.01
Total vegetables (c) ^{c***d**}	0.32±0.01	0.22±0.01	0.14±0.01	0.10±0.01
Dark green vegetables ^{c**}	0.06±0.01	0.03±0.01	0.01±0.00	0.00±0.00
Red orange vegetables	0.07±0.01	0.06±0.01	0.03±0.00	0.03±0.00
Starchy vegetables ^{c***d**}	0.15±0.01	0.10±0.01	0.10±0.01	0.06±0.01
Other vegetables ^{d*}	0.02±0.00	0.02±0.00	0.01±0.00	0.00±0.00
Legumes	0.02±0.00	0.01±0.00	0.01±0.00	0.00±0.00
Grains (oz equivalent)	2.21±0.04	2.21±0.04	1.62±0.04	1.65±0.04
Whole grains ^{c**d*}	0.48±0.03	0.59±0.03	0.30±0.02	0.39±0.02
Protein foods (oz equivalents) ^{c***d***}	2.00±0.04	2.27±0.04	1.43±0.05	1.77±0.04
Milk (oz)	7.11±0.11	7.28±0.10	5.38±0.15	5.19±0.14

^aAnalyses of covariance, adjusting for sex and school socioeconomic status (and grade for elementary school data).

^bOne control school student did not eat the meal.

^cSignificant food selected difference between intervention and control elementary school students.

^dSignificant food consumed difference between intervention and control elementary school students.

* $P<0.05$.

** $P<0.01$.

*** $P<0.001$.

Table 3. Mean amounts of calories and foods selected and consumed at lunch by all intermediate school students observed in the two intervention and two control schools testing the new National School Lunch Program meal patterns for fruit and vegetables during fall 2011^a

Food	Food Selected		Food Consumed	
	Intervention (n=212)	Control (n=215)	Intervention (n=210) ^b	Control (n=214) ^b
	←—mean±standard error—→			
Calories ^{c**}	612±10.8	599±10.7	520±12.4	571±12.5
Fruit (c) ^{c***d***}	0.25±0.02	0.12±0.02	0.19±0.02	0.09±0.02
Juice (c) ^{d*}	0.27±0.02	0.22±0.02	0.21±0.02	0.17±0.02
Fruit and juice (c) ^{c***d***}	0.52±0.02	0.33±0.02	0.39±0.02	0.26±0.02
Total vegetables (c) ^{c***d***}	0.23±0.02	0.14±0.02	0.17±0.02	0.10±0.02
Dark green vegetables	0.02±0.01	0.03±0.01	0.01±0.00	0.01±0.00
Red orange vegetables	0.05±0.01	0.04±0.01	0.03±0.01	0.03±0.01
Starchy vegetables ^{c*d***}	0.12±0.01	0.06±0.01	0.10±0.01	0.06±0.01
Other vegetables	0.05±0.01	0.05±0.01	0.03±0.01	0.02±0.01
Legumes ^{c***d*}	0.03±0.01	0.01±0.01	0.03±0.01	0.01±0.01
Grains (oz equivalents) ^{c*}	2.56±0.09	2.65±0.09	2.13±0.09	2.43±0.09
Whole grains ^{c***d*}	0.55±0.05	0.70±0.05	0.37±0.04	0.56±0.04
Protein foods (oz equivalents)	2.15±0.06	2.18±0.06	1.84±0.08	2.05±0.08
Milk (oz)	5.89±0.25	5.64±0.25	4.68±0.27	5.07±0.26

^aAnalyses of covariance, adjusting for sex and school socioeconomic status (and grade for elementary school data).

^bTwo intervention and one control school students did not eat their selected meals.

^cSignificant food consumed difference between intervention and control intermediate school students.

^dSignificant food selected difference between intervention and control intermediate school students.

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

grains ($P < 0.001$), whole grains ($P < 0.05$), and protein foods ($P < 0.01$) than control school students (Table 4). There were no significant differences in the proportion of fruit or vegetables consumed by intermediate intervention or control school students (Table 4).

DISCUSSION

Our pilot study investigated changes in student food selection and consumption in response to the new NSLP guidelines for F/V. Compared with elementary students in control schools, significantly more elementary intervention students selected total vegetables, dark green, and starchy vegetables, resulting in significantly greater amounts of these foods on their trays and significantly greater amounts of total and starchy vegetables consumed. However, significantly fewer intervention elementary school students selected juice, whole grains, and protein foods, which led to significantly smaller amounts of these foods on their trays and actually consumed. Elementary school intervention students also selected and consumed significantly fewer calories for lunch than control school elementary students.

The results for the intermediate school intervention students were also encouraging; significantly more intermediate

school intervention students selected fruit, juice, legumes, and protein foods. The intermediate intervention school students, therefore, had significantly greater amounts of fruit, juice, legumes, and total and starchy vegetables on their trays, and consumed significantly more fruit, legumes, and total and starchy vegetables than intermediate control school students. The intermediate intervention school students also selected and consumed significantly lower amounts of whole grains and total grains and consumed significantly fewer calories than control school students.

There were no significant differences in the proportion of F/V consumed and, therefore, wasted, between the intervention and control condition students except that elementary school intervention students consumed a greater proportion of other vegetables than control school students. However, because more students were selecting F/V, total waste was greater.

The new menu patterns were implemented nationwide during fall 2012. A recent study¹¹ examined food selection and consumption before and after the new school guidelines were implemented and found a significant increase in the proportion of elementary and intermediate school students selecting fruit, but no difference in the proportion selecting

Table 4. Percent of food consumed by all observed students in schools testing the new National School Lunch Program meal patterns for fruit and vegetables during fall 2011^a

Food	Elementary School		Intermediate School	
	Intervention (n=535)	Control (n=613)	Intervention (n=210)	Control (n=214)
	←—————mean±standard error—————→			
Calories ^{b***}	75±1	77±1	85±2	97±2
Fruit	67±2	64±2	76±4	78±6
Juice	70±2	70±2	75±3	78±4
Fruit and juice	69±2	67±2	76±3	79±3
Total vegetables	40±17	63±18	53±5	52±6
Dark green vegetables	13±4	23±5	67±10	55±9
Red orange vegetables	38±4	43±4	40±8	40±9
Starchy vegetables	39±3	37±3	52±6	43±7
Other vegetables ^{c*}	28±4	15±5	29±8	30±9
Legumes	46±9	40±11	66±12	41±20
Grains ^{b***}	73±2	73±1	83±4	100±4
Whole grains ^{b*}	64±3	65±3	67±4	81±4
Protein foods ^{b**c**}	72±2	79±1	84±3	96±3
Milk	81±5	75±5	79±10	103±10

^aAnalyses of covariance, adjusting for sex and school socioeconomic status (and grade for elementary school data).

^bSignificant % consumption difference between intermediate intervention and control school students.

^cSignificant % consumption difference between elementary intervention and control school students.

* $P < 0.05$.

** $P < 0.01$.

*** $P < 0.001$.

vegetables. However, a greater proportion and amount of the vegetables were consumed, but there was no difference for fruit consumption.¹¹ These findings, plus the results from our study, are encouraging. Regular monitoring of student food selection and consumption at school and over 24 hours is needed to assess the influence of the new meal patterns on diet.

Fruit and vegetable availability on the school cafeteria serving line has been related to improved student F/V consumption. Using national data from 2005, Newman¹² found that more students in schools that met the 2012 NSLP F/V daily standards tried vegetables and ate significantly more total vegetables than students in schools where menus did not meet the daily standard (0.38 cup vs 0.30 cup). An environmental cafeteria intervention in elementary schools that included an extra fruit or vegetable in the lunch line daily, attractive food presentations, and verbal prompts from cafeteria staff resulted in a significant increase in fruit consumption.¹³ Other research has investigated cafeteria design and food presentation to influence student food selection behaviors.^{14,15} Longer-term studies are needed to identify successful strategies to improve student F/V selection and consumption.

Another reason for the improvement in fruit (intermediate) and vegetable (elementary and intermediate) selection and consumption may be the increased promotion and marketing

of the new menu pattern. There was signage about the new pattern in the cafeteria, and classroom teachers and parents received supporting materials. Whether students were aware of and responded to these components was not assessed in this study. This is an important area of research. For example, the use of attractive names for carrots on the serving line led to significantly higher consumption of carrots, and to a significantly higher proportion of vegetables selected in a previous study.¹⁶

The new NSLP meal guidelines also set minimum and maximum calorie levels for lunch meals. In this study, elementary school students selected meals with caloric content within the guidelines, although intervention school students selected significantly fewer calories than control school students (598 vs 614 kcal). However, actual mean consumption for both groups was lower than the guidelines, with intervention school students consuming significantly fewer calories than control school students (449 vs 469 kcal). Intermediate school students selected meals that met the new minimum level for NSLP meals (612 kcal for intervention and 599 kcal for control), but mean consumption was also lower than the guidelines and intervention school students consumed significantly fewer calories than control school students (520 vs 571 kcal). These values are lower than what has been found in previous research. For example, mean lunch intakes for elementary and middle school students

were 587 and 620 kcal, respectively, for students in a national study who completed 24-hour recalls during the 2004-2005 school year.¹⁷ In another study, middle school students who completed a lunch food record in the cafeteria during lunch reported a mean intake of 694 kcal.¹⁸ The lower values in our study may be the result of the more objective method of obtaining dietary intake (observation by trained staff) in contrast to the self-report method used in the previous studies that required portion size estimation by students. A Colorado study¹⁹ using an objective method of assessment—digital photography of the foods selected and left at the end of the meal—reported results that were similar to our study (elementary [426 kcal] and middle school [529 kcal] students). Whether student energy needs are met by school meals is an important area for future research. This is particularly significant for those children for whom the school lunch meals are an important safety net for meeting food needs.

The generally low consumption of fruit, vegetables, and whole grains by students is a concern and should prompt future research efforts associated with improving student preferences, selection, and consumption. Previous school studies have documented fruit waste of 45%,¹¹ 47%,²⁰ and 33% to 50%¹⁹; and vegetable waste of 40% to 90%.^{11,19,20} These rates are similar to those in our study: fruit waste of 33% to 36% for elementary intervention and control school students, respectively, and 24% to 22% for intermediate intervention and control school students, respectively; and vegetable waste of 60% to 37% for elementary intervention and control school students, respectively, and 47% to 48% for intermediate intervention and control school students, respectively (Table 4). Previous research indicates that targeted nutrition education may be a promising strategy for increasing F/V consumption. For example, one previous study found that nutrition classes were needed to enable elementary school students to select more fruit or vegetables when a salad bar was introduced.²¹ In the Institute of Medicine report,³ support for school foodservice staff was recommended. This support could include marketing and presentation tips, as well as training for the foodservice staff to offer positive encouragement about the foods during meal service. Some previous studies have noted improvements in student food selection and consumption with positive encouragement to students during NSLP meal selection.^{13,22,23} Other promising strategies from recent studies included taste testing for parents and students, marketing on the food line, and media campaigns.^{3,24-26}

There are several limitations to this study that should be noted. The participating school district only had 26% of students eligible for FRP meals; about 62% of Texas students were eligible for FRP meals during that school year.²⁷ Thus, the findings from this district might not generalize to others in Texas and the United States. Plus, the study was conducted in 12 schools in the Houston area, also limiting generalizability. The requirement to select at least one fruit or vegetable serving for a reimbursable meal was not implemented, which should prompt future studies to assess the influence of this piece of the new rule on student lunch consumption. In addition, the promotion and signage was only available in the intervention schools; perhaps the changes found for the intervention students were due to these strategies. Because this was a pilot study, with exploratory hypotheses on consumption, adjustments for multiple comparisons were not made.

CONCLUSIONS

The key finding from this study is that in response to piloting the new NSLP meal patterns for F/V, whereby students could select three servings of F/V per reimbursable meal, more total and starchy vegetables and fruit were consumed by intervention school students compared with students in the control schools. The proportion of the F/V consumed did not differ between the intervention and control school students. This is an important step in the right direction. Future research should continue to evaluate the effect of the new menu patterns on student food selection and consumption at school.

References

1. US Department of Agriculture. National School Lunch Program fact sheet. <http://www.fns.usda.gov/sites/default/files/NSLPFactSheet.pdf>. Published 2012. Updated September, 2013. Accessed October 29, 2013.
2. US Department of Agriculture. National School Lunch Program: Participation and meals served. <http://www.fns.usda.gov/pdf/slsummar.htm>. Published 2013. Updated October 17, 2013. Accessed October 29, 2013.
3. Stallings VA, Suitor CW, Taylor CL, eds. *School Meals: Building Blocks for Healthy Children*. Washington, DC: National Academies Press; 2009.
4. Nutrition Standards in the National School Lunch and School Breakfast Programs; Final Rule. 77 *Federal Register* 4088-4167 (2012).
5. US Department of Agriculture. Healthy, Hunger-Free Kids Act of 2010. http://www.fns.usda.gov/cnd/Governance/Legislation/CNR_2_010.htm. Published 2010. Updated August 2, 2013. Accessed August 27, 2013.
6. US Department of Agriculture. Menu planning in the National School Lunch Program. http://www.fns.usda.gov/cnd/menu/menu_planning.doc. Published September 1, 2000. Accessed October 29, 2013.
7. Hanks AS, Wansink B, Just DR. Reliability and accuracy of real-time visualization techniques for measuring school cafeteria tray waste: Validating the quarter-waste method. *J Acad Nutr Diet*. 2014;114(3):470-474.
8. Cullen KW, Himes JH, Baranowski T, et al. Validity and reliability of a behavior-based food coding system for measuring fruit, 100% fruit juice, vegetable, and sweetened beverage consumption: Results from the Girls Health Enrichment Multisite Studies. *Prev Med*. 2004;38(suppl):S24-S33.
9. Nutrition Coordinating Center. *NDSR 2013 User Manual*. Minneapolis, MN: University of Minnesota; 2013.
10. Cotton PA, Subar AF, Friday JE, Cook A. Dietary sources of nutrients among US adults, 1994 to 1996. *J Am Diet Assoc*. 2004;104(6):921-930.
11. Cohen JF, Richardson S, Parker E, Catalano PJ, Rimm EB. Impact of the new U.S. Department of Agriculture school meal standards on food selection, consumption, and waste. *Am J Prev Med*. 2014;46(4):388-394.
12. Newman C. Fruit and vegetable consumption by school lunch participants: Implications for the success of new nutrition standards, ERR-154. <http://www.ers.usda.gov/publications/err-economic-research-report/err154.aspx>. Published August, 2013. Updated August 27, 2013. Accessed September 3, 2013.
13. Perry CL, Bishop DB, Taylor GL, et al. A randomized school trial of environmental strategies to encourage fruit and vegetable consumption among children. *Health Educ Behav*. 2004;31(1):65-76.
14. Hanks AS, Just DR, Wansink B. Smarter lunchrooms can address new school lunchroom guidelines and childhood obesity. *J Pediatr*. 2013;162(4):867-869.
15. Wansink B, Just DR, Hanks AS, Smith LE. Pre-sliced fruit in school cafeterias: Children's selection and intake. *Am J Prev Med*. 2013;44(5):477-480.
16. Wansink B, Just DR, Payne CR, Klinger MZ. Attractive names sustain increased vegetable intake in schools. *Prev Med*. 2012;55(4):330-332.

17. Gordon A, Fox MK, Clark M, et al. *School Nutrition Dietary Assessment Study III: Volume II: Student Participation and Dietary Intakes—Final Report*. Princeton, NJ: Mathematica Policy Research; 2007.
18. Cullen KW, Watson KB, Dave JM. Middle-school students' school lunch consumption does not meet the new Institute of Medicine's National School Lunch Program recommendations. *Public Health Nutr.* 2011;14(10):1876-1881.
19. Smith SL, Cunningham-Sabo L. Food choice, plate waste and nutrient intake of elementary- and middle-school students participating in the US National School Lunch Program. *Public Health Nutr.* 2014;17(6):1255-1263.
20. Cohen JF, Richardson S, Austin SB, Economos CD, Rimm EB. School lunch waste among middle school students: Nutrients consumed and costs. *Am J Prev Med.* 2013;44(2):114-121.
21. Suarez-Balcazar Y, Redmond L, Kouba J, et al. Introducing systems change in the schools: The case of school luncheons and vending machines. *Am J Community Psychol.* 2007;39(3-4):335-345.
22. Hendy HM. Comparison of five teacher actions to encourage children's new food acceptance. *Ann Behav Med.* 1999;21(1):20-26.
23. Schwartz MB. The influence of a verbal prompt on school lunch fruit consumption: A pilot study. *Int J Behav Nutr Phys Act.* 2007;4:6.
24. Fulkerson JA, French SA, Story M, Nelson H, Hannan PJ. Promotions to increase lower-fat food choices among students in secondary schools: Description and outcomes of TACOS (Trying Alternative Cafeteria Options in Schools). *Public Health Nutr.* 2004;7(5):665-674.
25. Wechsler H, Basch CE, Zybert P, Shea S. Promoting the selection of low-fat milk in elementary school cafeterias in an inner-city Latino community: Evaluation of an intervention. *Am J Public Health.* 1998;88(3):427-433.
26. US Department of Agriculture. Team nutrition. <http://www.fns.usda.gov/tn/team-nutrition>. Published 2013. Updated March 19, 2014. Accessed March 31, 2014.
27. Center for Public Policy Priorities. State of Texas Children 2012 -Texas KIDS COUNT annual data book. http://www.cppp.org/sotc/pdf/TKC2012_Final_1-page_view.pdf. Published April 5, 2012. Accessed October 29, 2013.

AUTHOR INFORMATION

K. W. Cullen is a professor, T.-A. Chen is a biostatistician, and J. M. Dave is an assistant professor, US Department of Agriculture/Agriculture Research Service Children's Nutrition Research Center, Baylor College of Medicine, Houston, TX. H. Jensen is a professor, Department of Economics, Iowa State University, Ames.

Address correspondence to: Karen W. Cullen, DrPH, 1100 Bates St, Houston, TX 77030-2600. E-mail: kcullen@bcm.edu

STATEMENT OF POTENTIAL CONFLICT OF INTEREST

The authors have no potential conflict of interest to report.

FUNDING/SUPPORT

This study was supported by funding from the National Institutes of Health to Karen W. Cullen (no. R01HD068349). The project was funded in part by federal funds from the USDA/Agricultural Research Service under cooperative agreement no. 6250-51000-053.