

5-7-2008

A New Scorpionate Ligand: Tris(4,4-dimethyl-2-oxazoliny)borate and Its Zirconium(IV) Complexes

James Francis Dunne
Iowa State University, jfdunne@iastate.edu

Jiachun Su
Iowa State University, jcsu@iastate.edu

Arkady Ellern
Iowa State University, ellern@iastate.edu

Aaron D. Sadow
Iowa State University, sadow@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/chem_pubs

 Part of the [Chemistry Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/chem_pubs/226. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Chemistry at Iowa State University Digital Repository. It has been accepted for inclusion in Chemistry Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

A New Scorpionate Ligand: Tris(4,4-dimethyl-2-oxazoliny)borate and Its Zirconium(IV) Complexes

Abstract

The first example of a new class of oxazoline-based scorpionate ligand, tris(4,4-dimethyl-2-oxazoliny)phenyl borate, [ToM]⁻, is prepared by reaction of 2-lithio-4,4-dimethyl-2-oxazolide and 0.3 equiv of dichlorophenylborane. The steric bulk of this ligand is greater than that of tris(3,5-Me₂-pyrazoly)borate (Tp^{*}), as quantified by comparison of solid angles of crystallographically characterized zirconium(IV) complexes.

Keywords

Oxazoline (Oxa), scorpionate ligand, chemical modification, ligands, nitrogen compounds, zirconium

Disciplines

Chemistry

Comments

Reprinted (adapted) with permission from *Organometallics* 27 (2008): 2399, doi:[10.1021/om800252p](https://doi.org/10.1021/om800252p). Copyright 2008 American Chemical Society.

A New Scorpionate Ligand: Tris(4,4-dimethyl-2-oxazoliny)borate and Its Zirconium(IV) Complexes

James F. Dunne, Jiachun Su, Arkady Ellern, and Aaron D. Sadow*

Department of Chemistry, U.S. DOE Ames Laboratory, Iowa State University, Ames, Iowa 50011-3111


Received March 18, 2008

Summary: The first example of a new class of oxazoline-based scorpionate ligand, tris(4,4-dimethyl-2-oxazoliny)phenyl borate, $[\text{To}^{\text{M}}]^-$, is prepared by reaction of 2-lithio-4,4-dimethyl-2-oxazolide and 0.3 equiv of dichlorophenylborane. The steric bulk of this ligand is greater than that of tris(3,5-Me₂-pyrazolyl)borate (Tp^*), as quantified by comparison of solid angles of crystallographically characterized zirconium(IV) complexes.

Sterically encumbered tris(pyrazolyl)borate ligands, such as Tp^* (HB(3,5-Me₂-pyrazolyl)₃), Tp^{iBuMe} (HB(3-*i*Bu-5-Me-pyrazolyl)₃), and Tp^{Mes} (HB(3-mesitylpyrazolyl)₃), have recently captured attention as useful ancillary ligands in early transition metal chemistry.^{1–4} Early transition metal complexes of these bulky ligands as well as those of parent tris(pyrazolyl)borate (Tp) show interesting stoichiometric chemistry^{2–8} and display catalytic activity in reactions such as olefin polymerization.^{2,4,8a,9,10} However, unsymmetrically substituted Tp derivatives (e.g., 3-mesitylpyrazolyl) are susceptible toward isomerization processes, giving mixtures of 3- and 5-substituted tris(pyrazolyl)bo-

rate complexes.⁴ In highly reactive complexes such as $[\text{Zr}(\kappa^3\text{-Tp}^*)(\text{CH}_2\text{C}_6\text{H}_5)_2]^+$, ligand decomposition through B–N cleavage hinders possible utility in catalysis.² In fact, isomerization via 1,2-sigmatropic shifts and B–N bond cleavage are common reaction pathways for pyrazolylborate ligands in a range of transition metal complexes.^{1c,d,3,11}

To avoid this decomposition mechanism, we designed the tris(oxazoliny)borate scorpionate ligand that contains B–C linkages, proposed to have greater resistance to cleavage and isomerization processes.¹² The synthesis of tris(4,4-dimethyl-2-oxazoliny)phenyl borate, $[\text{To}^{\text{M}}]^-$, the first example of this new ligand class, and its coordination chemistry in new zirconium (IV) complexes are reported here. Lithium tris(4,4-dimethyl-2-oxazolin-2-yl)phenyl borate ($\text{Li}[\text{To}^{\text{M}}]$) is prepared by deprotonation of 4,4-dimethyl-2-oxazoline with *n*-BuLi (1.05 equiv) at –78 °C followed by addition of 0.30 equiv of dichlorophenylborane (eq 1).



Quantitative deprotonation of the oxazoline is necessary, as mixtures of 2*H*-oxazoline, 2-lithio-oxazolide, and dichlorophenylborane produce $\text{Li}[\text{To}^{\text{M}}]$ and several unidentified products. Independent experiments show that interaction of 4,4-dimethyl-2-oxazoline and PhBCl_2 in THF accounts for two of the side products. Also, a slight excess of 2-lithio-oxazolide (0.30 equiv of PhBCl_2) is necessary, as 0.33 equiv of PhBCl_2 and shorter reaction times (<26 h) afford inseparable mixtures of $\text{Li}[\text{To}^{\text{M}}]$ and a species assigned as bis(4,4-dimethyl-2-oxazolin-2-yl)phenylborane on the basis of integration of its ¹H NMR spectrum and its ¹¹B NMR chemical shift (–7.26 ppm).

* Corresponding author. E-mail: sadow@iastate.edu.

(1) (a) Calabrese, J. C.; Trofimenko, S.; Thompson, J. S. *J. Chem. Soc., Chem. Commun.* **1986**, 1122–1123. (b) Trofimenko, S.; Calabrese, J. C.; Thompson, J. S. *Inorg. Chem.* **1987**, *26*, 1507–1514. (c) Trofimenko, S.; Calabrese, J. C.; Domaille, P. J.; Thompson, J. S. *Inorg. Chem.* **1989**, *28*, 1091–1101. (d) Trofimenko, S. *Chem. Rev.* **1993**, *93*, 943. (e) Trofimenko, S. *Scorpionates—The Coordination Chemistry of Polypyrazolylborate Ligands*; Imperial College Press: London, 1999.

(2) Lee, H.; Jordan, R. F. *J. Am. Chem. Soc.* **2005**, *127*, 9384–9385.

(3) (a) Kersten, J. L.; Kucharczyk, R. R.; Yap, G. P. A.; Rheingold, A. L.; Theopold, K. H. *Chem.—Eur. J.* **1997**, *3*, 1668–1674. (b) Hess, A.; Horz, M. R.; Liable-Sands, L. M.; Lindner, D. C.; Rheingold, A. L.; Theopold, K. H. *Angew. Chem., Int. Ed.* **1999**, *38*, 166–168. (c) Qin, K.; Incarvito, C. D.; Rheingold, A. L.; Theopold, K. H. *Angew. Chem., Int. Ed.* **2002**, *41*, 2333–2335. (d) Qin, K.; Incarvito, C. D.; Rheingold, A. L.; Theopold, K. H. *J. Am. Chem. Soc.* **2002**, *124*, 14008–14009.

(4) (a) Murtuza, S.; Casagrande, O. L.; Jordan, R. F. *Organometallics* **2002**, *21*, 1882–1890. (b) Michiue, K.; Jordan, R. F. *Organometallics* **2004**, *23*, 460–470.

(5) (a) Reger, D. L.; Tarquini, M. E. *Inorg. Chem.* **1982**, *21*, 840–842. (b) Reger, D. L.; Tarquini, M. E. *Inorg. Chem.* **1983**, *22*, 1064–1068. (c) Reger, D. L.; Tarquini, M. E.; Lebioda, L. *Organometallics* **1983**, *2*, 1763–1769.

(6) (a) Kresinski, R. A.; Hamor, T. A.; Jones, C. J.; McCleverty, J. A. *J. Chem. Soc., Dalton Trans.* **1991**, 603–608. (b) Kresinski, R. A.; Isam, L.; Hamor, T. A.; Jones, C. J.; McCleverty, J. A. *J. Chem. Soc., Dalton Trans.* **1991**, 1835–1842.

(7) Antiñolo, A.; Carrillo-Hermosilla, F.; Corrochano, A. E.; Fernández-Baeza, J.; Lanfranchi, M.; Otera, A.; Pellinghelli, M. A. *J. Organomet. Chem.* **1999**, *577*, 174–180.

(8) (a) Long, D. P.; Bianconi, P. A. *J. Am. Chem. Soc.* **1996**, *118*, 12453–12454. (b) Blackwell, J.; Lehr, C.; Sun, Y.; Piers, W. E.; Pearce-Batchelder, S. D.; Zaworotko, M. J.; Young, V. G. *Can. J. Chem.* **1997**, *75*, 701–711. (c) Long, D. P.; Chandrasekaran, A.; Day, R. O.; Bianconi, P. A.; Rheingold, A. L. *Inorg. Chem.* **2000**, *39*, 4476–4487.

(9) Nakazawa, H.; Ikai, S.; Imaoka, K.; Kai, Y.; Yano, T. *J. Mol. Catal. A* **1998**, *132*, 33–41.

(10) (a) Furlan, L. G.; Gil, M. P.; Casagrande, O. L., Jr. *Macromol. Rapid Commun.* **2000**, *21*, 1054–1057. (b) Gil, M. P.; Casagrande, O. L., Jr. *J. Organomet. Chem.* **2004**, *689*, 286–292.


(11) (a) Albinati, A.; Bovens, M.; Rügger, H.; Venanzi, L. M. *Inorg. Chem.* **1997**, *36*, 5991–5999. (b) Brunker, T. J.; Hascall, T.; Cowley, A. R.; Rees, L. H.; O'Hare, D. *Inorg. Chem.* **2001**, *40*, 3170–3176. (c) Kückmann, T. I.; Abram, U. *Z. Anorg. Allg. Chem.* **2004**, *630*, 783–785. (d) Biagini, P.; Calderazzo, F.; Marchetti, F.; Romano, A. M.; Spera, S. *J. Organomet. Chem.* **2006**, *691*, 4172–4180. (e) Zhao, N.; Van Stipdonk, M. J.; Bauer, C.; Campana, C.; Eichhorn, D. M. *Inorg. Chem.* **2007**, *46*, 8662–8667.

(12) Although metal-mediated B–C bond cleavages are known, these reactions are less facile than B–N bond cleavage. For representative examples, see: (a) Bianchini, C.; Meli, A.; Laschi, F.; Vizza, F.; Zanello, P. *Inorg. Chem.* **1989**, *28*, 227–233. (b) Konze, W. V.; Scott, B. L.; Kubas, G. J. *J. Chem. Soc., Chem. Commun.* **1999**, 1807–1808. (c) Strunkina, L. I.; Minacheva, M. Kh.; Lyssenko, K. A.; Petrovskii, P. V.; Burlakov, V. V.; Rosenthal, U.; Shur, V. B. *Russ. Chem. Bull., Int. Ed.* **2006**, *55*, 174–176. (d) Cho, J.; Yap, G. P. A.; Riordan, C. G. *Inorg. Chem.* **2007**, *36*, 11308–11315.

The ^1H NMR spectra of $\text{Li}[\text{To}^{\text{M}}]$ dissolved in benzene- d_6 , chloroform- d_1 , or methylene chloride- d_2 contained broad resonances at 1.4 and 3.3 ppm as well as aryl resonances. In contrast, the 4,4-dimethyl and methylene groups (at 1.19 and 3.60 ppm) on equivalent oxazoline rings gave sharp singlet resonances in acetonitrile- d_3 . The formation of the borate center is supported by the integrated ratio of oxazoline methyl to phenyl resonances (18:5 H) as well as the broad singlet observed at -16.9 ppm in the ^{11}B NMR spectrum.¹³

This salt elimination route to $\text{Li}[\text{To}^{\text{M}}]$ is related to the synthesis of bis(oxazolinyldiphenylborate ligands (borabox).¹⁴ Borabox, which also is prepared via reaction of 2-lithium oxazolidine and a chloroborane electrophile, forms in 4–12 h. In contrast to this short reaction time, preparation of the neutral ligand 1,1,1-tris(oxazolinyldiphenylborate) (tris-ox) requires heating THF solutions of lithium bis(oxazoline) and 2-bromo-2-oxazoline at reflux for 5 days.¹⁵ Interestingly, the reaction conditions needed for synthesis of $\text{Li}[\text{To}^{\text{M}}]$ fall between these two ligands.

The acidic pro-ligand $\text{H}[\text{To}^{\text{M}}]$ is prepared by flash chromatography of $\text{Li}[\text{To}^{\text{M}}]$ on silica gel in $\text{C}_6\text{H}_{14}/i\text{-PrOH}/\text{NEt}_3$ (15:1:1), giving the product as an off-white solid (eq 2).¹⁴



The ^1H NMR spectrum of $\text{H}[\text{To}^{\text{M}}]$ (benzene- d_6) contained sharp singlet resonances at 1.02 (Me) and 3.55 ppm (CH_2), in contrast to the broad resonances observed for $\text{Li}[\text{To}^{\text{M}}]$ in this solvent. In acetonitrile- d_3 , the oxazoline peaks are shifted downfield by 0.1 and 0.3 ppm (to 1.29 and 3.90 ppm) in comparison to $\text{Li}[\text{To}^{\text{M}}]$. Crystals of $\text{H}[\text{To}^{\text{M}}]$ are obtained from the chromatography solvent mixture after slow evaporation. Although the crystals diffracted weakly, single-crystal X-ray diffraction confirms the constitution and connectivity of $[\text{To}^{\text{M}}]^-$, in particular the formation of a tetrahedral borate center containing four B–C bonds (Figure 1).

The symmetry of $[\text{To}^{\text{M}}]^-$ in the solid state is C_s , as two of the oxazolines' C=N are coplanar ($\text{N1}=\text{C7}-\text{C17}=\text{N3}$ dihedral = 0°), oriented toward each other as if to form a bidentate chelate, and related by a mirror plane. That mirror plane contains the phenyl group and the third oxazoline. However, the ^1H NMR spectrum of $\text{H}[\text{To}^{\text{M}}]$ (benzene- d_6 or acetonitrile- d_3) indicates that all three oxazoline groups are equivalent in solution.

We have prepared $[\text{To}^{\text{M}}]\text{Zr}(\text{IV})$ complexes through salt metathesis with $\text{Li}[\text{To}^{\text{M}}]$ and ZrCl_4 as well as via amine elimination with $\text{H}[\text{To}^{\text{M}}]$ and $\text{Zr}(\text{NMe}_2)_4$. A salt elimination reaction occurs when a mixture of $\text{Li}[\text{To}^{\text{M}}]$ and 1.05 equiv of ZrCl_4 suspended in methylene chloride is stirred for 4 days,

(13) Kidd, R. G. In *NMR of Newly Accessible Nuclei*; Laszlo, P., Ed.; Academic Press: New York, 1983; Vol. 2, pp 49–77.

(14) (a) Mazet, C.; Kohler, V.; Pfaltz, A. *Angew. Chem., Int. Ed.* **2005**, *44*, 4888–4891. (b) Mazet, C.; Roseblade, S.; Kohler, V.; Pfaltz, A. *Org. Lett.* **2006**, *8*, 1879–1882.

(15) (a) Bellemin-Laponnaz, S.; Gade, L. H. *J. Chem. Soc., Chem. Commun.* **2002**, 1286–1287. (b) Bellemin-Laponnaz, S.; Gade, L. H. *Angew. Chem., Int. Ed.* **2002**, *41*, 3473–3475. (c) Ward, B. D.; Bellemin-Laponnaz, S.; Gade, L. H. *Angew. Chem., Int. Ed.* **2005**, *44*, 1668–1671. (d) Lukesova, L.; Ward, B. D.; Bellemin-Laponnaz, S.; Wadepohl, H.; Gade, L. H. *J. Chem. Soc., Dalton Trans.* **2007**, 920–922.






Figure 1. ORTEP diagram of $\text{H}[\text{To}^{\text{M}}]$ at 50% probability. Hydrogen atoms are omitted for clarity.

giving a yellow solution of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ and suspended lithium chloride (eq 3).



The same compound is also obtained by heating a mixture of $\text{Li}[\text{To}^{\text{M}}]$ and ZrCl_4 in toluene to reflux for 24 h. In fact, $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ is recovered without decomposition after isolated material is redissolved and heated in toluene at reflux for 7 days. C_{3v} -Symmetric $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ is characterized by equivalent oxazoline groups that are observed in the ^1H NMR spectrum (benzene- d_6) as singlet resonances at 3.28 (CH_2) and 1.36 ppm (Me). A comparison of the IR spectra of $\text{Li}[\text{To}^{\text{M}}]$ versus $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ showed the stretching frequency of the C=N double bond shifts from 1607 cm^{-1} to lower energy at 1545 cm^{-1} .


Amine elimination also proved to be a viable route for the preparation of $[\text{To}^{\text{M}}]\text{Zr}$ compounds.¹⁶ Thus, reaction of $\text{Zr}(\text{NMe}_2)_4$ and $\text{H}[\text{To}^{\text{M}}]$ in benzene yields $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{NMe}_2)_3]$ as a white solid (73%, eq 4). A micromolar scale reaction was monitored in situ by ^1H NMR spectroscopy, revealing that 1 equiv of NHMe_2 is rapidly formed (<10 min). The ^1H NMR spectrum (benzene- d_6) of the C_{3v} -symmetric product contained one set of oxazoline resonances (3.41 and 1.09 ppm) and a singlet corresponding to equivalent NMe_2 groups (3.15 ppm).



In contrast, salt metathesis of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ and 3 equiv of LiNMe_2 (benzene- d_6 , room temperature, 24 h) provides a C_s -symmetric product. The ^1H NMR spectrum of the crude reaction mixture revealed a 1:1 ratio of $[\text{To}^{\text{M}}]:\text{NMe}_2$ groups,

providing support for its assignment as $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{NMe}_2)\text{Cl}_2]$. Under more forcing conditions (6 equiv of LiNMe_2 , 24 h, 60 °C), $\text{Zr}(\text{NMe}_2)_4$ and $\text{Li}[\text{To}^{\text{M}}]$ are obtained. Although substitution of one chloride in $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ by $[\text{NMe}_2]^-$ occurs, formation of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{NMe}_2)_3]$ via salt elimination has not been observed.

To further explore chloride substitutions, a toluene solution of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})\text{Cl}_3]$ and $\text{KO-}t\text{-Bu}$ was stirred overnight at room temperature to provide $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$.



The ^1H NMR spectrum of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$ highlights its C_s -symmetry. As a result of the mirror plane that includes the *tert*-butoxide oxygen, zirconium center, and *trans*-oxazoline, the $-\text{CMe}_2-$ and $-\text{CH}_2-$ groups on that ring are equivalent and appear as singlet resonances (1.52 and 3.38, respectively, labeled as *trans* in the Newman projection in Figure 2).

Although the other two oxazoline rings are related by the mirror plane, the $-\text{CH}_2-$ within each ring are *endo* or *exo* with respect to the *tert*-butoxide and were observed as two coupled doublets (3.38 and 3.25 ppm, $^2J_{\text{HH}} = 8.72$ Hz). The methyl substituents in the CMe_2 groups of these oxazolines are also inequivalent (but not coupled), resulting in two singlet resonances (1.45 and 1.05 ppm are *exo* and *endo*, respectively, assigned with a NOESY experiment). Cross-peaks in a $^1\text{H}-^{15}\text{N}$ HMBC experiment verify that these two peaks are *cis* to the *tert*-butoxide, as they correlate to a ^{15}N resonance at 239 ppm (*N trans* is at 241 ppm).^{11a,17}

A solid-state structure of the product confirmed the N,N,N- κ^3 coordination of $[\text{To}^{\text{M}}]^-$ to zirconium, as well as the overall connectivity of the molecule (Figure 3). The zirconium center's geometry is distorted octahedral with a relatively short Zr–O bond (1.891(3) Å) for a six-coordinate zirconium *tert*-butoxide. Interestingly, $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$ is C_1 -symmetric in the solid state due to inequivalent twist conformations of the three oxazoline rings such that the methyl groups on the 4-position are positioned equatorial (pointed toward zirconium) or axial. Thus, all $\text{Zr}\cdots\text{C}$ (4-Me) distances are inequivalent, ranging from 3.76 to 4.23 Å, creating an unusual steric pocket around the zirconium center. The size of this pocket is best described by the solid angle for $[\text{To}^{\text{M}}]^-$,^{18,19} determined from the crystallographic coordinates to be 5.40 steradians (164° corresponding




Figure 2. Newman projection illustrating the C_s -symmetry of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$. The groups attached to N represent CH_2 and CMe_2 substituents on the oxazoline.




Figure 3. ORTEP diagram of $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$ at 50% probability. Hydrogen atoms are omitted for clarity.

to 43% of a sphere's surface). For comparison, the solid angle for Tp^* is 3.62 steradians (130° corresponding to 30% of a sphere's surface, determined for $\text{Tp}^*\text{Zr}(\text{CH}_2\text{C}_6\text{H}_5)_3$).²

These values for solid angles show that the 4,4-dimethyloxazoline substituents in $[\text{To}^{\text{M}}]^-$ encompass the zirconium center to a greater extent than the 3,5-dimethylpyrazole groups in Tp^* . Thus, tris(4,4-dimethyl-2-oxazolynyl)borate may offer enhanced steric protection for highly reactive metal centers, such as those of cationic zirconium alkyl and hydride complexes.² Additionally, optically active C_3 -symmetric tris(oxazolynyl)borates are readily accessible due to the availability of enantiopure 2*H*-oxazolines.²⁰ These chiral tris(oxazolynyl)borates will complement the chemistry of the related optically active tris(pyrazolyl)borates pioneered by Tolman.²¹ We are currently investigating our achiral complexes as catalysts, as well as preparing chiral analogues for stereoselective polymerizations.¹⁵

Acknowledgment. We thank the U.S. DOE office of Basic Energy Sciences, through the Catalysis Science Grant No. AL-03-380-011, the Roy J. Carver Charitable Trust, and Iowa State University for financial support. We also thank Dr. Bruce Fulton for assistance with ^{15}N NMR measurements.

Supporting Information Available: Experimental procedures, characterization, and X-ray crystallographic data are available free of charge at <http://pubs.acs.org>.

OM800252P

(16) (a) Diamond, G. M.; Rodewald, S.; Jordan, R. F. *Organometallics* **1995**, *14*, 5–7. (b) Diamond, G. M.; Jordan, R. F.; Petersen, J. L. *Organometallics* **1996**, *15*, 4030–4037.

(17) Foltz, C.; Enders, M.; Bellemin-Lapponnaz, S.; Wadepl, H.; Gade, L. H. *Chem.-Eur. J.* **2007**, *13*, 5994–6008.

(18) (a) White, D.; Coville, N. J. *Adv. Organomet. Chem.* **1994**, *36*, 95–158. (b) Taverner, B. C. *J. Comput. Chem.* **1996**, *17*, 1612–1623. (c) The solid angles were calculated from X-ray coordinates of $[\text{Zr}(\kappa^3\text{-Tp}^*)(\text{CH}_2\text{C}_6\text{H}_5)_3]$ (ref 2) and $[\text{Zr}(\kappa^3\text{-To}^{\text{M}})(\text{O-}t\text{-Bu})\text{Cl}_2]$ using the program *Steric* running the algorithms described in ref 18a. (d) For comparison, the cone angles were also calculated using *Steric*: $[\text{To}^{\text{M}}]^-$, 242° ; $[\text{Tp}^*]^-$, 246° .

(19) Taverner, B. C. *Steric Version. 1.1*: Structural Chemistry, University of the Witwatersrand: South Africa, 1995.

(20) (a) Leonard, W. R.; Romine, J. L.; Meyers, A. I. *J. Org. Chem.* **1991**, *56*, 1961–1963. (b) Kamata, K.; Agata, I.; Meyers, A. I. *J. Org. Chem.* **1998**, *63*, 3113–3116.

(21) (a) LeCloux, D. D.; Tolman, W. B. *J. Am. Chem. Soc.* **1993**, *115*, 1153–1154. (b) Kitajima, N.; Tolman, W. B. *Prog. Inorg. Chem.* **1995**, 419–531.