

7-2018

Bacterial wilt symptoms are impacted by host age and involve net downward movement of *Erwinia tracheiphila* in muskmelon

Q. Liu

Iowa State University

Gwyn A. Beattie

Iowa State University, gbeattie@iastate.edu

E. Saalau Rojas

University of Massachusetts Amherst

Mark L. Gleason

Iowa State University, mgleason@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/plantpath_pubs

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/plantpath_pubs/258. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Plant Pathology and Microbiology at Iowa State University Digital Repository. It has been accepted for inclusion in Plant Pathology and Microbiology Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Bacterial wilt symptoms are impacted by host age and involve net downward movement of *Erwinia tracheiphila* in muskmelon

Abstract

Cucurbit bacterial wilt, caused by *Erwinia tracheiphila*, is a damaging disease of cucurbit crops in the Midwest and Northeast U.S. Current management of bacterial wilt relies primarily on insecticide applications to control striped and spotted cucumber beetles (*Acalymma vittatum* and *Diabrotica undecimpunctata howardi*, respectively), which vector *E. tracheiphila*. Development of alternative management strategies is constrained by a lack of understanding of bacterial wilt etiology. The impact of host age on rate on symptom development and extent of bacterial movement in the xylem of muskmelon (*Cucumis melo* cv. Athena) was evaluated following wound inoculation of 2- to 8-week-old plants in growth chamber experiments. Wilting occurred more rapidly in plants after inoculating *E. tracheiphila* into 2- or 4-week-old plants than 6- or 8-week-old plants. Recovery of viable cells from stem segments revealed that vascular spread of *E. tracheiphila* was more extensive below than above the inoculation point. These findings provide experimental evidence that host age impacts the rate of symptom development in cucurbit bacterial wilt and that movement of the xylem-inhabiting pathogen *E. tracheiphila* within muskmelon plants occurs primarily in the downward direction.

Keywords

Phytopathogenic bacteria, vascular wilts, vegetable diseases, ontogenic resistance

Disciplines

Agricultural Science | Agriculture | Plant Pathology

Comments

This is a manuscript of an article published as Liu, Q., G.A. Beattie, E. Saalau Rojas and M.L. Gleason. 2018. Bacterial wilt symptoms are impacted by host age and involve net downward movement of *Erwinia tracheiphila* in muskmelon. *European Journal of Plant Pathology* 151:803-810. doi: [10.1007/s10658-018-1418-7](https://doi.org/10.1007/s10658-018-1418-7). Posted with permission.

1 BACTERIAL WILT SYMPTOMS ARE IMPACTED BY HOST AGE AND INVOLVE NET
2 DOWNWARD MOVEMENT OF *ERWINIA TRACHEIPHILA* IN MUSKMELON
3

4 Q. Liu¹, G. A. Beattie¹, E. Saalau Rojas², and M. L. Gleason¹

5 ¹Department of Plant Pathology and Microbiology, Iowa State University, Ames 50010

6 ² University of Massachusetts-Amherst, Cranberry Station, East Wareham, MA 02538
7

8 Corresponding author: Mark L. Gleason. Email: mgleason@iastate.edu. Telephone: 1-515-294-
9 0579
10
11
12
13
14
15
16
17
18
19

20 **ACKNOWLEDGEMENTS**

21 This research was funded by a Specialty Crop Research Initiative (SCRI) Program Grant
22 (2012-51181-20295) from the U.S. Department of Agriculture National Institute of Food and
23 Agriculture (NIFA). We thank Jean C. Batzer and Xiaoyu Zhang for technical advice and
24 assistance.
25

26 ABSTRACT

27 Cucurbit bacterial wilt, caused by *Erwinia tracheiphila*, is a damaging disease of cucurbit
28 crops in the Midwest and Northeast U.S. Current management of bacterial wilt relies primarily
29 on insecticide applications to control striped and spotted cucumber beetles (*Acalymma vittatum*
30 and *Diabrotica undecimpunctata howardi*, respectively), which vector *E. tracheiphila*.
31 Development of alternative management strategies is constrained by a lack of understanding of
32 bacterial wilt etiology. The impact of host age on rate on symptom development and extent of
33 bacterial movement in the xylem of muskmelon (*Cucumis melo* cv. Athena) was evaluated
34 following wound inoculation of 2- to 8-week-old plants in growth chamber experiments. Wilting
35 occurred more rapidly in plants after inoculating *E. tracheiphila* into 2- or 4-week-old plants
36 than 6- or 8-week-old plants. Recovery of viable cells from stem segments revealed that vascular
37 spread of *E. tracheiphila* was more extensive below than above the inoculation point. These
38 findings provide experimental evidence that host age impacts the rate of symptom development
39 in cucurbit bacterial wilt and that movement of the xylem-inhabiting pathogen *E. tracheiphila*
40 within muskmelon plants occurs primarily in the downward direction.

41
42 KEYWORDS

43 Phytopathogenic bacteria; vascular wilts; vegetable diseases; ontogenic resistance
44

45 INTRODUCTION

46 *Erwinia tracheiphila* (Smith), the causal agent of cucurbit bacterial wilt, is transmitted by
47 striped (*Acalymma vittatum* (F.)) and spotted (*Diabrotica undecimpunctata howardi* (Barber))
48 cucumber beetles. Bacterial wilt causes severe losses in many cultivated cucurbit crops,
49 primarily in the genera *Cucurbita* and *Cucumis*. Economic losses of cucurbit crops from bacterial
50 wilt can reach 75% (Zehnder et al. 1997), and muskmelon (*Cucumis melo* L.) and cucumber
51 (*Cucumis sativus* L.) are among the most susceptible crops (Sherf 1986). Epidemics have
52 occurred primarily in the Midwest and Northeast U.S. as well as in southern Quebec, Canada
53 (Brust 1997, Fleischer et al. 1999, Toussaint et al. 2013); the disease has been restricted
54 primarily to these areas with the exception of a recent report of bacterial wilt on pumpkin and
55 watermelon in New Mexico (Sanogo et al. 2011).

56 Overwintering of *E. tracheiphila* occurs in the foregut and hindgut of striped cucumber
57 beetles (Garcia-Salazar et al. 2000). It is transmitted when *E. tracheiphila*-infested frass of adult
58 beetles comes into contact with fresh wounds on leaves, stems, or floral nectaries (Sasu et al.
59 2010) and the bacteria invade the xylem. *Erwinia tracheiphila* cells multiply in the xylem and
60 can block water flow (Main and Walker 1971). Infected plants initially exhibit wilting of leaves
61 near the infection site, followed by wilting of vines and eventual collapse and plant death.

62 Current management for cucurbit bacterial wilt relies primarily on insecticide
63 applications (Cavanagh et al. 2009), but this approach is costly and potentially risks the health of
64 humans, pollinators, insectivorous birds and other ecosystem service providers (Cavanagh et al.
65 2009, Potts et al. 2010). Although watermelon (*Citrullus lanatus* var. *lanatus*) is resistant to
66 bacterial wilt and a few cultivars of cucumber (*Cucumis sativus*) are tolerant, resistant cultivars
67 are not commercially available for most cucurbit crops. Therefore, a clearer understanding of the
68 bacterial wilt infection process may yield insights to assist plant breeders in developing such
69 cultivars. A starting point toward this goal is to understand the impact of host age on disease
70 progress, as this may help breeders develop efficient protocols for screening candidate lines for
71 resistance. Moreover, tracing patterns of pathogen movement in the xylem could result in a

72 deeper understanding of disease development.

73 The impact of plant age on wilt symptom development has been investigated for several
 74 xylem-inhabiting bacterial pathogens. Plant age affects bacterial wilt and canker of tomato
 75 caused by *Clavibacter michiganensis* subsp. *michiganensis*; tomato transplants up to the 17- to
 76 18-leaf stage wilted and died after inoculation, whereas plants that were inoculated after that
 77 stage exhibited only mild symptoms (Sharabani et al. 2013). Similarly, *Ralstonia solanacearum*
 78 caused earlier symptoms, higher disease incidence, and greater disease severity of bacterial wilt
 79 of tomato in 2- to 3-week-old seedlings than in 5- to 6-week-old plants (Thomas and Upreti
 80 2014). For cucurbit bacterial wilt, the relationship between plant age and symptom development
 81 has been examined only for the relatively resistant host genera *Cucurbita* and *Citrullus*. In
 82 pumpkin (*Cucurbita pepo* L.), which exhibits intermediate resistance to *E. tracheiphila* (Sherf
 83 1986), seedlings inoculated at the cotyledon stage were more susceptible than older plants and
 84 resistance increased sharply with plant age (Brust 1997). Seedlings of watermelon (*Citrullus*
 85 *lanatus*), which is highly resistant to *E. tracheiphila* (Watterson et al. 1971), exhibited more rapid
 86 symptom development than older plants and resistance increased with age (Watterson et al.
 87 1971). However, no such relationships have been assessed for the highly susceptible cucurbits in
 88 the genus *Cucumis*, such as muskmelon or cucumber, which are at the greatest risk of economic
 89 losses from bacterial wilt.

90 The mechanisms by which *E. tracheiphila* causes bacterial wilt have not yet been
 91 investigated in detail. Absence of evidence for pectolytic enzyme production suggests that
 92 physical occlusion by the bacteria is the primary cause of xylem dysfunction (Main and Walker,
 93 1971, Watterson, Williams, and Durbin, 1971). Moreover, the presence of strands of ooze as
 94 infected stems are cut and drawn apart suggests the presence of extracellular polysaccharides,
 95 which could also contribute to xylem blockage, as with wilt by *R. solanacearum*, *Pantoea*
 96 *stewartii* and *Xylella fastidiosa* (Ayers et al. 1979, Beck von Bodman et al. 1998, Saile et al.
 97 1997). Factors that contribute to the virulence of other bacterial xylem pathogens, such as
 98 biofilm formation, quorum sensing, outer membrane vesicle production and motility (Herrera et
 99 al. 2008, Ionescu et al. 2014, Koutsoudis et al. 2006, Meng et al. 2005), have not yet been
 100 examined in *E. tracheiphila*. Recently, observational studies using a constructed bioluminescent
 101 strain of *E. tracheiphila* provided the first evidence that the bacterium can move both upward in
 102 muskmelon seedlings and downward into the roots following inoculation (Vrisman et al. 2016).
 103 The objectives of the present study were to i) quantify the impact of host age on development of
 104 wilt symptoms in muskmelon and ii) trace the movement of *E. tracheiphila* in the xylem
 105 following inoculation.

106

107 MATERIALS AND METHODS

108 **Plant growth conditions.** Muskmelon (*Cucumis melo* cv. Athena) seeds were planted in
 109 a 1:1:1 matrix of peat moss, coarse perlite, and Metro-Mix 300 (Sun Gro Horticulture,
 110 Vancouver, BC, Canada). A single seed was planted in each 650 cm³ pot. The pots were
 111 incubated at 26°C under a daily regimen of 14 h light and 10 h darkness under ambient relative
 112 humidity (RH) in growth chambers. Plants were watered daily and fertilized weekly (NPK: 15-5-
 113 15: Miracle Gro®, The Scotts Co., Marysville, OH).

114 **Bacterial strain and inoculum preparation.** *E. tracheiphila* strain SCR3 (Saalau Rojas
 115 et al. 2012) was used in this study. This strain is a spontaneous rifampicin-resistant derivative of
 116 an isolate from a symptomatic muskmelon plant grown in Iowa, U.S. Pathogenicity of this
 117 isolate was confirmed by puncture inoculation of the first true leaf of 2-week-old muskmelon

118 plants in growth chamber trials (Saalau Rojas et al. 2012). To prepare cells for plant inoculation
119 assays, SCR3 was recovered from -80°C storage on solid nutrient agar peptone medium (de
120 Mackiewicz et al. 1998) that was amended with rifampicin (75 µg/ml) (NAP-Rif). Cells were
121 grown at 27°C for 3 days, then transferred to fresh NAP-Rif medium and grown at 27°C for
122 another 3 days. Bacterial suspensions were prepared by recovering SCR3 colonies from the
123 surface of solid NAP-Rif medium and suspending them in 10 mM phosphate-buffered saline
124 (PBS) to a concentration of approximately 2.5×10^8 CFU/ml, based on a standard curve relating
125 cell density to optical density at 540 nm.

126 **Symptom expression experiments.** Muskmelon seeds were planted 2, 4, and 6 weeks
127 before inoculation in the first experiment and 2, 4, 6 and 8 weeks before inoculation in a second
128 experiment, in order to create multiple cohorts of plants that varied in age at the time of
129 inoculation. In each experiment, plant age at inoculation was considered a treatment, and each
130 treatment included four single-seedling replicates. A 100-µl droplet of inoculum was applied at
131 the base of the adaxial surface of the youngest fully expanded leaf, followed by puncturing the
132 leaf at the site of the droplet with a 28.6-mm-diameter, 60-pin florist's pin frog (Kenzan Pin
133 Frog, sold by www.save-on-crafts.com). Next, the pipette tip was rubbed lightly against the
134 punctured site in order to ensure maximal contact of the inoculum with the puncture wounds and
135 an additional 100 µl of suspension was applied to the punctured site on the leaf, after which the
136 pipette tip was again rubbed lightly on the site. Control plants were inoculated with PBS (10
137 mM) buffer in the same manner as described above. After inoculation, plants were incubated in a
138 growth chamber at 26°C under a daily regimen of 14 h light and 10 h darkness and ambient RH.
139 Numbers of wilted and asymptomatic leaves on each plant and their locations relative to the site
140 of inoculation were determined daily until all *E. tracheiphila*-inoculated plants displayed wilt
141 symptoms. Rate of wilting was determined by estimating the number of days for 50% of the
142 plant to show wilt symptoms.

143 **Pathogen movement experiments.** Plants that were 2, 4, 6, and 8 weeks old were
144 inoculated as described above, with plant age at inoculation considered as a treatment. On days
145 3, 7, 14, and 21 after inoculation, four plants of each treatment were chosen arbitrarily for
146 destructive sampling. Stem segment samples (5 cm long and devoid of nodes) from each
147 treatment were excised above and below the point of inoculation. In the first run of the
148 experiment, all internodes were sampled for the plants that were 2 or 4 weeks old at inoculation
149 (referred to here as 2-week-old and 4-week-old plants), whereas every third internode was
150 sampled in older plants (referred to here as 6-week-old and 8-week-old plants). In the second run
151 of the experiment, every internode was sampled from the 2- and 4-week-old plants, whereas
152 every second internode was sampled from the 6- and 8-week-old plants. These internode stem
153 segments were surface-sterilized by spraying them with 70% ethanol and then air dried on sterile
154 paper towels. A stem segment was cut transversely at its midpoint, the cut surface was imprinted
155 on the surface of NAP-Rif medium, and the resulting imprint was streaked for single colonies.
156 The presence of colonies exhibiting *E. tracheiphila* morphology was recorded after 4 days of
157 incubation at 27°C. The number of wilted and asymptomatic leaves per plant was also counted
158 on each sampling date. Sampling was terminated in each treatment when all leaves had wilted.
159 Bacterial movement in the upward direction vs. the downward direction was compared on the
160 basis of the mean number of internodes from the site of inoculation to the furthest internode from
161 which *E. tracheiphila* cells were recovered.

162 **Data analysis.** In the symptom expression experiments, disease progress was determined
163 based on the area under the disease progress curve (AUDPC) using the trapezoidal method

164 (Simko and Piepho 2012) and the means were compared with a Fisher's least significant
165 difference test using SAS 9.1 (SAS Institute Inc., Cary, NC, USA). For each treatment in the
166 pathogen movement experiments, the mean number of internodes from which *E. tracheiphila*
167 was recovered above vs. below the point of inoculation was compared using a Student's *t*-test.
168

169

170 RESULTS

171 **Rate of wilting.** Leaves on the inoculated muskmelon plants began to wilt as early as 4
172 days after inoculation. In both runs of the experiment, the inoculated leaves wilted rapidly
173 regardless of plant age at inoculation (Table 1). In the first experiment, the plants that were 2
174 weeks old at inoculation wilted significantly ($p < 0.05$) faster than those that were 4 or 6 weeks
175 old, as reflected in the mean number of days from inoculation until wilting of 50% of leaves and
176 area under the disease progress curve (AUDPC) (Simko et al. 2012). Similarly, plants that were 4
177 weeks old at the time of inoculation wilted significantly faster than those that were 6 weeks old.
178 In the second run of the experiment, although the plants in each age group wilted more slowly
179 than similarly-aged plants in the first experiment, the plants that were younger at the time of
180 inoculation again wilted faster than those that were older. Specifically, the 2- or 4-week-old
181 plants wilted significantly ($p < 0.05$) faster than the 6- and 8-week-old plants based on the number
182 of days required for plants within each treatment to display 50% of wilted leaves and on the
183 AUDPC (Table 1). No wilting was observed on the control plants in either experiment.

184 **Bacterial movement.** In each of two replicate experiments to evaluate bacterial
185 movement from the site of inoculation, *E. tracheiphila* cells were first recovered from stem
186 samples 7 days postinoculation (dpi) and were isolated from sites both above and below the
187 inoculation point (Fig. 1). There was evidence of upward movement of *E. tracheiphila*,
188 determined by recovery of bacteria from stem segments above the inoculation site at 7, 14 and
189 21 dpi, and at sites as far as 5 and 8 internodes above the inoculation site in the first and second
190 runs of the experiment, respectively. Stem growth above the inoculation point continued until 14
191 dpi on plants that had been 4, 6, or 8 weeks old at inoculation, but the pathogen was not
192 recovered from the uppermost internodes of the stem even at 21 dpi, indicating that its movement
193 in the upward direction was limited.

194 At each sampling time, *E. tracheiphila* moved much further downward than upward, with
195 movement sometimes limited by reaching the lowest possible node (Fig. 1). In contrast to the
196 upward movement to a maximum of 5 to 8 internodes above the inoculation site, bacteria moved
197 downward as far as 9, 13, and 23 nodes by 7, 14 and 21 dpi, respectively, in the first run of the
198 experiment regardless of plant age at inoculation (Fig. 1A), and to similar distances in the second
199 run of the experiment (Fig. 1B). On plants inoculated at 6 or 8 weeks of age, *E. tracheiphila* cells
200 were detected in nearly all of the stem segments sampled below the inoculation point at 21 dpi,
201 and the distance of movement of the bacterium at 14 and 21 dpi was significantly ($p < 0.05$)
202 greater below than above the point of inoculation.

203 In both experiments, the 2-week-old and 4-week-old plants died before reaching 14 and
204 21 dpi respectively; thus, data for these treatments are not presented for those time points. By 21
205 dpi in both runs of the experiment, the stems of all of the plants that were 6 weeks old at
206 inoculation had withered and died above the inoculation point; isolations from that portion of the
207 plant in the first run were not attempted because the pathogen does not survive in dead host
208 tissue (Latin 2000). In general, *E. tracheiphila* reached the lowest node of each plant by the time
209 of the plant's death.

210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255

DISCUSSION

Our results provide the first experimental evidence that the rate of wilting of *Cucumis* sp. crop by *E. tracheiphila* is impacted by host age, with young plants developing wilt symptoms significantly faster than older ones based on the time required for 50% of the leaves to wilt. Although grower guides frequently state that young cucurbit plants are more susceptible to infection than older plants (Brust 1997a, Watterson et al. 1971), experimental evidence supporting this assertion is absent for *Cucumis* spp. In the Brust (1997a) study, more wilt was observed in pumpkin seedlings that had been inoculated at the cotyledon stage than at the later stage of ≥ 1 true leaf, although most of the seedlings with true leaves never showed wilt and recovered from the inoculation. The general trend observed in the Brust study – increasing resistance with increasing plant age – agrees with results of the present study of muskmelon.

Knowledge that susceptibility to bacterial wilt decreases as muskmelon plants age has important management implications. For example, our evidence that plants are more susceptible to bacterial wilt when they are young can help to inform optimal timing for reduced-insecticide management strategies such as the deployment of row covers as protective barriers against cucumber beetles (Mueller et al. 2006, Saalau Rojas et al. 2011). In particular, although row covers should be deployed during the most susceptible period to suppress bacterial wilt, deciding when to remove them should factor in both the probability of pathogen transmission by cucumber beetles (Saalau Rojas et al. 2011) and plant phenology related to resistance. In addition, clarifying responses to inoculation as a function of seedling age should help plant breeders to optimize screening assays for bacterial wilt resistance. For example, using plants that are 4 weeks old at inoculation might be a cost-effective option because they are relatively small and thus require minimal growth space, and moderate in susceptibility when compared with 2-, 6-, and 8-week-old plants. Moreover, their requirement for about 8 dpi to show symptoms enables sufficient observation time to compare symptoms among breeding lines. However, additional, season-long experiments in the field would be needed in order to comprehend impact on yield and fruit quality when plants become infected at later stages of the growing season.

The mechanism responsible for a decreased wilting rate as muskmelon plants age is unclear. Ontogenic resistance could result from factors such as increased production of phytochemicals and/or the development of physical barriers that slow disease progress (Panter and Jones 2002). Alternatively, the pathogen may be diluted if plant growth exceeds pathogen growth, which could slow the rate of wilting. The correlation of plant age with plant size makes it difficult to separate the impacts of age versus size; this is particularly true for cucurbit crops, most of which increase rapidly in size during the early part of the growing season. It is possible that both ontogenic and plant size-related factors may operate to slow wilting in older, larger plants. Further experimentation will be required to unravel the causal mechanisms of this phenomenon.

Our experiments are the first to document that internal movement of *E. tracheiphila* following infection is more rapid in a downward than upward direction. In common with other xylem-limited vascular wilt diseases, symptom expression as indicated by visible wilt progressed distally beyond the nodes from which the pathogen was recovered, presumably as sieve plates became blocked and water flow ceased (Holland et al. 2014, McElrone et al. 2003). Such a blockage of the upward movement of water inside the vascular system may help to explain our observation that *E. tracheiphila* moved in a primarily downward rather than upward direction from the inoculation point. Vrisman et al (2016) provided observational evidence that a

256 bioluminescent-labeled strain of *E. tracheiphila* moved not only upward from the inoculated leaf
 257 of a muskmelon seedling but also downward into the roots. The mechanism driving bacterial
 258 movement against the xylem flow is unclear, but is consistent with the movement exhibited by
 259 two other xylem-limited vascular wilt pathogens: *Xylella fastidiosa*, the causal agent of Pierce's
 260 disease of grape (Meng et al. 2005), and *Acidovorax avenae* subsp. *citrulli*, the causal agent of
 261 bacterial fruit blotch of cucurbits (Bahar et al. 2010). *X. fastidiosa*, a nonflagellated bacterial
 262 pathogen, was shown to spread in the xylem via motility mediated by type IV pili (Meng et al.
 263 2005). Type IV pili, which are like grappling hooks, enable the bacterial cells to jerk forward
 264 along a surface in a form of motility known as twitching (Mattick 2002). *X. fastidiosa* mutants
 265 defective in these pili showed reduced downward colonization of the xylem (Meng et al. 2005).
 266 As a xylem-limited pathogen, *E. tracheiphila* may employ a similar mechanism of movement.
 267 This is consistent with the recent discovery that the putative type IV pili genes are conserved
 268 among *Erwinia* spp., including *E. tracheiphila* (Shapiro 2012). Our study, the first to quantify
 269 the relative movement of the pathogen upward and downward following infection, is a further
 270 step in understanding the movement of *E. tracheiphila* in the xylem and helps to set a foundation
 271 for evaluating the role of pilus genes in downward movement.

272

273 LITERATURE CITED

- 274 Ayers, A. R., Ayers, S. B., and Goodman, R. N. (1979). Extracellular polysaccharide of *Erwinia*
 275 *amylovora*: a correlation with virulence. *Appl. Environ. Microb.*, 38, 659-666.
- 276 Bahar, O., De la Fuente, L., and Burdman, S. (2010). Assessing adhesion, biofilm formation and
 277 motility of *Acidovorax citrulli* using microfluidic flow chambers. *FEMS Microbiol. Lett.*,
 278 312, 33-39.
- 279 Beck von Bodman, S., Majerczak, D. R., and Coplin, D. L. (1998). A negative regulator
 280 mediates quorum-sensing control of exopolysaccharide production in *Pantoea stewartii*
 281 subsp. *stewartii*. *Proc. Natl. Acad. Sci. USA*, 95, 7687-7692.
- 282 Brust, G. E. (1997a). Differential susceptibility of pumpkins to bacterial wilt related to plant
 283 growth stage and cultivar. *Crop Prot*, 16, 411-414.
- 284 Brust, G. E. (1997b). Seasonal variation in percentage of striped cucumber beetles (Coleoptera:
 285 Chrysomelidae) that vector *Erwinia tracheiphila*. *Environ. Entomol.*, 26, 580-584.
- 286 Cavanagh, A., Hazzard, R., Adler, L.S., and Boucher, J. (2009). Using trap crops for control of
 287 *Acalymma vittatum* (Coleoptera: Chrysomelidae) reduces insecticide use in butternut
 288 squash. *J. Econ. Entomol.*, 102, 1101-1107.
- 289 de Mackiewicz, D., Gildow, F. E., Blua, M., Fleischer, S. J., and Lukezic, F. L. (1998).
 290 Herbaceous weeds are not ecologically important reservoirs of *Erwinia tracheiphila*.
 291 *Plant Dis.*, 82, 521-529.
- 292 Fleischer, S. J., de Mackiewicz, D., Gildow, F. E., and Lukezic, F. L. (1999). Serological
 293 estimates of the seasonal dynamics of *Erwinia tracheiphila* in *Acalymma vittata*
 294 (Coleoptera : Chrysomelidae). *Environ. Entomol.*, 28, 470-476.
- 295 Garcia-Salazar, C., Gildow, F. E., Fleischer, S. J., Cox-Foster, D., and Lukezic, F. L. (2000).
 296 Alimentary canal of adult *Acalymma vittata* (Coleoptera : Chrysomelidae): morphology
 297 and potential role in survival of *Erwinia tracheiphila* (Enterobacteriaceae). *Can.*
 298 *Entomol.*, 132, 1-13.
- 299 Herrera, C. M., Koutsoudis, M. D., Wang, X. L., and von Bodman, S. B. (2008). *Pantoea*
 300 *stewartii* subsp. *stewartii* exhibits surface motility, which is a critical aspect of Stewart's
 301 wilt disease development on maize. *Mol. Plant-Microb. Interact.*, 21, 1359-1370.

- 302 Holland, R. M., Christiano, R. S. C., Gamliel-Atinsky, E., and Scherm, H. (2014). Distribution of
 303 *Xylella fastidiosa* in blueberry stem and root sections in relation to disease severity in the
 304 field. *Plant Dis.*, 98, 443-447.
- 305 Ionescu, M., Zaini, P. A., Baccari, C., Tran, S., da Silva, A. M., and Lindow, S. E. (2014).
 306 *Xylella fastidiosa* outer membrane vesicles modulate plant colonization by blocking
 307 attachment to surfaces. *Proc. Natl. Acad. Sci. USA*, 111, E3910-E3918.
- 308 Koutsoudis, M. D., Tsaltas, D., Minogue, T. D., and von Bodman, S. B. (2006). Quorum-sensing
 309 regulation governs bacterial adhesion, biofilm development, and host colonization in
 310 *Pantoea stewartii* subspecies *stewartii*. *Proc. Natl. Acad. Sci. USA*, 103, 5983-5988.
- 311 Latin, R.X. (2000). Bacterial wilt. In: APSnet Features: Scary Diseases Haunt Pumpkins and
 312 Other Cucurbits.
 313 <http://www.apsnet.org/publications/apsnetfeatures/Pages/BacterialWilt.aspx>
- 314 Main, C. E., and Walker, J. C. (1971). Physiological responses of susceptible and resistant
 315 cucumber to *Erwinia tracheiphila*. *Phytopathology*, 61, 518-522.
- 316 Mattick, J. S. (2002). Type IV pili and twitching motility. *Annu. Rev. Microbiol.*, 56, 289-314.
- 317 McElrone, A. J., Seral, J. L., and Forseth, I. N. (2003). Interactive effects of water stress and
 318 xylem-limited bacterial infection on the water relations of a host vine. *J. Exp. Bot.*, 54,
 319 419-430.
- 320 Meng, Y. Z., Li, Y. X., Galvani, C. D., Hao, G. X., Turner, J. N., Burr, T. J., and Hoch, H. C.
 321 (2005). Upstream migration of *Xylella fastidiosa* via pilus-driven twitching motility. *J.*
 322 *Bacteriol.*, 187, 5560-5567.
- 323 Mueller, D. S., Gleason, M. L., Sisson, A. J., and Massman, J. M. (2006). Effect of row covers
 324 on suppression of bacterial wilt of muskmelon in Iowa. *Plant Health Progress*,
 325 doi:10.1094/PHP-2006-1020-02-RS.
- 326 Panter, S. N., and Jones, D. A. (2002). Age-related resistance to plant pathogens. *Adv. Bot. Res.*,
 327 38, 251-280.
- 328 Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., and Kunin, W. E.
 329 (2010). Global pollinator declines: Trends, impacts and drivers. *Trends Ecol. Evol.*, 25,
 330 345-353.
- 331 Saalau Rojas, E., and Gleason, M. L. (2012). Epiphytic survival of *Erwinia tracheiphila* on
 332 muskmelon (*Cucumis melo* L.). *Plant Dis.*, 96, 62-66.
- 333 Saalau Rojas, E., Gleason, M. L., Batzer, J. C., and Duffy, M. (2011). Feasibility of delaying
 334 removal of row covers to suppress bacterial wilt of muskmelon (*Cucumis melo*). *Plant*
 335 *Dis.*, 95, 729-734.
- 336 Saile, E., McGarvey, J. A., Schell, M. A., and Denny, T. P. (1997). Role of extracellular
 337 polysaccharide and endoglucanase in root invasion and colonization of tomato plants by
 338 *Ralstonia solanacearum*. *Phytopathology*, 87, 1264-1271.
- 339 Sanogo, S., Etarock, B. F., and Clary, M. (2011). First report of bacterial wilt caused by *Erwinia*
 340 *tracheiphila* on pumpkin and watermelon in New Mexico. *Plant Dis.*, 95, 1583-1583.
- 341 Sasu, M. A., Seidl-Adams, I., Wall, K., Winsor, J. A., and Stephenson, A. G. (2010). Floral
 342 transmission of *Erwinia tracheiphila* by cucumber beetles in a wild *Cucurbita pepo*.
 343 *Environ. Entomol.*, 39, 140-148.
- 344 Shapiro, L. (2012). A to ZYMV guide to *Erwinia tracheiphila* infection: an ecological and
 345 molecular study. Doctoral thesis. Pennsylvania State University, 160 pp.

- 346 Sharabani, G., Shtienberg, D., Borenstein, M., Shulhani, R., Lofthouse, M., Sofer, M.,
347 Chalupowicz, L., Barel, V., and Manulis-Sasson, S. (2013). Effects of plant age on
348 disease development and virulence of *Clavibacter michiganensis* subsp. *michiganensis* on
349 tomato. *Plant Pathol.*, 62, 1114-1122.
- 350 Sherf, A. F., MacNab, A. A. (1986). Bacterial wilt. Pages 307-311 In: *Vegetable Diseases and*
351 *Their Control*. Wiley Interscience, New York.
- 352 Simko, I., and Piepho, H. P. (2012). The area under the disease progress stairs: calculation,
353 advantage, and application. *Phytopathology*, 102, 381-389.
- 354 Thomas, P., and Upreti, R. (2014). Influence of seedling age on the susceptibility of tomato
355 plants to *Ralstonia solanacearum* during protray screening and at transplanting. *Am. J.*
356 *Plant Sci.*, 5, 1755-1762.
- 357 Toussaint, V., Ciotola, M., Cadieux, M., Racette, G., Duceppe, M. O., and Mimee, B. (2013).
358 Identification and temporal distribution of potential insect vectors of *Erwinia*
359 *tracheiphila*, the causal agent of bacterial wilt of cucurbits. *Phytopathology*, 103, 147-
360 147.
- 361 Vrisman, C.M., Deblais, L., Rajashekara, G., and Miller, S.A. (2016). Differential colonization
362 dynamics of cucurbit hosts by *Erwinia tracheiphila*. *Phytopathology*, 106, 684-692.
- 363 Watterson, J. C., Williams, P. H., and Durbin, R. D. (1971). Response of cucurbits to *Erwinia*
364 *tracheiphila*. *Plant Dis. Rep.*, 55, 816-819.
- 365 Zehnder, G., Klopper, J., Yao, C. B., and Wei, G. (1997). Induction of systemic resistance in
366 cucumber against cucumber beetles (Coleoptera: Chrysomelidae) by plant growth-
367 promoting rhizobacteria. *J. Econ. Entomol.*, 90, 391-396.
- 368
- 369

370 **Table 1.** Impact of plant age at the time of inoculation with *Erwinia tracheiphila* on the rate of
 371 wilt and disease progression in muskmelon (*Cucumis melo* cv. Athena) plants in two growth
 372 chamber experiments.

Expt ^a	Plant age at inoculation (weeks)	Mean days to wilting of the inoculated leaf ^b	Mean days to wilting of 50% of the leaves ^{b,c}	AUDPC ^{b,d}
1	2	5.3 a	5.3 a	1,667 a
	4	5.0 a	11.3 b	1,062 b
	6	4.7 a	14.0 c	777 c
2	2	5.7 a	5.7 a	2,539 a
	4	6.0 a	8.5 a	2,289 a
	6	6.0 a	18.0 b	1,545 b
	8	7.3 a	20.7 b	1,240 b


373 ^a Two independent experiments were performed. Each experiment included four replicate plants
 374 per treatment.

375 ^b In each experiment, the data were subjected to Fisher's least significant difference (LSD) test to
 376 determine differences among the means using SAS 9.1 (SAS Institute Inc., Cary, NC, USA).
 377 Values in a column that are followed by the same letter do not differ significantly ($p < 0.05$).
 378 $n = 4$.

379 ^c Data include the inoculated leaf.

380 ^d AUPDC, area under the disease progress curve (Simko and Piepho 2012).

381



382
383

384 **Fig. 1** Directional movement of *E. tracheiphila* in stems of muskmelon (*Cucumis melo* cv.
385 Athena) plants inoculated at different ages. The results of two replicate experiments are shown in
386 (a) and (b). Data shown are the mean number of internodes from the site of inoculation to the
387 point where *E. tracheiphila* cells were recovered (black bars) or were not recovered (hatched
388 bars) from stem segments. Plants were 2, 4, 6, or 8 weeks old when the youngest true leaf was
389 inoculated. The zero point on the y-axis indicates the point of inoculation. Positive numbers on
390 the y-axis indicate the number of internodes above the inoculation point, whereas negative
391 numbers indicate number of internodes below the inoculation point. Data are means of four
392 replicates per treatment. * indicates sampling times for which the bacterial movement from the
393 site of inoculation was significantly (Student's *t*-test, $p < 0.05$) greater in the downward than
394 upward direction. n = 4. # = indicates samples that were too desiccated to attempt to recover *E.*
395 *tracheiphila*.