

2-2018

Genomes of the Hymenoptera

Michael G. Branstetter
U.S. Department of Agriculture

Anna K. Childers
U.S. Department of Agriculture

Diana Cox-Foster
U.S. Department of Agriculture

Keith R. Hopper
U.S. Department of Agriculture

Karen M. Kapheim
Utah State University

See next page for additional authors

Follow this and additional works at: https://lib.dr.iastate.edu/eeob_ag_pubs

 Part of the [Behavior and Ethology Commons](#), [Entomology Commons](#), and the [Genetics and Genomics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/eeob_ag_pubs/269. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

Genomes of the Hymenoptera

Abstract

Hymenoptera is the second-most sequenced arthropod order, with 52 publically archived genomes (71 with ants, reviewed elsewhere), however these genomes do not capture the breadth of this very diverse order (Figure 1, Table 1). These sequenced genomes represent only 15 of the 97 extant families. Although at least 55 other genomes are in progress in an additional 11 families (see Table 2), stinging wasps represent 35 (67%) of the available and 42 (76%) of the in progress genomes. A more comprehensive catalog of hymenopteran genomes is needed for research into the evolutionary processes underlying the expansive diversity in terms of ecology, behavior, and physiological traits within this group. Additional sequencing is needed to generate an assembly for even 0.05% of the estimated 1 million hymenopteran species, and we recommend premier level assemblies for at least 0.1% of the >150,000 named species dispersed across the order. Given the haplodiploid sex determination in Hymenoptera, haploid male sequencing will help minimize genome assembly issues to enable higher quality genome assemblies.

Disciplines

Behavior and Ethology | Ecology and Evolutionary Biology | Entomology | Genetics and Genomics

Comments

This article is published as Branstetter, M., Childers, A.K., Cox-Foster, D., Hopper, K.R., Kapheim, K.M., Toth, A.L., Worley, K.C. 2017. Genomes of Hymenoptera. *Current Opinion in Insect Science* 25: 65-75. doi: [10.1016/j.cois.2017.11.008](https://doi.org/10.1016/j.cois.2017.11.008).

Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

Authors

Michael G. Branstetter, Anna K. Childers, Diana Cox-Foster, Keith R. Hopper, Karen M. Kapheim, Amy L. Toth, and Kim C. Worley



Genomes of the Hymenoptera

Michael G Branstetter¹, Anna K Childers², Diana Cox-Foster¹,
Keith R Hopper³, Karen M Kapheim⁴, Amy L Toth⁵ and
Kim C Worley⁶

Hymenoptera is the second-most sequenced arthropod order, with 52 publically archived genomes (71 with ants, reviewed elsewhere), however these genomes do not capture the breadth of this very diverse order (Figure 1, Table 1). These sequenced genomes represent only 15 of the 97 extant families. Although at least 55 other genomes are in progress in an additional 11 families (see Table 2), stinging wasps represent 35 (67%) of the available and 42 (76%) of the in progress genomes. A more comprehensive catalog of hymenopteran genomes is needed for research into the evolutionary processes underlying the expansive diversity in terms of ecology, behavior, and physiological traits within this group. Additional sequencing is needed to generate an assembly for even 0.05% of the estimated 1 million hymenopteran species, and we recommend premier level assemblies for at least 0.1% of the >150,000 named species dispersed across the order. Given the haplodiploid sex determination in Hymenoptera, haploid male sequencing will help minimize genome assembly issues to enable higher quality genome assemblies.

Addresses

¹ Pollinating Insect-biology, Management, Systematics Research Unit, USDA-ARS, Logan, UT 84322, United States

² Bee Research Laboratory, USDA-ARS, Beltsville, MD 20705, United States

³ Beneficial Insects Introduction Research Unit, USDA-ARS, Newark, DE 19713, United States

⁴ Utah State University, Department of Biology, Logan, UT 84322, United States

⁵ Iowa State University, Department of Ecology, Evolution, and Organismal Biology and Department of Entomology, Ames, IA 50011, United States

⁶ Human Genome Sequencing Center, and Department of Molecular and Human Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, TX 77030, United States

Current Opinion in Insect Science 2018, 25:65–75

This review comes from a themed issue on **Insect genomics**

Edited by **Christopher Childers, Anna Childers and Stephen Richards**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 22nd November 2017

<https://doi.org/10.1016/j.cois.2017.11.008>

2214-5745/© 2017 Elsevier Inc. All rights reserved.

Introduction


The order Hymenoptera is both ancient and hyper-diverse, having fossils that date back to the Triassic [1], and a richness that exceeds 150,000 described and one million estimated species [2,3]. One of the ‘big four’ insect orders, Hymenoptera comprise diverse species including sawflies and wood wasps (‘Symphyta’), parasitoid wasps (‘Parasitica’), and stinging wasps (Aculeata), which includes the ubiquitous and ecologically dominant ants, bees and social wasps. With an astonishing diversity of biologically interesting traits Hymenoptera have significant economic impact (e.g. biological control and pollination) and thus was one of the first insect orders to benefit from genome sequencing [4]. Despite additional genomes sequenced (Table 1) or in preparation (Figure 1, Table 2), genomic resources are lacking for most major lineages, which comprise 28 superfamilies, 97 families, and 8422 genera. Herein we discuss insights gained and opportunities for improvements for hymenopteran genomics (noting that ants are reviewed separately). Our proposed roadmap for future hymenopteran genome sequencing optimizes benefit to the research community.

Features of hymenopteran genomes

Hymenopteran genomes possess some notable and unique features. Hymenoptera are haplodiploid: unfertilized eggs produce haploid males and fertilized eggs produce diploid females [5]. Haplodiploidy engenders interesting biology related to sex determination [6], control of sex ratios, and relatedness [7], but is also useful for whole genome sequencing. Enough DNA can sometimes be extracted from a single large haploid male to provide material for whole genome sequencing without genetic variation (e.g. [8]). Another notable feature of some Hymenoptera is extremely high recombination rates, especially in social species such as the honey bee, where recombination rates are among the highest known for any organism [9].

Among sequenced examples, hymenopteran genomes are moderate in size (80% are between 180 and 340 Mb) with a few exceptions [10,11,12**]. Most possess 12,000–20,000 genes (note counts are highly annotation-pipeline and assembly contiguity dependent [13*]), with a relatively low content of repetitive and transposable elements. One unusual feature is low GC content, which ranges from 30 to 45% depending on the species [8].

Figure 1



Hymenoptera phylogeny and genome assemblies. Phylogeny based upon [83**,88**], with dotted lines marking lineages of uncertain placement, branches not to scale (=cladogram). Major groups shown on right, individual superfamilies listed with columns indicating the numbers of families, genera, species, genomes in NCBI (in progress).

Although the reason for low GC content is not yet understood, it may be related to GC-biased gene conversion and high recombination rates [14]. Due to their relatively small size and simple structure, hymenopteran genomes are readily assembled and highly tractable for genome sequencing [15*].

Potential genomic insights into Hymenoptera biology

The Hymenoptera include everything from herbivores to pollinators to predators to parasites, and the species filling each of these ecological roles are endowed with a complex set of physiological and behavioral adaptations. Genomic studies have informed applied questions related to pollinator health [16] and biological control using parasitoid wasps [17]. Comparative genomics approaches have unlocked a deeper understanding of the molecular evolutionary basis for these adaptations, including the

evolution of eusociality [8,12**,18], social parasitism [19,20*], venom function [19,20*,21,22*], and behavioral host specificity [23,24*].

Pollinator health and management research has greatly benefited from the availability of reference genomes for many important pollinators (Table 1). Genomic approaches to pollinator health allow us to screen for disease, elucidate the effects of parasites, and investigate the immune response to environmental stressors and pathogens. Examples include developing new biomarkers of honey bee colony health [25–29], characterizing environmental stress responses and optimal developmental temperature regimes for the alfalfa leafcutting bee (an intensively managed solitary pollinator) [30,31] and developing other genomic and transcriptomic indicators of health in wild and managed pollinators [32–39]. Honey bee genomic nutrition research has been fruitful in

generating insights into mechanisms of bee health [40–43]. Expanding this emerging field to native, unmanaged bees can provide a comparative perspective on how wild bee health is influenced by environmental diet restriction due to habitat loss [44]. This approach can also improve our understanding of innate and acquired immunity function among bees, which is critical for management and conservation efforts [45,46].

The vast majority of Hymenoptera species are parasitoids (green in Figure 1 and Table 2) as the majority of insect species are attacked by at least one species of parasitic Hymenoptera [47–51], with complex and intimate host-parasitoid interactions [52] and narrow host ranges [53] which may drive ecological speciation [54,55]. Parasitoids are economically important as biological control agents of invasive pests, reducing pest abundance and impact, providing a safe, cost-effective alternative to insecticides [56], so understanding the genetic architecture and evolution of traits like climatic adaptation and host specificity is critical. Such research is revealing diverse mechanisms by which parasitoid wasps overcome the host immune system, including venoms, immunosuppressive factors such as polydnviruses and virus-like particles, specialized embryonic cells (teratocytes), and larval secretions [57–59]. Parasitoids can also co-opt host endocrine systems, disrupting host development [60] or manipulating host behavior [61,62]. Recent findings have revealed wasp-microbe associations resulting in evolutionary changes that affect the interactions with their host, suggesting much remains to be discovered. Specifically, turnover of integrated viral genomes within Braconidae and Ichneumonidae wasp genomes are more complex than once anticipated [63,64,65]. Genomes in progress for three parasitoids with a common host have revealed that each has separate mechanisms to overcome host immunity and other defenses (M. Strand, personal communication). Comparative genomics can reveal more information than phylogeny alone as illustrated by the rich collection of research stemming from genomes in the genus *Nasonia* (Pteromalidae; [66] introduces the collection), which also leveraged the extensive history of *Nasonia* genetics research [67]. Comparable research on species from other families will be needed to fully realize the potential of genomics for parasitic Hymenoptera.

Mechanisms and evolution of social behavior are an emphasis of Hymenoptera genomic research. Aculeata are known for sophisticated social behavior within colonies of some ants, bees, and wasps, with eusociality arising multiple times within this clade [68,69]. Glimpses into the molecular basis for this cooperative lifestyle were among the most anticipated findings from the initial sequencing of the honey bee genome [4] while more recent comparative genomic approaches have placed these initial findings within a broader phylogenetic context [8,12,18]. Comparative methods

are particularly useful for understanding the molecular basis for hymenopteran behavior given the impressive diversity of social niches [70,71]. Some insights into eusociality from over 20 eusocial hymenopteran genomes include: 1) the importance of both novel and conserved genes in the regulation of caste phenotypes [19,72], 2) caste-related genes appear to be faster-evolving compared to the rest of the genome suggesting they are targets for adaptive evolution [73,74], and 3) the importance of gene regulation (e.g. changes in gene network structure, epigenetic mechanisms such as DNA methylation) in phenotypic plasticity in eusocial species [13,18].

While most of the focus of hymenopteran genomics has been on social evolution, the Hymenoptera offer many opportunities to further investigate the molecular basis for other aspects of development and animal behavior including maternal care, social parasitism, and foraging/hunting behavior. In some cases, the conceptual framework and predictions are already in place (e.g. social parasitism [75], development [76–81]), others can be enhanced by those generated for other taxonomic groups (e.g. venom evolution in snakes [82], genomic basis of diet shifts [83,84]).

Taxa that are underrepresented in genomics

Given the deep evolutionary distances between major hymenopteran families [85], selecting appropriate taxa for comparison presents a substantial challenge. Crown-group Hymenoptera originated 250–300 mya and spans evolutionary distances more than 3-fold those of modern birds [1,85,86]. Bees shared a common ancestor ~100 mya, and bees + apoid wasps diverged from ants ~145 mya, similar to the split between marsupial and placental mammals (>160 mya) [87]. Increasing the number of taxa sampled within each lineage, as well as the total number of lineages sampled will provide higher resolution to interpret genomic signatures of key phenotypes. In addition, we suggest specific groups that may be particularly appropriate for additional comparisons.

Expanding sampling of Hymenoptera genomes to include one or several genomes per family would provide a useful framework for future evolutionary studies within the order. Although significant recent progress has been made resolving higher-level phylogenetic relationships [85,88,89,90], important uncertainties remain at the superfamily (Figure 1) and family levels, especially outside of the Aculeata. Having genomic data for all 97 families would help overcome common phylogenetic problems, such as insufficient data, incomplete lineage sorting, base composition bias, and long-branch attraction. Multiple carefully chosen representatives of each family would even further reduce issues related to poor taxon sampling [85]. While producing phylogenies with entire genomes is still too expensive for most multi-taxon

studies, having a high quality genome for each family would enable more economical reduced-representation phylogenomics across more taxa [91,92]. Having a complete sampling of Hymenoptera genomes at the family level would also provide a better evaluation of genome size variation and gene family evolution over the entire order. For future developmental biology examinations, selected species should include samples that are readily available for experiments such as embryological and larval time series sampling for gene expression and localization.

Parasitoids are an immensely diverse group for those interested in biological control applications and evolutionary biology [93,94], however the genomes of only 14 species in five families have been published, with draft genomes of ~30 more species (one additional family) in progress (Figure 1, Table 1). Comparative genomics studies of the four families with poly embryonic development would provide insight to this unique development strategy [80]. Representatives of two have been sequenced (Table 1, *M. cingulum* (Braconidae) and *C. floridanum* (Encyrtidae)). Future initiatives should

prioritize at least one polyembryonic species from the Platygasteridae and Dryinidae families. We also recommend at least one species from each parasitoid family, because different families of parasitoids use different hosts taxa with very different biologies. Understanding host specialization and mechanisms parasitoids use to overcome host immunity thus requires broad sampling. Including additional samples to allow comparisons of parasitoids sharing the same host, such as *Drosophila*, and pairing with genomes of the hosts will greatly enable comparative studies across lineages and host/parasite interactions [95].

Comparative genomic approaches to understanding the evolution of eusociality will benefit from increasing the depth of taxonomic coverage within key wasp and bee families, especially the Vespidae, Apidae and Halictidae. Each of these families includes a diversity of social lifestyles, along with closely related solitary individuals. In a few key lineages such as vespine wasps and carpenter bees (Apidae: Xylocopinae), there are also opportunities to compare genomes within lineages in which the entire

Table 1

Hymenoptera genomes in NCBI. Family Formicidae 19 genomes are omitted (reviewed elsewhere).

Family	Genus and Species	Representative Assembly	Reference
Agaonidae	<i>Ceratosolen solmsi</i>	GCF_000503995.1	[106]
Apidae	<i>Apis cerana</i>	GCF_001442555.1	[107]
Apidae	<i>Apis dorsata</i>	GCF_000469605.1	[108]
Apidae	<i>Apis florea</i>	GCF_000184785.2	[12**]
Apidae	<i>Apis mellifera</i> †	GCF_000002195.4	[4,13*]
Apidae	<i>Bombus impatiens</i> †	GCF_000188095.1	[109]
Apidae	<i>Bombus terrestris</i> †	GCF_000214255.1	[109]
Apidae	<i>Ceratina calcarata</i>	GCF_001652005.1	[110]
Apidae	<i>Eufriesea mexicana</i>	GCF_001483705.1	[12**]
Apidae	<i>Euglossa dilemma</i>	GCA_002201625.1	[10]
Apidae	<i>Habropoda laboriosa</i> †	GCF_001263275.1	[12**]
Apidae	<i>Melipona quadrifasciata</i>	GCA_001276565.1	[12**]
Braconidae	<i>Cotesia vestalis</i>	GCA_000956155.1	[111]
Braconidae	<i>Diachasma alloenum</i>	GCF_001412515.1	[112]
Braconidae	<i>Fopius arisanus</i>	GCF_000806365.1	[17]
Braconidae	<i>Macrocentrus cingulum</i>	GCA_002156465.1	[113]
Braconidae	<i>Microplitis demolitor</i>	GCF_000572035.2	[65]
Cephalidae	<i>Cephus cinctus</i>	GCF_000341935.1	[114]
Diprionidae	<i>Neodiprion lecontei</i>	GCF_001263575.1	[115]
Encyrtidae	<i>Copidosoma floridanum</i>	GCF_000648655.1	[116]
Figitidae	<i>Leptopilina clavipes</i>	GCA_001855655.1	[117]
Halictidae	<i>Dufourea novaeangliae</i>	GCF_001272555.1	[12**]
Halictidae	<i>Lasioglossum albipes</i>	GCA_000346575.1	[118]
Megachilidae	<i>Megachile rotundata</i> †	GCF_000220905.1	[12**]
Orussidae	<i>Orussus abietinus</i>	GCF_000612105.1	[119]
Pteromalidae	<i>Nasonia giraulti</i>	GCA_000004775.1	[120*]
Pteromalidae	<i>Nasonia longicornis</i>	GCA_000004795.1	[120*]
Pteromalidae	<i>Nasonia vitripennis</i>	GCF_000002325.3	[120*]
Pteromalidae	<i>Trichomalopsis sarcophagae</i>	GCA_002249905.1	[22*]
Tenthredinidae	<i>Athalia rosae</i>	GCF_000344095.1	[119]
Trichogrammatidae	<i>Trichogramma pretiosum</i>	GCF_000599845.1	[116]
Vespidae	<i>Polistes canadensis</i>	GCF_001313835.1	[18]
Vespidae	<i>Polistes dominula</i>	GCF_001465965.1	[8]

† Important pollinators.

* Genome unpublished but included in collection.

spectrum of sociality occurs, from solitary to weakly social to highly eusocial [71].

Although social wasps have been a model lineage for understanding the evolution of sociality [96], there are only two published genomes for this group, both within the paper wasp genus *Polistes* [8,18]. Current projects to fill these gaps, will deliver genomic insight into the full spectrum of sociality evolution [97•]. The sweat bees (Halictidae) are also particularly useful for comparative studies of social evolution. This family has at least two independent origins of eusociality, at least a dozen lineages that have reverted back to a solitary lifestyle [98–100], and several species in the genera *Halictus*, *Lasioglossum*, and *Megalopta* with facultative expression of eusociality (where individuals or populations vary in solitarily or socially nesting). We propose a strategy of dense sampling at fine phylogenetic scales for key species with variation in social behavior to enhance our understanding of social evolution, providing mechanistic explanations for behavioral responses to environmental factors, such as latitude or elevation [101–103], genetic influences [70•] and *within* population differences in reproductive strategy [104].

Development of genomic resources for taxa from the non-social aculeate wasp families will provide a critical picture of the genomic substrate from which eusociality evolved, and additionally provide insight into the genomic signatures of other behavioral innovations, such as central place foraging (i.e., nesting), maternal care, and social parasitism. Particularly useful families include the spider wasps (Pompilidae), velvet ants (aka ‘cow killers’) (Mutillidae), thread-waisted wasps such as mud daubers (Sphecidae), and cicada killers and bee wolves (Crabronidae, including an independent origin of wasp eusociality). These groups are closely related to vespids and bees, and exhibit remarkable diversity in parental care, dietary breadth, parasitism, and venom function. Additionally, because wasp families are cosmopolitan and conspicuous, their genomes can be studied in the context of well-described behavior and natural history information.

Challenges

Several challenges may impede sequencing and assembly, as well as annotation, of the genomes of Hymenoptera including sample quantity, heterozygosity, and availability; functional interpretation; and unknowns. Though obtaining high quality DNA may be difficult because samples are lacking (especially for parasitoids), for Chalcidoidea, the US National Science Foundation has funded a phylogenetics project that could make available high molecular weight DNA from 388 species in 294 genera in 24 families (John M. Heraty, personal communication). For bees, pinned specimens for many groups may be available from the U.S. National Pollinating Insect Collection at the USDA-ARS-PWA Pollinating Insect

Research Unit. In addition, ARS scientists are routinely collecting new material in North America and could provide genome quality specimens (recently collected and identified) to interested collaborators for some families found in North America. However, some available specimens (including many parasitoids) are very small and yield limited amounts of DNA. Although enough DNA for short-read sequencing library preparation is usually available, extractions from many individuals are needed to provide sufficient DNA for long-read technologies and heterozygosity in these pooled samples contributes to assembly challenges. Theoretically inbreeding for multiple generations can solve this, however some species cannot be inbred (e.g. those with complementary sex determination, in which homozygosity causes mortality or sterile diploid males). Concerning annotation, genes that underlie interesting differences in biology often evolve rapidly, making it difficult to find homologs in species where gene functions are well known. Therefore there is a need for experimentally determined functions (e.g. tissue specific expression and gene knockout/knock in). This is particularly true where differences in expression underlie differences in biology and distant actuators/enhancers and epigenetic modifications, may underlie differences in some key traits (e.g. diapause differences among *Nasonia vitripennis* populations [105•]). Finally, newly sequenced hymenopteran species may have genomic features that reduce genome assembly quality including large size, complex structure and high GC content.

A complete catalog is possible

Although many strategies can be envisioned for future hymenopteran genomics, we suggest a three-pronged strategy focused on breadth, biology and diversity. Increasing the breadth of genome coverage across taxa of this hyper-diverse order will provide long-term scientific benefits. We propose obtaining high quality genomes from at least one representative of each family (81 samples). For the 15 families with more than 50 genera or more than 100 species we propose sampling an additional 0.05% of the species (27 samples, Table 2). For families of particular interest such as pollinators and parasitic wasps, we propose adding 42 species, bringing the total proposed premier genomes to 150 or 0.1% of the species. Premier genomes produced today use long-read sequencing, de novo contig assembly, long-range scaffolding (with Hi-C, BioNano Genomics), and short-read base quality improvement. Such genomes are more contiguous, complete and correct than genomes generated with Sanger or short-read sequencing. Such premier genomes are less expensive than in the past, but still expensive so this may not be possible for all of these samples. Lower cost short read assemblies with high fidelity short contigs and scaffolds or 30x comparative mapping studies can be used to fill in the phylogenetic sampling with additional closely related species. Other useful types of genomic data have been produced (ENCODE) or proposed (FAANG) to

Table 2

Hymenoptera Families with Genera and Species counts and Genomes in NCBI, otherwise published (Pub), in progress (Prog), and proposed here(Prop). Classification citations (Class) follow [2,3] with updates from [47,85,121]. Counts of genera and species are modified from [2] with richness sources cited (Rich). Parasitica* and Symphyta* are paraphyletic. #20 are low coverage genomes.**

	Superfamily	Family	Genomes					Species	Rich	Class		
			Total	NCBI	Pub	Prog	Prop				Genera	
Aculeata/Vespomorpha (stinging wasps)	Thynnoidea	Chyphotidae	-	-	-	-	1	2	65	[123]	[121]	
		Thynnidae	-	-	-	-	2	67	1000	[123]	[121]	
	Tiphioidea	Tiphidae	-	-	-	-	2	53	1000	[2]	[121]	
		Pompiloidea	Pompilidae	-	-	-	-	3	125	4855	[2]	[121]
	Mutillidae		-	-	-	-	2	200	4252	[2]	[121]	
	Sapygidae		-	-	-	-	1	12	66	[2]	[121]	
	Myrmosidae		-	-	-	-	1	10	50	[2]	[121]	
	Sierolomorphae		Sierolomorphidae	-	-	-	-	1	2	11	[2]	[85**]
	Vespoidea	Rhopalosomatidae	-	-	-	-	1	4	72	[2]	[121]	
		Vespidae	14	2	-	12	-	268	4932	[2]	[121]	
	Formicoidea	Formicidae	20	19	1	n/a	n/a	299	12,199	[2]	[121]	
	Apoidea	Ampulicidae	1	-	-	1	-	6	200	[2]	[121]	
		Heterogynidae	-	-	-	-	1	1	8	[124]	[121]	
	Apoidea: Anthophila	Sphecidae	-	-	-	-	1	17	716	[2]	[121]	
		Crabronidae	1	-	-	1	3	242	8773	[2]	[121]	
		Apidae	12	11	-	1	-	209	5749	[2]	[121]	
		Andrenidae	1	-	-	1	-	77	2917	[2]	[121]	
		Megachilidae	3	1	-	2	-	76	4096	[2]	[121]	
		Melittidae	-	-	-	-	1	15	187	[2]	[121]	
		Halictidae	19	2	-	17	-	79	4327	[2]	[121]	
		Stenotritidae	-	-	-	-	1	2	21	[2]	[121]	
		Colletidae	-	-	-	-	2	86	2547	[2]	[121]	
		Scolioidea	Scoliidae	-	-	-	-	1	143	560	[2]	[121]
	Chrysoidea	Bradynobaenidae s.s.	-	-	-	-	1	8	123	[2]	[121]	
		Bethylidae	1	-	-	1	1	84	2340	[2]	[121]	
		Chrysididae	6	-	-	6	-	81	2500	[2]	[121]	
		Dryinidae	-	-	-	-	2	41	1605	[2]	[121]	
		Embolemidae	-	-	-	-	1	2	39	[2]	[121]	
		Plumariidae	-	-	-	-	1	7	22	[2]	[121]	
		Sclerogibbidae	-	-	-	-	1	3	20	[2]	[121]	
		Scolecobryidae	-	-	-	-	1	4	6	[2]	[121]	
	Parasitica* (parasitic wasps)	Megalyroidea	Megaluridae	-	-	-	-	1	8	43	[2]	[3]
	Parasitica* (parasitic wasps)	Trigonalyoidea	Trigonidae	-	-	-	-	1	16	92	[2]	[3]
Evanoidea		Aulacidae	-	-	-	-	1	2	185	[2]	[3]	
		Evaniidae	-	-	-	-	1	21	449	[2]	[3]	
		Gasteruptionidae	-	-	-	-	1	6	496	[2]	[3]	
Stephanoidea		Stephanidae	1	-	-	1	-	11	342	[2]	[3]	
		Chalcidoidea	Eulophidae	1	-	-	1	2	334	4969	[2]	[3]
Pteromalidae			5	4	-	1	-	619	3544	[2]	[3]	
Torymidae			1	-	-	1	-	82	900	[2]	[3]	
Agaonidae			1	1	-	-	-	20	762	[2]	[3]	
Leucospidae			-	-	-	-	1	4	134	[2]	[3]	
Eurytomidae			-	-	-	-	2	97	1453	[2]	[3]	
Azotidae			-	-	-	-	1	1	92	[2]	[2]	
Chalcididae			-	-	-	-	2	90	1469	[2]	[3]	
Cynipencyrtidae			-	-	-	-	1	1	1	[2]	[2]	
Encyrtidae			1	1	-	-	1	493	4058	[2]	[3]	
Eriaporidae			-	-	-	-	1	5	22	[2]	[2]	
Trichogrammatidae			1	1	-	-	-	97	881	[2]	[3]	
Aphelinidae			-	-	-	-	1	29	1078	[2]	[3]	
Eucharitidae			-	-	-	-	1	57	427	[2]	[3]	
Eupelmidae			-	-	-	-	1	51	931	[2]	[3]	
Mymaridae			1	-	-	1	1	96	1437	[2]	[3]	
Ormyridae			-	-	-	-	1	3	125	[2]	[3]	
Perilampidae			-	-	-	-	1	17	284	[2]	[3]	
Rotoitidae			-	-	-	-	1	2	2	[2]	[3]	
Signiphoridae			-	-	-	-	1	4	78	[2]	[3]	
Tanaostigmatidae		-	-	-	-	1	9	93	[2]	[3]		
Tetracampidae		-	-	-	-	1	15	44	[2]	[3]		

Table 2 (Continued)

	Superfamily	Family	Genomes					Species	Rich	Class		
			Total	NCBI	Pub	Prog	Prop				Genera	
Symphyta* (sawflies)	Diapriidae	Diapriidae	23	1	–	21#	–	190	2048	[2]	[3]	
		Ismaridae	–	–	–	–	1	1	29	[2]	[2]	
	Mymarommatoidea	Monomachidae	–	–	–	–	1	2	30	[2]	[3]	
		Maamingidae	–	–	–	–	1	1	2	[2]	[3]	
	Proctotrupoidea	Mymarommatidae	–	–	–	–	1	3	10	[2]	[3]	
		Peleciniidae	–	–	–	–	1	1	3	[2]	[3]	
		Vanhorniidae	–	–	–	–	1	3	5	[2]	[3]	
		Roproniidae	–	–	–	–	1	2	20	[2]	[3]	
		Austroniidae	–	–	–	–	1	1	3	[2]	[2]	
		Heloridae	–	–	–	–	1	1	12	[2]	[3]	
		Peradeniidae	–	–	–	–	1	1	2	[2]	[3]	
		Proctotrupidae	–	–	–	–	1	28	403	[2]	[3]	
		Proctorenyxidae	–	–	–	–	1	2	2	[2]	[3]	
		Platygastroidea	Platygastridae	–	–	–	–	3	236	5385	[2]	[3]
	Cynipoidea	Cynipidae	–	–	–	–	2	74	1412	[2]	[3]	
		Figitidae	1	1	–	–	1	148	1571	[2]	[3]	
	Ichneumonoidea	Liopteridae	–	–	–	–	1	10	153	[2]	[3]	
		Ibaliidae	–	–	–	–	1	3	20	[2]	[3]	
		Austrocynipidae	–	–	–	–	1	1	1	[2]	[3]	
		Braconidae	5	4	–	1	4	1057	19,205	[2]	[3]	
	Ceraphronoidea	Ichneumonidae	3	–	–	3	9	1575	24,025	[2]	[3]	
		Megaspilidae	–	–	–	–	1	15	304	[2]	[3]	
	Orussoidea	Ceraphronidae	–	–	–	–	1	12	299	[2]	[3]	
		Orussidae	1	1	–	–	–	16	82	[2]	[3]	
	Cephoidea	Cephidae	2	1	–	1	–	21	160	[2]	[3]	
	Xiphidriidea	Xiphidriidae	–	–	–	–	1	28	146	[2]	[3]	
	Siricoidea	Siricidae	–	–	–	1	1	11	111	[2]	[3]	
		Anaxyelidae	–	–	–	–	1	1	1	[2]	[3]	
	Xyeloidea	Xyelidae	1	–	–	1	–	5	63	[2]	[3]	
	Tenthredinoidea	Argidae	–	–	–	–	1	58	897	[2]	[3]	
		Blasticotomidae	–	–	–	–	1	2	12	[2]	[3]	
		Cimbicidae	–	–	–	–	1	16	182	[2]	[3]	
		Diprionidae	–	–	–	–	1	11	136	[2]	[3]	
		Heptamelidae	–	–	–	–	1	2	38	[125,126]	[3]	
		Pergidae	–	–	–	–	1	60	442	[2]	[3]	
		Tenthredinidae	1	1	–	–	2	398	5462	[2]	[3]	
		Pamphilioidea	Pamphiliidae	–	–	–	–	1	1	42	[2]	[3]
			Megalodontesidae	–	–	–	–	1	10	291	[2]	[3]
		Family counts			26	15	1	20	81			
	Sum totals			128	52	1	75#	108	8422	152,658		

annotate other animals. Including these data or at least transcript sequencing (with high quality long read data) is a cost effective adjunct to enhance the value of lower quality assemblies. With available samples, robust technologies and more reliable methods, this is a feasible task with the potential to impact many fields of study for years into the future.

Acknowledgements

Funding for this work was provided by the NIH NHGRI Large-Scale Sequencing Program grant to Richard Gibbs (U54 HG003273); KK is supported by Utah State University and project 1297 to the Utah Agricultural Experiment Station, AT is supported by NSF-IOS # 1456283 (Behavioral Systems Cluster). We thank Alexander Wild for photographs. We thank Gerald Thomas Wright for production assistance. We thank researchers for sharing their genome sequencing plans with us for this publication. USDA is an equal opportunity provider and employer.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.cois.2017.11.008](https://doi.org/10.1016/j.cois.2017.11.008).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Grimaldi D, Engel MS: *Evolution of the Insects*. Cambridge University Press; 2005.
2. Aguiar AP, Deans AR, Engel MS, Forshage M, Huber JT, Jennings JT, Johnson NF, Lelej AS, Longino JT, Lohrmann V *et al.*: **Order Hymenoptera**. In *Animal Biodiversity: An Outline of Higher-level Classification and Survey of Taxonomic Richness (Addenda 2013)*, vol 3703. Edited by Zhang Z-Q. Zootaxa; 2013:51.

3. Sharkey MJ: **Phylogeny and classification of Hymenoptera.** *Zootaxa* 2007, **1668**:521-548.
4. Honeybee Genome Sequencing Consortium: **Insights into social insects from the genome of the honeybee *Apis mellifera*.** *Nature* 2006, **443**:931-949.
5. Heimpel GE, de Boer JG: **Sex determination in the hymenoptera.** *Annu Rev Entomol* 2008, **53**:209-230.
6. Evans JD, Shearman DC, Oldroyd BP: **Molecular basis of sex determination in haplodiploids.** *Trends Ecol Evol* 2004, **19**:1-3.
7. Trivers RL, Hare H: **Haplodiploidy and the evolution of the social insect.** *Science* 1976, **191**:249-263.
8. Standage DS, Berens AJ, Glastad KM, Severin AJ, Brendel VP, Toth AL: **Genome, transcriptome and methylome sequencing of a primitively eusocial wasp reveal a greatly reduced DNA methylation system in a social insect.** *Mol Ecol* 2016, **25**:1769-1784.
9. Wilfert L, Gadau J, Schmid-Hempel P: **Variation in genomic recombination rates among animal taxa and the case of social insects.** *Heredity (Edinb)* 2007, **98**:189-197.
10. Brand P, Saleh N, Pan H, Li C, Kapheim KM, Ramirez SR: **The nuclear and mitochondrial genomes of the facultatively eusocial orchid bee euglossa dilemma.** *G3 (Bethesda)* 2017, **7**:2891-2898.
11. Ardila-Garcia AM, Umphrey GJ, Gregory TR: **An expansion of the genome size dataset for the insect order Hymenoptera, with a first test of parasitism and eusociality as possible constraints.** *Insect Mol Biol* 2010, **19**:337-346.
12. Kapheim KM, Pan H, Li C, Salzberg SL, Puiu D, Magoc T, Robertson HM, Hudson ME, Venkat A, Fischman BJ *et al.*: **Social evolution, genomic signatures of evolutionary transitions from solitary to group living.** *Science* 2015, **348**:1139-1143.
- Most of the 20,000 bee species live solitary lifestyles, but group living has evolved independently at least three times in the Apidae and Halictidae. These families represent a striking amount of social diversity that spans a broad range of group sizes, degree of cooperation, and adaptations to colony life. This study comparing genomes from both solitary and social bees from these families suggested that changes in gene regulatory function accompany evolutionary transitions among social lifestyles, but that the specifics of these changes are largely lineage specific.
13. Elsik CG, Worley KC, Bennett AK, Beye M, Camara F, Childers CP, de Graaf DC, Debyser G, Deng J, Devreese B *et al.*: **Finding the missing honey bee genes: lessons learned from a genome upgrade.** *BMC Genomics* 2014, **15**:86.
- Increases in genome assembly contiguity and completeness, along with transcriptome evidence for gene prediction, led to a 50% larger honey bee gene set and aided the detection of potential lineage-specific genes. Relative contributions of each evidence type on the composition of the final official gene set are discussed.
14. Kent CF, Minaei S, Harpur BA, Zayed A: **Recombination is associated with the evolution of genome structure and worker behavior in honey bees.** *Proc Natl Acad Sci U S A* 2012, **109**:18012-18017.
15. Richards S, Murali SC: **Best practices in insect genome sequencing: what works and what doesn't.** *Curr Opin Insect Sci* 2015, **7**:1-7.
- This review discusses issues of insect genome sequencing including the opportunities provided by new technologies and computational methods as well as genomic and biological features of insects that impact genome sequencing projects.
16. Cox-Foster DL, Conlan S, Holmes EC, Palacios G, Evans JD, Moran NA, Quan PL, Briese T, Hornig M, Geiser DM *et al.*: **A metagenomic survey of microbes in honey bee colony collapse disorder.** *Science* 2007, **318**:283-287.
17. Geib SM, Liang GH, Murphy TD, Sim SB: **Whole genome sequencing of the braconid parasitoid wasp fopius arisanus, an important biocontrol agent of pest tepritid fruit flies.** *G3 (Bethesda)* 2017, **7**:2407-2411.
18. Patalano S, Vlasova A, Wyatt C, Ewels P, Camara F, Ferreira PG, Asher CL, Jurkowski TP, Segonds-Pichon A, Bachman M *et al.*: **Molecular signatures of plastic phenotypes in two eusocial insect species with simple societies.** *Proc Natl Acad Sci U S A* 2015, **112**:13970-13975.
19. Smith CR, Helms Cahan S, Kemena C, Brady SG, Yang W, Bornberg-Bauer E, Eriksson T, Gadau J, Helmkamp M, Gotzek D *et al.*: **How do genomes create novel phenotypes? insights from the loss of the worker caste in ant social parasites.** *Mol Biol Evol* 2015, **32**:2919-2931.
20. Wallberg A, Pirk CW, Allsopp MH, Webster MT: **Identification of multiple loci associated with social parasitism in honeybees.** *PLoS Genet* 2016, **12**:e1006097.
- Cape bees reproduce asexually and exhibit phenotypes that facilitate social parasitism. This comparison to other African bees identified divergent regions of the genome with signatures of positive selection that likely underlie the genetic basis of social parasitism.
21. Martinson EO, Martinson VG, Edwards R, Mrinalini, Werren JH: **Laterally transferred gene recruited as a venom in parasitoid wasps.** *Mol Biol Evol* 2016, **33**:1042-1052.
22. Martinson EO, Mrinalini, Kelkar YD, Chang CH, Werren JH: **The evolution of venom by co-option of single-copy genes.** *Curr Biol* 2017, **27**:2007-2013 e2008.
- Rapid turnover of venom genes was used to study how new gene functions evolve. Co-option of single copy genes via evolution of cis-regulatory expression may be an underappreciated mechanism of rapid evolutionary change.
23. Desneux N, Barta RJ, Hoelmer KA, Hopper KR, Heimpel GE: **Multifaceted determinants of host specificity in an aphid parasitoid.** *Oecologia* 2009, **160**:387-398.
24. Robertson HM, Gadau J, Wanner KW: **The insect chemoreceptor superfamily of the parasitoid jewel wasp *Nasonia vitripennis*.** *Insect Mol Biol* 2010, **19**:121-136.
- Analysis of the published *Nasonia vitripennis* genome revealed 47 functional gustatory receptor genes and 225 functional odorant receptor genes. *Nasonia* may need this diversity of chemoreceptor genes to find mates and hosts or avoid harmful substances.
25. Badaoui B, Fougeroux A, Petit F, Anselmo A, Gorni C, Cucurachi M, Cersini A, Granato A, Cardeti G, Formato G *et al.*: **RNA-sequence analysis of gene expression from honeybees (*Apis mellifera*) infected with *Nosema ceranae*.** *PLOS ONE* 2017, **12**:e0173438.
26. Dolezal AG, Toth AL: **Honey bee sociogenomics: a genome-scale perspective on bee social behavior and health.** *Apidologie* 2014, **45**:375-395.
27. Doublet V, Poeschl Y, Gogol-Doring A, Alaux C, Annoscia D, Aurori C, Barribeau SM, Bedoya-Reina OC, Brown MJ, Bull JC *et al.*: **Unity in defence: honeybee workers exhibit conserved molecular responses to diverse pathogens.** *BMC Genomics* 2017, **18**:207.
28. Grozinger CM, Robinson GE: **The power and promise of applying genomics to honey bee health.** *Curr Opin Insect Sci* 2015, **10**:124-132.
29. Zanni V, Galbraith DA, Annoscia D, Grozinger CM, Nazzi F: **Transcriptional signatures of parasitization and markers of colony decline in Varroa-infested honey bees (*Apis mellifera*).** *Insect Biochem Mol Biol* 2017, **87**:1-13.
30. Torson AS, Yocum GD, Rinehart JP, Kemp WP, Bowsher JH: **Transcriptional responses to fluctuating thermal regimes underpinning differences in survival in the solitary bee *Megachile rotundata*.** *J Exp Biol* 2015, **218**:1060-1068.
31. Torson AS, Yocum GD, Rinehart JP, Nash SA, Kvidera KM, Bowsher JH: **Physiological responses to fluctuating temperatures are characterized by distinct transcriptional profiles in a solitary bee.** *J Exp Biol* 2017, **220**:3372-3380.
32. Lozier JD, Zayed A: **Bee conservation in the age of genomics.** *Conserv Genet* 2016, **18**:713-729.
33. Trapp J, McAfee A, Foster LJ: **Genomics, transcriptomics and proteomics: enabling insights into social evolution and disease challenges for managed and wild bees.** *Mol Ecol* 2017, **26**:718-739.

34. Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R: **Climate-associated phenological advances in bee pollinators and bee-pollinated plants.** *Proc Natl Acad Sci U S A* 2011, **108**:20645-20649.
35. Budge GE, Garthwaite D, Crowe A, Boatman ND, Delaplane KS, Brown MA, Thygesen HH, Pietravalle S: **Evidence for pollinator cost and farming benefits of neonicotinoid seed coatings on oilseed rape.** *Sci Rep* 2015, **5**:12574.
36. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE: **Global pollinator declines: trends, impacts and drivers.** *Trends Ecol Evol* 2010, **25**:345-353.
37. Whitehorn PR, O'Connor S, Wackers FL, Goulson D: **Neonicotinoid pesticide reduces bumble bee colony growth and queen production.** *Science* 2012, **336**:351-352.
38. Ziska LH, Pettis JS, Edwards J, Hancock JE, Tomecek MB, Clark A, Duker JS, Loladze I, Polley HW: **Rising atmospheric CO₂ is reducing the protein concentration of a floral pollen source essential for North American bees.** *Proc Biol Sci* 2016:283.
39. Kerr JT, Pindar A, Galpern P, Packer L, Potts SG, Roberts SM, Rasmont P, Schweiger O, Colla SR, Richardson LL *et al.*: **Relocation risky for bumblebee colonies – response.** *Science* 2015, **350**:287.
40. Annoscia D, Zanni V, Galbraith D, Quirici A, Grozinger C, Bortolomeazzi R, Nazzi F: **Elucidating the mechanisms underlying the beneficial health effects of dietary pollen on honey bees (*Apis mellifera*) infested by Varroa mite ectoparasites.** *Sci Rep* 2017, **7**:6258.
41. Corby-Harris V, Jones BM, Walton A, Schwan MR, Anderson KE: **Transcriptional markers of sub-optimal nutrition in developing *Apis mellifera* nurse workers.** *BMC Genomics* 2014, **15**:134.
42. Wang H, Zhang S-W, Zeng Z-J, Yan W-Y: **Nutrition affects longevity and gene expression in honey bee (*Apis mellifera*) workers.** *Apidologie* 2014, **45**:618-625.
43. Wheeler MM, Robinson GE: **Diet-dependent gene expression in honey bees: honey vs. sucrose or high fructose corn syrup.** *Sci Rep* 2014, **4**:5726.
44. Brunner FS, Schmid-Hempel P, Barribeau SM: **Protein-poor diet reduces host-specific immune gene expression in *Bombus terrestris*.** *Proc Biol Sci* 2014:281.
45. Barribeau SM, Sadd BM, du Plessis L, Brown MJ, Buechel SD, Cappelle K, Carolan JC, Christiaens O, Colgan TJ, Eriş S *et al.*: **A depauperate immune repertoire precedes evolution of sociality in bees.** *Genome Biol* 2015, **16**:83.
46. Lopez-Urbe MM, Sconiers WB, Frank SD, Dunn RR, Tarpy DR: **Reduced cellular immune response in social insect lineages.** *Biol Lett* 2016, **12**:20150984.
47. Heraty JM, Burks RA, Craud A, Gibson GAP, Liljeblad J, Munro J, Rasplus J-Y, Delvare G, Janta P, Gumovsky A *et al.*: **A phylogenetic analysis of the megadiverse Chalcidoidea (Hymenoptera).** *Cladistics* 2013, **29**:466-542.
48. Gauld ID, Janzen DH: **The systematics and biology of the Costa Rican species of parasitic wasps in the Thyreodon genus-group (Hymenoptera: Ichneumonidae).** *Zool J Linn Soc* 2004, **141**:297-351.
49. Kazmer DJ, Kazmer DJ, Maiden K, Ramualde N, Coutinot D: **Reproductive compatibility, mating behavior, and random amplified polymorphic dna variability in some *Aphelinus asychis* (Hymenoptera: Aphelinidae) derived from the old world.** *Ann Entomol Soc Am* 1996, **89**:212-220.
50. Pinto JD, Platner GR, Stouthamer R: **The systematics of the *Trichogramma minutum* species complex (Hymenoptera: Trichogrammatidae), a group of important North American biological control agents: the evidence from reproductive compatibility and allozymes.** *Biol Control* 2003, **27**:167-180.
51. Stireman JO, Nason JD, Heard SB, Seehawer JM: **Cascading host-associated genetic differentiation in parasitoids of phytophagous insects.** *Proc Biol Sci* 2006, **273**:523-530.
52. Godfray H: *Parasitoids: Behavioral and Evolutionary Ecology.* Princeton, NJ: Princeton University Press; 1994.
53. Strand MR, Obyrck JJ: **Host specificity of insect parasitoids and predators. Many factors influence the host ranges of insect natural enemies.** *BioScience* 1996, **46**:422-429.
54. Forbes AA, Powell TH, Stelinski LL, Smith JJ, Feder JL: **Sequential sympatric speciation across trophic levels.** *Science* 2009, **323**:776-779.
55. Schluter D: **Evidence for ecological speciation and its alternative.** *Science* 2009, **323**:737-741.
56. Cock MJW, Murphy ST, Kairo MTK, Thompson E, Murphy RJ, Francis AW: **Trends in the classical biological control of insect pests by insects: an update of the BIOCAT database.** *BioControl* 2016, **61**:349-363.
57. Burke GR, Strand MR: **Systematic analysis of a wasp parasitism arsenal.** *Mol Ecol* 2014, **23**:890-901.
58. Moreau SJ, Asgari S: **Venom proteins from parasitoid wasps and their biological functions.** *Toxins (Basel)* 2015, **7**:2385-2412.
59. Vass E, Nappi AJ: **Developmental and immunological aspects of *Drosophila*-parasitoid relationships.** *J Parasitol* 2000, **86**:1259-1270.
60. Beckage NE, Gelman DB: **Wasp parasitoid disruption of host development: implications for new biologically based strategies for insect control.** *Annu Rev Entomol* 2004, **49**:299-330.
61. Dheilly NM, Maure F, Ravallec M, Galinier R, Doyon J, Duval D, Leger L, Volkoff AN, Misse D, Nidelet S *et al.*: **Who is the puppet master? Replication of a parasitic wasp-associated virus correlates with host behaviour manipulation.** *Proc Biol Sci* 2015, **282**:20142773.
62. Mohan P, Sinu PA: **Parasitoid wasp usurps its host to guard its pupa against hyperparasitoids and induces rapid behavioral changes in the parasitized host.** *PLOS ONE* 2017, **12**:e0178108.
63. Drezen JM, Gauthier J, Josse T, Bezier A, Herniou E, Huguet E: **Foreign DNA acquisition by invertebrate genomes.** *J Invertebr Pathol* 2017, **147**:157-168.
64. Drezen JM, Leobold M, Bezier A, Huguet E, Volkoff AN, • Herniou EA: **Endogenous viruses of parasitic wasps: variations on a common theme.** *Curr Opin Virol* 2017, **25**:41-48.
- Review of endogenous viruses and their co-option by parasitic wasps for parasitism and reproductive success. Sequencing additional parasitic wasp genomes will shed more light on the co-evolution of parasitic wasps with these 'domesticated' viruses.
65. Burke GR, Walden KK, Whitfield JB, Robertson HM, Strand MR: **Widespread genome reorganization of an obligate virus mutualist.** *PLoS Genet* 2014, **10**:e1004660.
66. Brown S: **The *Nasonia* genome sequence: finding gems in the jewel (wasp) box.** *Insect Mol Biol* 2010, **19** v-vii.
67. Werren JH, Loehlin DW: **The parasitoid wasp *Nasonia*: an emerging model system with haploid male genetics.** *Cold Spring Harb Protoc* 2009, **2009** pdb emo134.
68. Bank S, Sann M, Mayer C, Meusemann K, Donath A, Podsiadlowski L, Kozlov A, Petersen M, Krogmann L, Meier R *et al.*: **Transcriptome and target DNA enrichment sequence data provide new insights into the phylogeny of vespid wasps (Hymenoptera: Aculeata: Vespidae).** *Mol Phylogenet Evol* 2017, **116**:213-226.
69. Bradley TJ, Briscoe AD, Brady SG, Contreras HL, Danforth BN, Dudley R, Grimaldi D, Harrison JF, Kaiser JA, Merlin C *et al.*: **Episodes in insect evolution.** *Integr Comp Biol* 2009, **49**:590-606.
70. Kocher SD, Paxton RJ: **Comparative methods offer powerful insights into social evolution in bees.** *Apidologie* 2014, **45**:289-305.
- Review of the diversity of social behavior displayed from bees, including four independent origins of eusociality and reversals to solitary behavior in some lineages, which provides a powerful framework for comparative work on the evolution of this trait.
71. Rehan SM, Toth AL: **Climbing the social ladder: the molecular evolution of sociality.** *Trends Ecol Evol* 2015, **30**:426-433.

72. Berens AJ, Hunt JH, Toth AL: **Comparative transcriptomics of convergent evolution: different genes but conserved pathways underlie caste phenotypes across lineages of eusocial insects.** *Mol Biol Evol* 2015, **32**:690-703.

First study to compare differentially expressed transcripts between castes in three separate eusocial hymenopteran lineages. Convergent evolution for the social caste system was not found to utilize a common set of genes, nor was lineage specific genes. Instead the authors found that similar gene networks and molecular functions underlie this key social trait.

73. Harpur BA, Kent CF, Molodtsova D, Lebon JM, Alqarni AS, Owayss AA, Zayed A.: **Population genomics of the honey bee reveals strong signatures of positive selection on worker traits.** *Proc Natl Acad Sci U S A* 2014, **111**:2614-2619.
74. Johnson BR, Tsutsui ND: **Taxonomically restricted genes are associated with the evolution of sociality in the honey bee.** *BMC Genomics* 2011, **12**:164.
75. Cini A, Patalano S, Segonds-Pichon A, Busby GB, Cervo R, Sumner S: **Social parasitism and the molecular basis of phenotypic evolution.** *Front Genet* 2015, **6**:32.
76. Corona M, Libbrecht R, Wheeler DE: **Molecular mechanisms of phenotypic plasticity in social insects.** *Curr Opin Insect Sci* 2016, **13**:55-60.
77. Cridge AG, Lovegrove MR, Skelly JG, Taylor SE, Petersen GEL, Cameron RC, Dearden PK: **The honeybee as a model insect for developmental genetics.** *Genesis* 2017:55.
78. Lynch JA: **The expanding genetic toolbox of the wasp *Nasonia vitripennis* and its relatives.** *Genetics* 2015, **199**:897-904.
79. Sehnaal F, Svacha P, Zrzavy J: **Evolution of insect metamorphosis.** In *Metamorphosis. Postembryonic Reprogramming of Gene Expression in Amphibian and Insect Cells*. Edited by Gilbert L, Tata J, Atkinson B. Academic Press; 1996:3-58.
80. Smith MS, Milton I, Strand MR: **Phenotypically plastic traits regulate caste formation and soldier function in polyembryonic wasps.** *J Evol Biol* 2010, **23**:2677-2684.
81. Zhurov V, Terzin T, Grbic M: **(In)discrete charm of the polyembryony: evolution of embryo cloning.** *Cell Mol Life Sci* 2007, **64**:2790-2798.
82. Casewell NR, Wuster W, Vonk FJ, Harrison RA, Fry BG: **Complex cocktails: the evolutionary novelty of venoms.** *Trends Ecol Evol* 2013, **28**:219-229.
83. Babbitt CC, Warner LR, Fedrigo O, Wall CE, Wray GA: **Genomic signatures of diet-related shifts during human origins.** *Proc Biol Sci* 2011, **278**:961-969.
84. Groen SC, Whiteman NK: **Using *Drosophila* to study the evolution of herbivory and diet specialization.** *Curr Opin Insect Sci* 2016, **14**:66-72.
85. Branstetter MG, Danforth BN, Pitts JP, Faircloth BC, Ward PS, Buffington ML, Gates MW, Kula RR, Brady SG: **Phylogenomic insights into the evolution of stinging wasps and the origins of ants and bees.** *Curr Biol* 2017, **27**:1019-1025.

One of the first (see also 90 below) large-scale phylogenomic studies of the stinging wasps. Ultra-Conserved Element (UCE) enrichment methods were used to sample 854 UCE loci from 187 taxa, including 30 of 31 aculeate families. The results produced a highly supported phylogeny of stinging wasps, recovering ants as the sister group to apooid wasps plus bees. Taxon sampling experiments also demonstrated the significant impact taxon choice can have on the phylogeny, even when using genomic data.

86. Prum RO, Berv JS, Dornburg A, Field DJ, Townsend JP, Lemmon EM, Lemmon AR: **A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing.** *Nature* 2015, **526**:569-573.
87. Williamson TE, Brusatte SL, Wilson GP: **The origin and early evolution of metatherian mammals: the Cretaceous record.** *Zookeys* 2014:1-76.
88. Johnson BR, Borowiec ML, Chiu JC, Lee EK, Atallah J, Ward PS: **Phylogenomics resolves evolutionary relationships among ants, bees, and wasps.** *Curr Biol* 2013, **23**:2058-2062.

89. Misof B, Liu S, Meusemann K, Peters RS, Donath A, Mayer C, Frandsen PB, Ware J, Flouri T, Beutel RG *et al.*: **Phylogenomics resolves the timing and pattern of insect evolution.** *Science* 2014, **346**:763-767.

90. Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, Podsiadlowski L, Petersen M, Lanfear R *et al.*: **Evolutionary history of the Hymenoptera.** *Curr Biol* 2017, **27**:1013-1018.

This report developed a time-calibrated and statistically solid phylogenetic tree of Hymenoptera from analyzing more than 3000 protein coding genes in 173 insect species, derived primarily from transcriptome data collected by the 1KITE consortium. Initial Hymenoptera diversification placed at ~281 mya. Sawflies and parasitoid wasps were each found to be monophyletic, Trigonalioidea were identified as the sister group to Aculeata and the evolutionary root of bees was located within the Crabronidae. This understanding of the evolutionary history of Hymenoptera allows for a clear classification system and provides a strong foundation from which to choose species for comparative efforts across a wide range of interesting traits.

91. Branstetter MG, Longino JT, Ward PS, Faircloth BC, Price S: **Enriching the ant tree of life: enhanced UCE bait set for genome-scale phylogenetics of ants and other Hymenoptera.** *Methods Ecol Evol* 2017, **8**:768-776.
92. Lemmon EM, Lemmon AR: **High-throughput genomic data in systematics and phylogenetics.** *Annu Rev Ecol Syst* 2013, **44**:99-121.
93. Kuhn KL, Duan JJ, Hopper KR: **Next-generation genome sequencing and assembly provides tools for phylogenetics and identification of closely related species of *Spathius*, parasitoids of *Agrilus planipennis* (emerald ash borer).** *Biol Control* 2013, **66**:77-82.
94. Yao Y-X, Duan JJ, Hopper KP, Mottern JL, Gates MW: **A new species of *Oobius* Trjapitzin (Hymenoptera: Encyrtidae) from the Russian far east that parasitizes eggs of emerald ash borer (Coleoptera: Buprestidae).** *Ann Entomol Soc Am* 2016, **109**:629-638.
95. Salazar-Jaramillo L, Jalvingh KM, de Haan A, Kraaijeveld K, Buermans H, Wertheim B: **Inter- and intra-species variation in genome-wide gene expression of *Drosophila* in response to parasitoid wasp attack.** *BMC Genomics* 2017, **18**:331.
96. Jandt JM, Bengston S, Pinter-Wollman N, Pruitt JN, Raine NE, Dornhaus A, Sih A: **Behavioural syndromes and social insects: personality at multiple levels.** *Biol Rev Camb Philos Soc* 2014, **89**:48-67.
97. Toth AL, Rehan SM: **Molecular evolution of insect sociality: an eco-evo-devo perspective.** *Annu Rev Entomol* 2017, **62**:419-442.
- This synthesis of the comparative and evolutionary genomics research in eusocial insects reviews numerous independently proposed mechanisms for the evolution of eusociality and considers them in a holistic framework. Areas requiring additional data and consideration include solitary and incipiently social species within each of the multiple independent origins of sociality among Hymenoptera in order to investigate the early stages of social evolution.
98. Brady SG, Sipes S, Pearson A, Danforth BN: **Recent and simultaneous origins of eusociality in halictid bees.** *Proc Biol Sci* 2006, **273**:1643-1649.
99. Gibbs J, Brady SG, Kanda K, Danforth BN: **Phylogeny of halictine bees supports a shared origin of eusociality for Halictus and Lasioglossum (Apoidea: Anthophila: Halictidae).** *Mol Phylogenet Evol* 2012, **65**:926-939.
100. Wcislo WT, Danforth BN: **Secondarily solitary: the evolutionary loss of social behavior.** *Trends Ecol Evol* 1997, **12**:468-474.
101. Davison PJ, Field J: **Social polymorphism in the sweat bee *Lasioglossum (Evylaeus) calceatum*.** *Insectes Soc* 2016, **63**:327-338.
102. Field J, Paxton RJ, Soro A, Bridge C: **Cryptic plasticity underlies a major evolutionary transition.** *Curr Biol* 2010, **20**:2028-2031.
103. Packer L: **Solitary and eusocial nests in a population of *Augochlorella striata* (Provancher) (Hymenoptera; Halictidae) at the northern edge of its range.** *Behav Ecol Sociobiol* 1990, **27**:339-344.

104. Kapheim KM, Smith AR, Nonacs P, Wcislo WT, Wayne RK: **Foundress polyphenism and the origins of eusociality in a facultatively eusocial sweat bee, *Megalopta genalis* (Halictidae)**. *Behav Ecol Sociobiol* 2013, **67**:331-340.
105. Pegoraro M, Bafna A, Davies NJ, Shuker DM, Tauber E: **DNA methylation changes induced by long and short photoperiods in *Nasonia***. *Genome Res* 2016, **26**:203-210.
 Female *Nasonia vitripennis* control diapause of their offspring by the pattern of DNA methylation of offspring genomes, and knockout of methylation genes disrupts the photoperiodic diapause response. This is the first example of DNA methylation affecting insect photoperiodic timing.
106. Xiao JH, Yue Z, Jia LY, Yang XH, Niu LH, Wang Z, Zhang P, Sun BF, He SM, Li Z *et al.*: **Obligate mutualism within a host drives the extreme specialization of a fig wasp genome**. *Genome Biol* 2013, **14**:R141.
107. Park D, Jung JW, Choi BS, Jayakodi M, Lee J, Lim J, Yu Y, Choi YS, Lee ML, Park Y *et al.*: **Uncovering the novel characteristics of Asian honey bee, *Apis cerana*, by whole genome sequencing**. *BMC Genomics* 2015, **16**:1.
108. Rueppell O, *et al.*: *Apis dorsata* Genome. In preparation.
109. Sadd BM, Barribeau SM, Bloch G, de Graaf DC, Dearden P, Elsik CG, Gadau J, Grimmelikhuijzen CJ, Hasselmann M, Lozier JD *et al.*: **The genomes of two key bumblebee species with primitive eusocial organization**. *Genome Biol* 2015, **16**:76.
110. Rehan SM, Glastad KM, Lawson SP, Hunt BG: **The genome and methylome of a subsocial small carpenter bee, *Ceratina calcarata***. *Genome Biol Evol* 2016, **8**:1401-1410.
111. Kim Y, Lee D, Kim Y: **Mutualism of polydnavirus with host wasp: an insight of evolutionary trajectory from phytophagous to parasitoid Hymenoptera**. In Preparation.
112. Forbes A, Robertson HM, Walden KK, Hood GR, Feder JL: **Draft genome of *Diachasma alloeum***. In Preparation.
113. Yin C, Hu J, Li F: **Genome analysis enriches our understanding of wasp parasitic tactics**. In preparation.
114. Wanner KW, Robertson H: ***Cephus cinctus* genome**. In Preparation.
115. Linnen C: ***Neodiprion lecontei* genome**. In preparation.
116. Richards S, *et al.*: **The 15k Pilot Initiative**. In preparation.
117. Kraaijeveld K, Anvar SY, Frank J, Schmitz A, Bast J, Wilbrandt J, Petersen M, Ziesmann T, Niehuis O, de Knijff P *et al.*: **Decay of sexual trait genes in an asexual parasitoid wasp**. *Genome Biol Evol* 2016, **8**:3685-3695.
118. Kocher SD, Li C, Yang W, Tan H, Yi SV, Yang X, Hoekstra HE, Zhang G, Pierce NE, Yu DW: **The draft genome of a socially polymorphic halictid bee, *Lasioglossum albipes***. *Genome Biol* 2013, **14**:R142.
119. Hatakeyama M: ***Athalia rosae* and *Orussus abietinus* genomes**. In preparation, 15k.
120. Werren JH, Richards S, Desjardins CA, Niehuis O, Gadau J, Colbourne JK, Nasonia Genome Working G, Werren JH, Richards S, Desjardins CA *et al.*: **Functional and evolutionary insights from the genomes of three parasitoid *Nasonia* species**. *Science* 2010, **327**:343-348.
 Genome sequences and comparative genomic analyses of three closely related parasitoids (*Nasonia vitripennis*, *N. giraulti*, and *N. longicornis*), which were the first annotated genomes for parasitoids. Analyses revealed diverse venom genes, a set of DNA methylation genes, lateral transfers of *Wolbachia* among *Nasonia* species, and rapid evolution of genes involved in nuclear-mitochondrial interactions that may mediate speciation.
121. Pilgrim EM, von Dohlen CD, Pitts JP: **Molecular phylogenetics of Vespoidea indicate paraphyly of the superfamily and novel relationships of its component families and subfamilies**. *Zool Scripta* 2008, **37**:539-560.
122. Goulet H, Huber JT (Eds): *Hymenoptera of the World: An Identification Guide to Families*. Ottawa, Ontario, Canada: Agriculture Canada; 1993.
123. Pitts JP. personal communication.
124. Pulawski WJ (Ed): *Catalog of Sphecidae sensu lato*. 2017. Available from: <https://www.calacademy.org/scientists/projects/catalog-of-sphecidae> (accessed 30.11.17).
125. Vikberg V, Liston A: **Taxonomy and biology of European Heptamelini (Hymenoptera, Tenthredinidae, Selandriinae)**. *Zootaxa* 2009, **2112**:1-24.
126. Malm T, Nyman T: **Phylogeny of the symphytan grade of Hymenoptera: new pieces into the old jigsaw(fly) puzzle**. *Cladistics* 2015, **31**:1-17.