

1997

Rapid Communication: A Restriction Fragment Length Polymorphism in the Ovine Prolactin Gene

Amy L. Vincent

Iowa State University, amyv@iastate.edu

Max F. Rothschild

Iowa State University, mfrothsc@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/ans_pubs

 Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), and the [Genetics and Genomics Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/ans_pubs/304. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Rapid Communication: A Restriction Fragment Length Polymorphism in the Ovine Prolactin Gene

Abstract

Polymorphism. A HaeIII PCR-RFLP was identified in the ovine prolactin (PRL) gene. Source and Description of Primers. Human genomic (Truong et al., 1984) and pig cDNA (GenBank accession no. X14068) sequences were compared to design primers to span the second intron of the prolactin gene.

Keywords

Sheep, PCR-RLFP, Prolactin Gene

Disciplines

Agriculture | Animal Sciences | Genetics and Genomics

Comments

This is an article from *Journal of Animal Science* 75 (1997): 1686, doi:/1997.7561686x. Posted with permission.

Rapid Communication: A Restriction Fragment Length Polymorphism in the Ovine *Prolactin* Gene¹

A. L. Vincent and M. F. Rothschild²

Department of Animal Science, Iowa State University, Ames 50011

Polymorphism. A *Hae*III PCR-RFLP was identified in the ovine *prolactin* (*PRL*) gene.

Source and Description of Primers. Human genomic (Truong et al., 1984) and pig cDNA (GenBank accession no. X14068) sequences were compared to design primers to span the second intron of the prolactin gene.

Primer Sequences. Forward primer: 5'-ACCTCTCTTCGAAATGTTCA-3'; reverse primer: 5'-CTGTTGGGCTTGCTCTTTGTC-3'.

Method of Detection. PCR amplification (25 μ L final volume) was performed using 25 ng of genomic DNA, 350 μ M each dNTP, .3 μ M each primer, 1.1 mM MgCl₂, 1 \times *Taq* Extender Buffer, 1 unit *Taq* Extender, and 1 unit *Taq* polymerase. The thermal cycler profile was 92°C for 2 min; 35 cycles of 92°C for 45 s, 56°C for 45 s, and 72°C for 3 min, followed by a final extension at 72°C for 7 min.

Description of Polymorphism. Twenty microliters of the 2.5-kb product was digested with *Hae*III to produce bands of approximately 1,400, 530, 360, and 150 bp for the A allele and bands of approximately 1,400, 510, 360, and 150 bp for the B allele (Figure 1).

Inheritance Pattern. The *PRL* *Hae*III polymorphism was observed to have a Mendelian inheritance pattern in nine two-generation sheep families of the AgResearch International Mapping Flock (Crawford et al., 1995).

Frequency. Frequencies for the A allele were .30 for crossbred whitefaced (n = 10), .25 for Suffolk (n = 6), .375 for Coopworth (n = 4), and .50 for Texel (n = 1).

Chromosomal Location. Unknown.

Comments. The ends of the sheep PCR product were sequenced to confirm that the product was *PRL*. The coding portion was 100% homologous to the corresponding region of the ovine *PRL* cDNA (GenBank accession no. M27057). Two Suffolk animals had a differing restriction pattern with *Hae*III, but this was not observed in any other animals. A polymorphism

was also identified with the restriction enzyme *Mse*I. This second polymorphism was observed to be in complete phase with the *Hae*III polymorphism in the families typed. Prolactin is an anterior pituitary hormone involved in many reproductive pathways.

Literature Cited

- Crawford, A. M., K. G. Dodds, A. J. Ede, C. A. Pierson, G. W. Montgomery, H. G. Garmonsway, A. E. Beattie, K. Davies, J. F. Maddox, S. W. Kappes, R. T. Stone, T. C. Nguyen, J. M. Penty, E. A. Lord, J. E. Broom, J. Buitkamp, W. Schwaiger, J. T. Epplen, P. Matthew, M. E. Matthews, D. J. Hulme, K. J. Beh, R. A. McGraw, and C. W. Beattie. 1995. An autosomal genetic linkage map of the sheep genome. *Genetics* 140:703.
- Truong, A. T., C. Duez, A. Belayew, A. Renard, R. Pictet, G. I. Bell, and J. A. Martial. 1984. Isolation and characterization of the human prolactin gene. *EMBO J.* 3(2):429.

Key Words: Sheep, PCR-RFLP, Prolactin Gene

J. Anim. Sci. 1997. 75:1686




Figure 1. *Hae*III RFLP in the ovine prolactin PCR fragments separated on a 3% NuSieve gel (FMC). Lane 1: 1-kb ladder (Promega); lane 2: AA genotype; lane 3: AB genotype; lane 4: BB genotype; lane 5: uncut PCR product.

¹The authors thank J. Medrano and N. Cockett for distribution of IMF DNA and D. Notter for additional sheep DNA. This work was supported in part by the Iowa Agriculture and Home Economics Experiment Station, Ames, Journal Paper No. 17192, Project No. 3231.

²To whom correspondence should be addressed.

Received January 6, 1997.

Accepted March 17, 1997.