

9-7-2018

# Environmentally induced phenotypic plasticity explains hatching synchrony in the freshwater turtle *Chrysemys picta*

Jessica K. McGlashan  
*University of Western Sydney*

Michael B. Thompson  
*The University of Sydney*

Fredric J. Janzen  
*Iowa State University, fjanzen@iastate.edu*

Ricky-John Spencer  
*University of Western Sydney*

Follow this and additional works at: [https://lib.dr.iastate.edu/eeob\\_ag\\_pubs](https://lib.dr.iastate.edu/eeob_ag_pubs)



Part of the [Population Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

The complete bibliographic information for this item can be found at [https://lib.dr.iastate.edu/eeob\\_ag\\_pubs/292](https://lib.dr.iastate.edu/eeob_ag_pubs/292). For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

---

# Environmentally induced phenotypic plasticity explains hatching synchrony in the freshwater turtle *Chrysemys picta*

## Abstract

Environmentally cued hatching allows embryos to alter the time of hatching in relation to environment through phenotypic plasticity. Spatially variable temperatures within shallow nests of many freshwater turtles cause asynchronous development of embryos within clutches, yet neonates still hatch synchronously either by hatching early or via metabolic compensation. Metabolic compensation and changes in circadian rhythms presumably enable embryos to adjust their developmental rates to catch up to more-advanced embryos within the nest. Hatchlings of the North American freshwater turtle *Chrysemys picta* usually overwinter within the nest and emerge the following spring, but still hatch synchronously via hatching early. Here we used rates of oxygen consumption and heart rate profiles to investigate the metabolic rates of clutches of *C. picta* developing in conditions that result in asynchronous development to determine if compensatory changes in metabolism occur during incubation. Embryos hatched synchronously and displayed circadian rhythms throughout incubation, but exhibited no evidence of metabolic compensation. Phenotypic traits of hatchlings, including body size and righting performance, were also not affected by asynchronous development. We conclude that less developed embryos of *C. picta* hatch synchronously with their clutchmates by hatching early, which does not appear to inflict a fitness cost to individuals. The ultimate mechanism for synchronous hatching in *C. picta* could be for hatchlings to ensure an optimal overwintering position within the centre of the nest. Consequently, immediate fitness costs will not hinder hatchling survival. The geographic location, as well as environmental and genetic factors unique to populations can all influence hatching behaviour in turtles through phenotypic plasticity. Hence, synchronous hatching is an adaptive bet-hedging strategy in turtles, but the mechanisms to achieve it are diverse.

## Keywords

circadian rhythms, convergent evolution, embryonic development, environmentally cued hatching, metabolic compensation, turtle

## Disciplines

Ecology and Evolutionary Biology | Population Biology | Terrestrial and Aquatic Ecology

## Comments

This is a manuscript of an article published as McGlashan, Jessica K., Michael B. Thompson, Fredric J. Janzen, and Ricky-John Spencer. "Environmentally induced phenotypic plasticity explains hatching synchrony in the freshwater turtle *Chrysemys picta*." *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology* (2018). doi: [10.1002/jez.2217](https://doi.org/10.1002/jez.2217). Posted with permission.

# **Title: Hatching plasticity in the freshwater turtle *Chrysemys picta* under variable incubation temperatures**

Jessica K. McGlashan<sup>1\*</sup>, Michael B. Thompson<sup>2</sup>, Fredric J. Janzen<sup>3</sup> and Ricky-John Spencer<sup>1</sup>

<sup>1</sup>School of Science and Health, University of Western Sydney, Locked Bag 1797, Penrith South DC, NSW 2751, Australia.

<sup>2</sup>School of Biological Sciences, Heydon-Laurence Building (A08), The University of Sydney NSW 2006, Australia

<sup>3</sup>Department of Ecology, Evolution and Organismal Biology, Iowa State University, 251 Bessey Hall, Ames, Iowa 50011 USA

\*Author for correspondence: Jessica McGlashan

Email: [j.mcglashan@westernsydney.edu.au](mailto:j.mcglashan@westernsydney.edu.au)

Phone: T +61 0433969804

**Running title:** Synchronous hatching in *Chrysemys picta*

**Key words:** circadian rhythms, convergent evolution, embryonic development, environmentally cued hatching, metabolic compensation, turtle.

## **Research Highlights**

Our study demonstrates that synchronous hatching in *Chrysemys picta* results from environmentally-induced phenotypic plasticity. We monitored heart rates and respiration rates during embryonic development to reveal that neonates can hatch 2-3 days earlier than expected without metabolic compensation or identifiable physiological consequence. Thus, synchronous hatching is likely a bet-hedging adaptive strategy in turtles and the mechanisms to achieve it are diverse, suggesting a likelihood of convergent evolution of hatching behaviour in turtles.

## **Abstract**

Environmentally cued hatching allows embryos to alter the time of hatching in relation to environment through phenotypic plasticity. Spatially variable temperatures within shallow nests of many freshwater turtles cause asynchronous development of embryos within clutches, yet neonates still hatch synchronously either by hatching early or via metabolic compensation. Metabolic compensation and changes in circadian rhythms presumably enable embryos to adjust their developmental rates to catch up to more-advanced embryos within the nest. Hatchlings of the North American freshwater turtle *Chrysemys picta* usually overwinter within the nest and emerge the following spring, but still hatch synchronously via hatching early. Here we used rates of oxygen consumption and heart rate profiles to investigate the metabolic rates of clutches of *C. picta* developing in conditions that result in asynchronous development to determine if compensatory changes in metabolism occur during incubation. Embryos hatched synchronously and displayed circadian rhythms throughout incubation, but exhibited no evidence of metabolic compensation. Phenotypic traits of hatchlings, including body size and righting performance, were also not affected by asynchronous development. We conclude that less developed embryos of *C. picta* hatch synchronously with their clutch-mates by hatching early, which does not appear to inflict a fitness cost to individuals. The ultimate mechanism for synchronous hatching in *C. picta* could be for hatchlings to ensure an optimal overwintering position within the centre of the nest. Consequently, immediate fitness costs will not hinder hatchling survival. The geographic location, as well as environmental and genetic factors unique to populations can all influence hatching behaviour in turtles through phenotypic plasticity. Hence, synchronous hatching is an adaptive bet-hedging strategy in turtles, but the mechanisms to achieve it are diverse.

## **Introduction**

Phenotypic plasticity is the interaction between environmental factors and genotype, which results in the expression of an environment-specific phenotype, and can impact fitness and survival (Scheiner, 1993; West-Eberhard, 1989, 2003). Variability in the environments experienced by embryos during development influences adaptive phenotypic plasticity (Scheiner, 1993; Scheiner & Lyman, 1991), altering the behaviour, morphology, and physiology of the embryo (Doody, 2011). Plasticity in developmental response to changes in the environment stimulates an expression of traits that reduce immediate threats to their survival, such as hatching behaviour, morphology, and performance (Doody, 2011; Shine, 2004).

Reptilian embryos can alter the timing of their hatching in response to environmental conditions through phenotypic plasticity within boundaries set by incubation temperature, which is known as environmentally-cued hatching (ECH) (Colbert, Spencer, & Janzen, 2010; McGlashan, Spencer, & Old, 2012; Spencer & Janzen, 2011; Spencer, Thompson, & Banks, 2001). There are three forms of ECH: early, synchronous, and delayed hatching (Doody, 2011; Spencer & Janzen, 2011). These mechanisms are not mutually exclusive and some species show multiple forms of ECH (Doody, 2011; Warkentin & Calswell, 2009). Timing of hatching can affect the survival of the offspring both immediately (risk of predation) and in the future (resource availability, size and performance of individual) (Brinkhof, Anton, Hage, & Simon, 1993; Colbert et al., 2010; O'Donoghue & Boutin, 1995; Sheldon, Kruuk, Merilä, & Crespi, 2003; Spencer & Janzen, 2011; Tucker, Paukstis, & Janzen, 2008). The ultimate mechanisms of ECH in reptiles are still poorly understood, however the proximate mechanisms yield variation in the developmental stage at hatching, and/or a reduction in the variation in incubation time, which may increase an individual's chance of survival through a

trade-off between the risks and benefits of hatching (Gomez-Mestre, Wiens, & Warkentin, 2008; Warkentin & Calswell, 2009).

Temperature significantly affects all biological processes from the molecular level, to organism-level processes, including metabolism and growth (Johnston, Vieira, & Hill, 1996; Packard, Packard, Miller, & Boardman, 1988; Rome, Stevens, & John-Alder, 1992; Wilhoft, 1958). Embryonic development and incubation period are sensitive to temperature changes, with warmer temperatures accelerating growth and developmental rate and therefore shortening incubation periods (Booth, 1998; Deeming & Ferguson, 1991; Monaghan, 2008). In the nests of many freshwater turtles, a vertical thermal gradient exists due to the diel fluctuations of temperature, and seasonal variation in temperatures (Telemeco et al., 2016; Thompson, 1988, 1997; Thompson, Packard, Packard, & Rose, 1996). Temperature thus affects eggs differently depending on their position in the nest (Georges, Limpus, & Stoutjesdijk, 1994), with eggs at the top of the nest generally experiencing warmer temperatures than eggs at the bottom, and these differences may be as high as 6 °C (Thompson, 1988, 1997). Such thermal variation within the nest has the potential to cause embryos to develop at different rates (asynchronous development) during the incubation period. Despite such variation in developmental rate between embryos, offspring of many species still hatch synchronously (Colbert et al., 2010; McGlashan et al., 2012; Spencer et al., 2001; Thompson, 1989).

Synchrony can occur when embryos delay hatching until stimulated by an environmental cue (Doody et al., 2001; Doody, Stewart, Camacho, & Christian, 2012; Webb, Choquenot, & Whitehead, 1986), or when less developed embryos, either adjust their developmental rate through metabolic compensation (McGlashan et al., 2012) or hatch at an earlier

developmental stage (Colbert et al., 2010; Spencer & Janzen, 2011). Previous experimental studies have shown that embryos that are initially less developed (i.e. incubating at cooler temperatures) advance their hatching date by 2-3 days to hatch at approximately the same time as their warmer incubated siblings (Colbert et al., 2010; McGlashan et al., 2012; Spencer et al., 2001). Under these conditions, embryos were seen to increase their developmental rates in the final third of incubation to “catch up” to the more advanced embryos, likely with a cost (increased heart and respiration rates, and reduced yolk sac volume), known as metabolic compensation (McGlashan et al., 2012). Precocial species exhibit a decline in metabolic rate several days before hatching (Booth, 2000; Booth & Astill, 2001; Du, Zhao, & Shine, 2010; McGlashan et al., 2012; Thompson, 1989). This period is considered a ‘resting stage’ where (1) tissue growth is essentially complete, (2) sensory, neuromuscular, and thermoregulatory systems mature (Vleck et al., 1979; Vleck, Vleck, & Hoyt, 1980; Webb et al., 1986; Thompson, 1989), and (3) any necessary acclimation can occur with minimal metabolic cost (McGlashan et al., 2012). Metabolic compensation is essential to survival as any reduction in developmental time can adversely affect growth and performance of hatchlings (Janzen, 1993; Vince & Chinn, 1971; Warkentin, 1995). Remarkably, metabolic compensation does not incur a reduction in hatchling size and locomotor performance (McGlashan, Loudon, Thompson, & Spencer, 2015; McGlashan et al., 2012).

Several factors have likely driven the evolution of synchronous hatching in turtles.

Synchronous hatching allows hatchlings to emerge from a nest together, thereby increasing their chance of survival by swamping predators, or reducing their exposure to prey switching generalist predators (Bradbury, Campana, Bentzen, & Snelgrove, 2004; Santos et al., 2016; Sih & Moore, 1993; Tucker et al., 2008; Warkentin, 1995; Warkentin, 2000). Embryos can hatch early and synchronously in response to a predatory attack (Vitt, 1991; Warkentin, 1995;

Warkentin, 2000), or when the risk of predation is reduced (Bradbury et al., 2004). Environmental cues may also stimulate synchronous hatching, as with pig-nosed turtles (*Carettochelys insculpta*), which can delay their hatching until an environmental trigger (hypoxia) stimulates hatching (Doody et al., 2001; Doody et al., 2012; Webb, Choquenot, & Whitehead, 1986). Painted turtles (*Chrysemys picta*) hatch synchronously apparently through early hatching, which could be an overwintering strategy to improve their positioning within the nest to increase their chance of survival until emerging the following spring (Colbert et al., 2010). In contrast, Murray River short-neck turtles (*Emydura macquarii*) and eastern long-necked turtles (*Chelodina longicollis*) both hatch synchronously as a predator avoidance strategy (Tucker et al., 2008) through metabolic compensation, which is evident through an increased rate of oxygen consumption ( $\dot{V}O_2$ ), and altered heart rate patterns during development (McGlashan et al., 2015; McGlashan et al., 2012). Circadian rhythms in heart rate also appear during development, and embryos might be able to detect the heart rate rhythm of their clutch-mates to communicate developmental stage, and thus respond to perceived differences by metabolically compensating (Aubret, Blanvillain, Bignon, & Kok, 2016; Loudon, Spencer, Strassmeyer, & Harland, 2013; McGlashan et al., 2015).

Offspring in many populations of freshwater turtles in the northern hemisphere overwinter in the natal nest, where hatchlings delay emergence until the following spring (Gibbons, 2013). *Chrysemys picta* frequently overwinter in the natal nest, yet they exhibit synchronous hatching (Colbert et al. 2010). It is not yet known whether *C. picta* also utilises metabolic compensation to hatch early. For *C. picta*, winter is a critical period for hatchling survival as temperatures within the nest can drop below -10 °C, and mortality due to freezing approaches 100% when temperatures fall to -14 °C (Packard, Fasano, Attaway, Lohmiller, & Lynch, 1997; Packard & Packard, 2004). Ensuring an optimum overwintering position in the nest,



where conditions are ideal for reducing energy expended or reducing the chance of mortality from freezing is a significant benefit of hatching early (Colbert et al., 2010; Spencer & Janzen, 2014). Still, early hatching *C. picta* has fitness costs, with a reduction in performance exhibited for up to 9 months after hatching (Colbert et al., 2010). This reduced performance suggests that embryos are missing an important stage of development, which may be vital for precocial species (Peterson & Kruegl, 2005; Vleck, Hoyt, & Vleck, 1979).

Synchronous hatching may be an adaptive “bet-hedging” strategy, with the mechanisms to achieve it differing among turtle species (Colbert et al., 2010; Doody et al., 2001; McGlashan et al., 2012). Metabolic compensation in Australian chelids enables embryos to complete development, which prevents them from being disadvantaged when they emerge from the nest (McGlashan et al., 2012). Metabolic compensation can be costly in terms of the yolk reserves necessary for post hatching growth and development (McGlashan et al., 2012; Radder, Shanbhag, & Saidapur, 2002; Radder, Shanbhag, & Saidapur, 2004). Because neonates of *C. picta* typically remain in the nest over winter, we would not expect to see metabolic compensation in this species, as utilising yolk reserves prior to hatching could be detrimental to their survival in the nest (Willette, Tucker, & Janzen, 2005). The aim of this study was to test the hypothesis that synchronous hatching occurs in *C. picta* by hatching early and not through metabolic compensation. We investigated metabolic rate by measuring  $\dot{V}O_2$  and heart rate profiles to detect variation among embryos in controlled conditions of synchronous and asynchronous incubation. Phenotypic and performance measures were also used to reflect developmental variation.

## **Materials and method**

### ***Egg collection and incubation***

Eggs of *C. picta* were collected from freshly built nests at the Thomson Causeway Recreation Area, Thomson, IL, USA from the 4<sup>th</sup>-7<sup>th</sup> June 2012. Eggs were excavated from nests within 5 h of oviposition and were placed in plastic containers with moist soil. Ten clutches (with an average of 11 eggs per clutch) were obtained over a two-day period and transported to Iowa State University. All eggs were uniquely marked using a HB pencil, weighed (g), and placed in plastic containers (200 x 100 x 50 mm), half buried in a 1:1 mixture of vermiculite and deionised water (-150 kPa) (Colbert et al., 2010; Spencer et al., 2001). To account for evaporation and maintain water potential of vermiculite, containers were weighed and rehydrated weekly. To counteract potential thermal gradients within incubators, the position of the containers were also rotated weekly (Janzen, 1993).

### ***Incubation method***

The treatment (asynchronous) group and the control (synchronous) group were set up using protocols similar to Spencer et al. (2001) and Colbert et al. (2010). Both the treatment and control groups contained five clutches, each with nine eggs, for a total of 45 eggs per treatment. To establish asynchrony within the five asynchronous clutches, three of the nine eggs from each clutch were incubated at 30 °C (asynchronous-more advanced), with the remaining six eggs (asynchronous-target) incubated at 26 °C for the first 7 d of incubation. After 7 d, the asynchronous-target eggs were reunited with the three asynchronous-more advanced eggs. The asynchronous-more advanced eggs were positioned between the asynchronous-target eggs (within 10 mm) (Fig. 1). All containers were then incubated at 28 °C until hatching. The remaining five clutches in the control (synchronous) group were arranged using the same procedure to control for egg movement in the experimental setup. Three eggs (synchronous-not advanced) were incubated at 26 °C for the 7 d period. The remaining six eggs (synchronous-target) were also incubated at 26 °C so that all nine eggs

were still at the same developmental stage when reunited (Spencer et al., 2001). After eggs were reunited, containers were incubated at 28 °C until hatching. Each clutch was incubated in a separate container throughout the experiment. Heart rates and  $\dot{V}O_2$  of target embryos from asynchronous and synchronous groups were monitored weekly and containers were opened every 2-3 d to allow fresh air exchange throughout incubation until the commencement of pipping/hatching.

### ***Oxygen consumption***

The rate of oxygen consumption ( $\dot{V}O_2$ ) was measured from week 4 to 8 of the eight-week incubation period. One target egg from each clutch (5 asynchronous and 5 synchronous), was selected for  $O_2$  analysis each week.  $\dot{V}O_2$  of each egg was measured twice within a 24 h period and the average of both measures was used. Oxygen (%) was measured in a closed system using an S-3A oxygen analyser (AEI Technologies, USA). The analyser was first calibrated (20.94 %  $O_2$ ) using 50 cc of atmospheric air injected using a syringe and syringe pump (Vleck, 1987). Airtight chambers were constructed from 500 mL metal paint cans and sealed with lids that were fitted with a 2-way stopcock. Eggs were placed on moist cotton balls (to keep them from desiccating) in the opened chamber and allowed to come to thermal equilibrium in the 28 °C incubator approximately 5 min before sealing. The lid was sealed and an initial ( $F_I$ ) air sample was taken from the outflow port with a syringe fitted with a stopcock. The chamber pressure was allowed to equilibrate before the outflow port was sealed and containers were carefully placed in the 28 °C incubator. A final ( $F_E$ ) sample was taken at the conclusion before the egg was returned to its original incubator container. To measure  $\dot{V}O_2$ , eggs remained in the air-tight chambers for 1 to 5 h, depending on the stage of incubation. A blank control was in place to account for any leakage or spontaneous change in  $O_2$  not caused by the egg. Temperature and air pressure were recorded at the beginning of the

experiment to allow correction to standard temperature pressure volume of each can.  $F_I$  and  $F_E$  samples were then put through the  $O_2$  analyser via two 3 mL syringe barrels, one with silica gel and the other with a combination of ascarite and soda lime to remove  $CO_2$  and moisture from the sample. Fractional  $O_2$  concentrations of  $F_I$  and  $F_E$  samples were analysed, and  $\dot{V}O_2$  was calculated using the equation  $\dot{V}O_2 = V \frac{(F_I - F_E)}{t(1 - F_E)}$  where  $V$  represents the dry gas volume in the chamber and  $t$  represents time between  $F_I$  and  $F_E$  samples (Vleck et al., 1979; Vleck et al., 1980; Vleck, 1987)

### ***Heart rate measurements***

Heart rates (bpm) were measured from weeks 3 to 8 of incubation to determine 1) heart rate profile, 2) circadian rhythm, and 3) metabolic compensation. Heart rates were used to infer metabolic rate (Du, Radder, Sun, & Shine, 2009; Wallace & Jones, 2008) of embryos and identify any difference between asynchronous and synchronous groups. One target egg from each clutch was removed from its incubator container and immediately placed in a Buddy digital egg monitor system (Avian Biotech, England) in complete darkness. Three readings were taken once heart rates had stabilised (within 2 min) (McGlashan et al., 2015; Sartori, Leite, Abe, Crossley Ii, & Taylor, 2015). Heart rates were calculated as the mean of the three readings, that were taken at 30 s intervals. Eggs were not disturbed during the recording period, thus any embryonic movement could be detected and distinguished from heart rate readings, in which case we waited for movement to cease and heart rate to stabilise before recording again (McGlashan et al., 2015). The heart rate of each egg was taken at four time intervals over a 24 h period (0500 h, 1100 h, 1700 h and 2300 h) once a week to identify circadian rhythms and the average daily heart rate for the weekly profile. Each week a different egg was used to record heart rate. Care was taken not to exceed this time due to a potential heating effect from the Buddy system (Sartori et al., 2015).

### *Pipping and post hatching measurements*

To determine incubation period, eggs were inspected for signs of pipping (initial perforation of the eggshell with caruncle) twice daily (Gutzke, Paukstis, & Packard, 1984). The incubation period was measured as time in days from oviposition until pipping. Neonates were allowed to hatch unassisted, after which they were weighed (g), straight carapace length and width, straight plastron length and width were measured (mm), and righting ability was recorded within 12 h of hatching. To measure righting ability, all hatchlings were placed on their back (carapace), and the time taken to right themselves was measured, as were latency to begin moving and total active righting time. All tests were terminated after 180 s. Measurements of righting ability were used as an index of neuromuscular development (Colbert et al., 2010; Peterson & Kruegl, 2005; Vleck et al., 1979; Vleck et al., 1980).

### *Data analysis*

Clutch was included as a random effect in all models and where treatment effects were significant, post-hoc Tukey tests were used to determine where pairwise differences occurred. Incubation period was analysed in all four independent variables (asynchronous-more advanced, asynchronous-target, synchronous-more advanced synchronous-target) using a linear mixed model (PROC MIXED; SAS, version 9.3, SAS Institute, Cary, NC) and egg mass was used as a covariate. Embryonic  $\dot{V}O_2$  and heart rates were also compared among weeks and among asynchronous and synchronous groups using PROC MIXED. Egg mass and week of incubation were included in the model as covariates.

Circadian rhythms were determined through daily heart rate fluctuations, which were calculated as a difference from the mean diel rate at each time point. Peaks were aligned with

time of day in both asynchronous and synchronous groups to determine circadian rhythms. Differences from the mean diel rate were analysed between asynchronous and synchronous groups each week and at each time point as a repeated measure using PROC MIXED with egg mass as a covariate. The differences between heart rates at each time point (i.e. -6 h to peak) within treatment groups were analysed using a one-way ANOVA.

Phenotypic variation in carapace length and width, plastron length and width, and hatchling mass were compared among asynchronous and synchronous groups using a multivariate analysis of variance (MANCOVA) with egg mass as a covariate (PROC GLM). Hatchling righting ability was analysed using PROC MIXED with hatchling mass as a covariate. A binomial test was used to assess propensity to right of hatchlings (PROC GLIMMIX).

All  $\dot{V}O_2$ , heart rate, and growth data analysed with PROC MIXED were log-transformed before analysis to ensure linearity in the models. Examination of residuals ensured that all assumptions of univariate parametric statistics were met. In the multivariate analysis, the assumption of multivariate normality could not be directly tested due to sample size limitations. Thus, Pillai's Trace was used as a test statistic because it is robust to violations of multivariate normality (Scheiner, 2001). Statistical significance was determined at the 0.05 Type I error level, and data are presented as the mean  $\pm$  SE.

## **Results**

### ***Incubation period***

Of the 89 eggs, 11 failed to hatch (2 asynchronous-target eggs, 4 synchronous-target eggs, 3 asynchronous-more advanced eggs, and 2 synchronous-not advanced eggs). Hatching synchrony occurred within clutches in both asynchronous (treatment) ( $t_{73} = 1.84$ ,  $p = 0.2641$ )

and synchronous groups (control) ( $t_{73} = 0.04$ ,  $p = 1$ ), with incubation period averaging  $54 \pm 0.3$  and  $55 \pm 0.3$  d, respectively (Fig. 2). The incubation period of eggs in asynchronous or synchronous clutches was significantly affected by treatment ( $F_{3,73} = 4.30$ ,  $p = 0.0076$ ). Pairwise comparisons showed that asynchrony was established, as the incubation period of asynchronous-more advanced and synchronous-not advanced eggs differed by 2 d ( $t_{73} = 2.84$ ,  $p = 0.0293$ ). Asynchronous-more advanced eggs also hatched earlier than synchronous-target eggs and synchronous-not advanced eggs within clutches ( $t_{73} = 3.25$ ,  $p = 0.0091$ ), but incubation period did not differ between asynchronous-target eggs and synchronous-target eggs ( $t_{73} = 1.90$ ,  $p = 0.2361$ ).

### ***Oxygen consumption***

Treatment and control groups did not differ in  $\dot{V}O_2$  at any time point during incubation ( $F_{1,30} = 0.01$ ,  $p = 0.9149$ ; Fig. 3), but there was a time effect ( $F_{4,30} = 3.33$ ,  $p = 0.0226$ ).  $\dot{V}O_2$  increased with egg mass ( $F_{1,30} = 15.78$ ,  $p = 0.0004$ ), and there was an interaction effect between egg mass and time ( $F_{4,30} = 0.27$ ,  $p = 0.0432$ ) but not between egg mass and treatment ( $F_{1,30} = 0.12$ ,  $p = 0.7289$ ), treatment and time ( $F_{4,30} = 0.27$ ,  $p = 0.8969$ ), or egg mass, treatment, and time ( $F_{4,30} = 0.28$ ,  $p = 0.8860$ ).  $\dot{V}O_2$  of asynchronous-target eggs peaked at 0.48 mL/h in week 6 of incubation and declined to 0.16 mL/h prior to hatching in week 8. Synchronous-target eggs peaked at 0.45 mL/h in week 5 then gradually declined to 0.17 mL/h in week 8.  $\dot{V}O_2$  increased from week 4-5 within each group ( $t_{30} = -5.15$ ,  $p < 0.0001$ ) and declined from week 7 to 8 of incubation ( $t_{30} = 6.77$ ,  $p < 0.0001$ ).

### ***Heart rate***

Heart rates did not differ significantly between asynchronous-target eggs and synchronous-target eggs at any time point during incubation ( $F_{1,36} = 1.19$ ,  $p = 0.2819$ ; Fig. 4). They were

further unaffected by egg mass and by interactions of treatment and egg mass. Heart rates did vary among weeks of incubation ( $F_{5,36} = 5.55$ ,  $p = 0.0007$ ), gradually decreasing from 96 ( $\pm 2.6$ ) to 60 ( $\pm 2.6$ ) bpm in asynchronous groups and from 100 to 67 bpm in synchronous groups from week 3 to 8 of incubation. There was no interaction effect of egg mass and time, treatment and time, nor egg mass, treatment, and time. Heart rate in both treatment groups declined from week 6 to 7 ( $t_{47} = 3.71$ ,  $p = 0.0069$ ) and week 7 to 8 ( $t_{47} = 3.60$ ,  $p = 0.0094$ ) of incubation.

Circadian rhythms in heart rates were present, with a clear peak in heart rates throughout the day in both asynchronous-target eggs (-6 h to peak-  $F_{1,14} = 18.92$ ,  $p = 0.0007$ ; peak to 6 h-  $F_{1,14} = 20.41$ ,  $p = 0.0005$ ; Fig. 5) and synchronous-target eggs (-6 h to peak-  $F_{1,14} = 5.59$ ,  $p = 0.03$ ; peak to 6 h-  $F_{1,14} = 24.69$ ,  $p = 0.0002$ ). Individuals showed minimum and maximum heart rates over a 24 h period that varied up to 6 bpm, but they were not correlated with the time of day in both asynchronous and synchronous groups. There was no significant difference between asynchronous and synchronous groups at each time point; -6 h ( $F_{1,30} = 0.22$ ,  $p = 0.64$ ), peak ( $F_{1,30} = 0.00$ ,  $p = 0.99$ ), 6 h ( $F_{1,30} = 1.39$ ,  $p = 0.25$ ), nor 12 h ( $F_{1,30} = 0.01$ ,  $p = 0.94$ ). There was a modest week effect ( $F_{4,30} = 2.80$ ,  $p = 0.044$ ) and egg mass by week interaction effect ( $F_{4,30} = 3.38$ ,  $p = 0.021$ ) during the peak time point, but further pairwise tests revealed no significant differences between weeks. No other main effects or interaction effects were statistically significant at any time point.

### ***Post hatching measurements***

Multivariate analysis of all measured phenotypes (carapace length and width, plastron length and width, and mass) post hatching revealed that hatchling size was not affected by the interaction between treatment and egg mass (Pillai = 0.2142,  $F_{15,174} = 0.89$   $p = 0.5742$ ).



Treatment alone also did not affect hatchling size (Pillai = 0.165,  $F_{10,114} = 1.03$   $p = 0.4265$ ), with egg mass alone explaining the majority of variation in body size (Pillai = 0.647,  $F_{5,56} = 20.54$   $p < 0.0001$ ). The average measurements for carapace length in asynchronous groups was  $25.60 \pm 0.4$  mm and synchronous group was  $25.57 \pm 0.4$  mm. Plastron length in asynchronous groups was  $24.72 \pm 0.32$  mm and synchronous groups was  $24.74 \pm 0.38$  mm, and hatchling mass was  $4.96 \pm 0.31$  g and  $4.99 \pm 0.20$  g respectively.

Performance and neuromuscular development, as assessed by righting ability, were not affected by treatment group. There was no significant difference between any of the treatment groups in ability to right successfully ( $F_{2,65} = 1.3$ ,  $p = 0.2783$ ). Latency to right, total time to right, and active righting time also were not affected by treatment group.

## **Discussion**

Here we show *Chrysemys picta* neonates hatch synchronously regardless of developmental stage, with less developed embryos hatching at the same time as their more advanced clutch-mates, with no indication of metabolic compensation. Asynchronous-target eggs hatched within 1 day of the asynchronous-more advanced eggs with which they were incubated with, and this was consistent with the synchronous group, where synchrony was expected. The incubation period of asynchronous-more advanced eggs was shortened compared to the synchronous-not advanced eggs, indicating that the asynchronous setup (Fig. 1) resulted in a substantial difference in incubation period, like that detected in prior work (Colbert et al., 2010).

There was no evidence of metabolic compensation inferred through heart rate and  $\dot{V}O_2$  of developing target eggs in both the asynchronous and synchronous clutches, as these measures

were similar between treatments at all time points.  $\dot{V}O_2$  peaked during weeks 5 and 6 of incubation and declined in the final 2 weeks, as was expected based on previous studies (Booth & Astill, 2001; Booth, Thompson, & Herring, 2000; Du, Thompson, & Shine, 2010; McGlashan et al., 2012; Thompson, 1989). Embryonic heart rates were also similar between target eggs in both asynchronous and synchronous clutches, and these gradually decreased from weeks 3 to 8 of incubation, similar to the pattern seen in some semi-precocial birds (Cain, Abbott, & Rogallo, 1967; Tazawa, 2005; Tazawa, Hiraguchi, Kuroda, Tullett, & Deeming, 1991; Tazawa, Kuroda, & Whittow, 1991) and snapping turtles (*Chelydra serpentina*) (Birchard & Reiber, 1996). The decline in metabolic rate during the resting period (80-90 % of development) prior to hatching indicated by both heart rate and  $\dot{V}O_2$  has also been detected in embryos of *Ch. longicollis* (McGlashan et al., 2015), *Ca. insculpta* (Webb et al., 1986), *E. macquarii* (McGlashan et al., 2012; Thompson, 1989), and the American alligator (*Alligator mississippiensis*) (Thompson, 1989). The resting period is when acclimation occurs, where metabolic costs for growth are not usually as high, and the effects of temperature are relatively less, in comparison to early development (Birchard & Reiber, 1996; Birchard & Reiber, 1995). Hence, if metabolic compensation was to occurring, it would be expected during this period, as seen in *E. macquarii* (McGlashan et al., 2012). Synchronous hatching with no evidence of metabolic compensation suggests that less advanced embryos are hatching early with the more-advanced eggs in the clutch, and possibly missing out on an important stage of development.

Circadian rhythms were present in developing *C. picta* embryos despite an absence of environmental cues, such as photoperiod. Over a 24 h period, heart rates increased and reached a 'peak', indicating high developmental activity, and then dropped, suggesting low developmental activity or daily 'resting'. These rhythms are not synchronised to the time of

day among eggs and there is no indication of synchrony within clutches when compared from week to week, as seen in *Ch. longicollis* and *E. macquarii* (Loudon et al., 2013; McGlashan et al., 2012). The mean weekly heart rate reflects metabolic rate of embryos, and deviations from the mean rate throughout the day suggest positive or negative compensatory changes in embryonic metabolism (McGlashan et al., 2015). Asynchronous groups did not exceed the potential for periods of higher peak daily rate or reduced resting periods relative to synchronous groups. There was no significant difference in any deviation from the mean between asynchronous and synchronous groups at each time point, implying an absence of metabolic compensation via diel heart rate fluctuations.

Hatchlings were not obviously disadvantaged by hatching early, in either growth or development. The lack of a differences in body size or mass of hatchlings between treatments, and their ability to right themselves, indicates that hatching early does not substantially affect development. Embryos that were developmentally behind during incubation hatched synchronously with more advanced embryos. There was no evidence to indicate metabolic compensation, nor any phenotypic or performance differences at the time of hatching, suggesting that there is potential for small shifts in hatching timing without observable metabolic or phenotypic costs. Although previous studies have found that synchronous hatching affects the performance of neonates, a difference in the length of time used to experimentally establish asynchrony might be the cause. Colbert et al. (2010) kept more advanced embryos at a warmer temperature for 11 d. Siblings were most likely too under-developed and, when stimulated to hatch synchronously by these more advanced clutch-mates, were ultimately disadvantaged in performance. Comparisons of the two studies on *C. picta* indicate that temporal differences in temperature extremes that result in a greater variation in developmental rates may also be biologically relevant.

In terms of biological significance, climatic extremes could affect incubation conditions (Walther et al., 2002) and therefore the level of asynchrony between clutch-mates. Eggs at the top of the nest experience greater temperature fluctuations throughout the day and are warmer for more than 75% of the day when compared to eggs at the bottom (Shine & Elphick, 2001; Telemeco et al., 2016; Thompson, 1988, 1997; Thompson et al., 1996). Air temperature, substrate thermal conductance, and chamber dimensions can also influence nest temperatures (Thompson, 1988). Each factor determines the extent to which temperature differences within a nest might yield variation in embryonic metabolism and growth. Natural conditions in nests could stimulate greater differences among clutch-mates than elicited experimentally, and thus result in neonates with reduced performance. Further studies are needed to determine the upper limit of synchronous hatching in natural conditions. The final 10% of incubation is when hatching synchrony can take place once hatching competence is achieved (Doody, 2011; Gomez-Mestre et al., 2008;). The asynchronous set up in this study was 12% of the total incubation period, but in Colbert et al. (2010) the set up was 20% of the total incubation period. Nevertheless, both this study and Colbert et al. (2010) demonstrated that synchronous hatching occurs in *C. picta* regardless of developmental differences at the beginning of incubation and the potential costs that may arise from hatching early. If the difference in development is small, then synchrony will not incur performance costs, but if the difference is great, then hatchling recruitment in some years may decline because more individuals are underdeveloped at hatching.

Although temperature is the major determinant for embryonic development in ectotherms (Booth, 1998; Deeming & Ferguson, 1991; Monaghan, 2008), hatching time is nonetheless plastic (Doody, 2011; Packard & Packard, 2000; Spencer & Janzen, 2011). Abiotic and biotic

forces unique to species and populations drive synchronous hatching, with environmental cues (e.g. heart rate, movement, embryo-embryo communication) within the nest potentially triggering metabolic compensation or early hatching (Doody, 2011). Of particular interest are the physiological mechanisms and endocrine responses, such as how hormones might alter development and behaviour. Thyroid hormones can induce changes to metabolism or hatching behaviour independent of temperature without any metabolic or post hatching cost (McGlashan, Thompson, Van Dyke, & Spencer, 2017; O'Steen & Janzen, 1999). Increased thyroid hormone concentrations strongly correlate with the final stages of embryogenesis (Dimond, 1954; Shepherdley et al., 2002), and could be important for hatching behaviour (McGlashan et al., 2017).

Embryonic development is complex and the environment in which embryos develop can influence hatching behaviour. By overwintering in the nest, hatchlings presumably experience a more favourable environment and reduced risk of predation when they emerge in the following spring (Gibbons, 2013; Gibbons & Nelson, 1978; Wilbur, 1975).

Synchrony by hatching early could ensure an optimal overwintering position in the nest for *C. picta* to reduce the chance of mortality from freezing, dehydration, and starvation from loss of energy reserves (Costanzo, Dinkelacker, Iverson, & Lee, 2004; Costanzo, Lee, & Ultsch, 2008; Spencer & Janzen, 2014). Winter nest temperatures are variable (Weisrock & Janzen, 1999) and could influence the overall energy expenditures of neonates in the nest. During winter, metabolic demands are generally low but maternally sourced yolk is utilised during this time, and the warmer the winter, the more energy is used (Muir, Dishong, Lee, & Costanzo, 2013; Willette et al., 2005). In *C. picta*, where warmer incubation conditions produce more females and cooler temperatures produce more males, males in these cooler overwintering nests utilise less yolk than females in these same nests, and the converse is true

for females from warmer nests (Spencer & Janzen, 2014). Thus, retaining yolk energy stores without metabolic compensation is imperative in preparation for atypical overwintering conditions, which can result in up to a 50% loss of energy (Spencer & Janzen, 2014).

The timing of hatching is a life-history trait influenced by the trade-off between the risk and benefit of hatching, including growth and development, hatchling energetics, and predator avoidance (Baker et al., 2013; Doody et al., 2001; Doody et al., 2012; Santos et al., 2016; Webb et al., 1986). Whether emergence is triggered immediately after hatching or several months later could depend on differing selective pressures, including geographic location, environmental and/or genetic factors unique to populations (Breitenbach, Congdon, & Sels, 1984). The phenomenon of synchronous hatching appears uniform in turtles, but the mechanisms to overcome the effect of variable temperatures on developmental rates vary considerably. Pleurodiran turtles metabolically compensate during incubation to hatch early and ensure synchronous hatching (McGlashan et al., 2012; McGlashan et al., 2015). Within cryptodires, at least one species delays hatching to synchronously emerge when environmental conditions are right (Doody et al., 2001; Doody et al., 2012; Webb, Choquenot, & Whitehead, 1986), whereas *C. picta* differ altogether by hatching early with no evidence of metabolic compensation during embryonic development and no evidence that hatchlings had a reduction in size or performance. Developmental plasticity allows embryos to hatch synchronously despite the variation in temperature, with various abiotic and biotic driving forces for individual species. In sum, synchronous hatching is an adaptive bet-hedging strategy in turtles characterised by diverse mechanisms to achieve it.

## **Acknowledgements**

Research was supported by a UWS SSH Equipment Fund Grant and NSF grant DEB-0640932, and was conducted under permits from the US Fish and Wildlife Service, Illinois Department of Natural Resources, and Iowa State University Institutional Animal Care and Use Committee. We thank the US Army Corps of Engineers for access to the Thomson Causeway, and T. S. Mitchell, J. Maciel, B. Bodensteiner and C. Hinsley for lab and field assistance. We also thank J. U. Van Dyke for statistical advice and helpful comments, B. P. Bentley for support and editorial help and D. Vleck for use of his laboratory and respirometry equipment.

## **Author contributions**

This manuscript is jointly authored, J.K.M., conceived the idea and, along with R-J.S., F.J.J. and M.B.T., designed the methodology. J.K.M. collected the data and carried out data analysis. All four authors contributed critically to the writing of the manuscript and gave final approval for the publication.

## **Competing interests**

The authors declare no competing or financial interests.

## **Funding**

Research was supported by a UWS SSH Equipment Fund Grant and NSF grant DEB-0640932.

## **Data accessibility**

Data will be made available from the Dryad Digital Repository upon acceptance of the manuscript.

## References

- Aubret, F., Blanvillain, G., Bignon, F., & Kok, P. J. R. (2016). Heartbeat, embryo communication and hatching synchrony in snake eggs. *Scientific Reports*, *6*, 23519. doi:10.1038/srep23519
- Baker, P. J., Costanzo, J. P., Iverson, J. B., & Lee, R. E. (2013). Seasonality and interspecific and intraspecific asynchrony in emergence from the nest by hatchling freshwater turtles. *Canadian Journal of Zoology*, *91*(7), 451-461. doi:10.1139/cjz-2012-0335
- Birchard, G., & Reiber, C. (1996). Heart rate during development in the turtle embryo: effect of temperature. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology*, *166*(8), 461-466. doi:10.1007/bf02338288
- Birchard, G. F., & Reiber, C. L. (1995). Growth, metabolism, and chorioallantoic vascular density of developing snapping turtles (*Chelydra serpentina*): influence of temperature. *Physiological Zoology*, *68*(5), 799-811. doi:10.2307/30163932
- Booth, D. T. (1998). Incubation of turtle eggs at different temperatures: do embryos compensate for temperature during development? *Physiological Zoology*, *71*(1), 23-26.
- Booth, D. T. (2000). Incubation of eggs of the Australian broad-shelled turtle, *Chelodina expansa* (Testudinata : Chelidae), at different temperatures: effects on pattern of oxygen consumption and hatchling morphology. *Australian Journal of Zoology*, *48*(4), 369-378.
- Booth, D. T., & Astill, K. (2001). Incubation temperature, energy expenditure and hatchling size in the green turtle (*Chelonia mydas*), a species with temperature-sensitive sex determination. *Australian Journal of Zoology*, *49*(4), 389-396. doi:http://dx.doi.org/10.1071/ZO01006
- Booth, D. T., Thompson, M. B., & Herring, S. (2000). How incubation temperature influences the physiology and growth of embryonic lizards. *Journal of Comparative Physiology B*, *170*(4), 269-276. doi:10.1007/s003600000097
- Bradbury, I. R., Campana, S. E., Bentzen, P., & Snelgrove, P. V. R. (2004). Synchronized hatch and its ecological significance in rainbow smelt *Osmerus mordax* in St. Mary's Bay, Newfoundland. *Limnology and Oceanography*, *49*(6), 2310-2315.



- Breitenbach, G. L., Congdon, J. D., & Sels, R. C. v. L. (1984). Winter temperatures of *Chrysemys picta* nests in Michigan: effects on hatchling survival. *Herpetologica*, 40(1), 76-81. doi:10.2307/3892463
- Brinkhof, M. W. G., Anton, J. C., Hage, F. J., & Simon, V. (1993). Timing of Reproduction and Fledging Success in the Coot *Fulica atra*: Evidence for a Causal Relationship. *Journal of Animal Ecology*, 62(3), 577-587.
- Cain, J. R., Abbott, U. K., & Rogallo, V. L. (1967). Heart rate of the developing chick embryo. *Experimental Biology and Medicine*, 126(2), 507-510. doi:10.3181/00379727-126-32490
- Carr, A., & Hirth, H. (1961). Social facilitation in green turtle siblings. *Animal Behaviour*, 9(1-2), 68-70. doi:10.1016/0003-3472(61)90051-3
- Colbert, P. L., Spencer, R.-J., & Janzen, F. J. (2010). Mechanism and cost of synchronous hatching. *Functional Ecology*, 24(1), 112-121. doi:10.1111/j.1365-2435.2009.01602.x
- Costanzo, J. P., Dinkelacker, Stephen A., Iverson, John B., & Lee, Richard E. J. (2004). Physiological ecology of overwintering in the hatchling painted turtle: multiple-scale variation in response to environmental stress. *Physiological and Biochemical Zoology*, 77(1), 74-99. doi:10.1086/378141
- Costanzo, J. P., Lee, R. E., & Ultsch, G. R. (2008). Physiological ecology of overwintering in hatchling turtles. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 309A(6), 297-379. doi:doi:10.1002/jez.460
- Deeming, D. C., & Ferguson, M. W. J. (1991). *Egg incubation: its effects on embryonic development in birds and reptiles*. Cambridge: Cambridge University Press.
- Dimond, S. M. T. (1954). The reactions of developing snapping turtles, *Chelydra serpentina serpentina* (Linné), to thiourea. *Journal of Experimental Zoology*, 127(1), 93-115. doi:10.1002/jez.1401270106
- Doody, J. S. (2011). Environmentally cued hatching in reptiles. *Integrative and Comparative Biology*, 51(1), 49-61. doi:10.1093/icb/icr043
- Doody, J. S., Georges, A., Young, J. E., Pauza, M. D., Pepper, A. L., Alderman, R. L., & Welsh, M. A. (2001). Embryonic aestivation and emergence behaviour in the pig-nosed turtle, *Carettochelys insculpta*. *Canadian Journal of Zoology*, 79(6), 1062-1072. doi:doi:10.1139/z01-061
- Doody, J. S., Stewart, B., Camacho, C., & Christian, K. (2012). Good vibrations? Sibling embryos expedite hatching in a turtle. *Animal Behaviour*, 83(3), 645-651. doi:http://dx.doi.org/10.1016/j.anbehav.2011.12.006

- Du, W.-G., Radder, R. S., Sun, B., & Shine, R. (2009). Determinants of incubation period: do reptilian embryos hatch after a fixed total number of heart beats? *Journal of Experimental Biology*, 212(9), 1302-1306. doi:10.1242/jeb.027425
- Du, W.-G., Thompson, M. B., & Shine, R. (2010). Facultative cardiac responses to regional hypoxia in lizard embryos. *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology*, 156(4), 491-494. doi:10.1016/j.cbpa.2010.04.005
- Du, W.-G., Zhao, B., & Shine, R. (2010). Embryos in the fast lane: high-temperature heart rates of turtles decline after hatching. *PLoS ONE*, 5(3), e9557. doi:10.1371/journal.pone.0009557
- Ferguson, M. W. J. (1985). Reproductive biology and embryology of the crocodylians. In *Biology of the Reptilia* (Vol. 14, pp. 329-491). New York: John Wiley & Sons.
- Georges, A., Limpus, C., & Stoutjesdijk, R. (1994). Hatchling sex in the marine turtle *Caretta caretta* is determined by proportion of development at a temperature, not daily duration of exposure. *Journal of Experimental Zoology*, 270(5), 432-444. doi:doi:10.1002/jez.1402700504
- Gibbons, J. W. (2013). A long-term perspective of delayed emergence (aka overwintering) in hatchling turtles: some they do and some they don't, and some you just can't tell. *Journal of Herpetology*, 47(2), 203-214. doi:10.1670/12-122
- Gibbons, J. W., & Nelson, D. H. (1978). The evolutionary significance of delayed emergence from the nest by hatchling turtles. *Evolution*, 32(2), 297-303.
- Gomez-Mestre, I., Wiens, J. J., & Warkentin, K. M. (2008). Evolution of adaptive plasticity: risk-sensitive hatching in neotropical leaf-breeding treefrogs. *Ecological Monographs*, 78(2), 205-224. doi:10.1890/07-0529.1
- Gutzke, W. H., Paukstis, G. L., & Packard, G. C. (1984). Pipping versus hatching as indices of time of incubation in reptiles. *Journal of herpetology*, 494-496.
- Ims, R. A. (1990). The ecology and evolution of reproductive synchrony. *Trends in Ecology and Evolution*, 5(5), 135-140. doi:10.1016/0169-5347(90)90218-3
- Janzen, F. J. (1993). An experimental analysis of natural selection on body size of hatchling turtles. *Ecology*, 74(2), 332-341. doi:10.2307/1939296
- Johnston, I. A., Vieira, V. L. A., & Hill, J. (1996). Temperature and ontogeny in ectotherms: muscle phenotype in fish. In C. U. Press (Ed.), *Animals and Temperature: Phenotypic and Evolutionary Adaptation* (I.A. Johnston, A.F. Bennett ed., Vol. 59). Cambridge: Cambridge University Press.

- Loudon, F. K., Spencer, R.-J., Strassmeyer, A., & Harland, K. (2013). Metabolic circadian rhythms in embryonic turtles. *Integrative and Comparative Biology*, *53*(1), 175-182. doi:10.1093/icb/ict040
- McGlashan, J. K., Loudon, F. K., Thompson, M. B., & Spencer, R.-J. (2015). Hatching behavior of eastern long-necked turtles (*Chelodina longicollis*): the influence of asynchronous environments on embryonic heart rate and phenotype. *Comparative Biochemistry and Physiology, Part A: Molecular & Integrative Physiology*, *188*, 58-64. doi:http://dx.doi.org/10.1016/j.cbpa.2015.06.018
- McGlashan, J. K., Spencer, R.-J., & Old, J. M. (2012). Embryonic communication in the nest: metabolic responses of reptilian embryos to developmental rates of siblings. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1734), 1709.
- McGlashan, J. K., Thompson, M. B., Van Dyke, J. U., & Spencer, R.-J. (2017). Thyroid hormones reduce incubation period without developmental or metabolic costs in Murray River short-necked turtles (*Emydura macquarii*). *Physiological and Biochemical Zoology*, *90*(1), 34-46. doi:10.1086/689744
- Monaghan, P. (2008). Early growth conditions, phenotypic development and environmental change. *363*(1497), 1635-1645. doi:10.1098/rstb.2007.0011
- Muir, T. J., Dishong, B. D., Lee, R. E., & Costanzo, J. P. (2013). Energy use and management of energy reserves in hatchling turtles (*Chrysemys picta*) exposed to variable winter conditions. *Journal of Thermal Biology*, *38*(6), 324-330. doi:http://dx.doi.org/10.1016/j.jtherbio.2013.04.003
- O'Donoghue, M., & Boutin, S. (1995). Does Reproductive Synchrony Affect Juvenile Survival Rates of Northern Mammals? *Oikos*, *74*(1), 115-121.
- O'Steen, S., & Janzen, Fredric J. (1999). Embryonic Temperature Affects Metabolic Compensation and Thyroid Hormones in Hatchling Snapping Turtles. *Physiological and Biochemical Zoology*, *72*(5), 520-533.
- Packard, G., Packard, M., Miller, K., & Boardman, T. (1988). Effects of temperature and moisture during incubation on carcass composition of hatchling snapping turtles (*Chelydra serpentina*). *Journal of Comparative Physiology B*, *158*(1), 117-125. doi:10.1007/BF00692735
- Packard, G. C., Fasano, S. L., Attaway, M. B., Lohmiller, L. D., & Lynch, T. L. (1997). Thermal environment for overwintering hatchlings of the painted turtle (*Chrysemys picta*). *Canadian Journal of Zoology*, *75*(3), 401-406. doi:10.1139/z97-050

- Packard, G. C., & Packard, M. J. (2000). Developmental plasticity in painted turtles, *Chrysemys picta*. *Functional Ecology*, 14(4), 474-483. doi:10.1046/j.1365-2435.2000.00448.x
- Packard, G. C., & Packard, M. J. (2004). To freeze or not to freeze: adaptations for overwintering by hatchlings of the North American painted turtle. *Journal of Experimental Biology*, 207(17), 2897-2906. doi:10.1242/jeb.01123
- Peterson, C. C., & Kruegl, A. (2005). Peaked temporal pattern of embryonic metabolism in an emydid turtle (*Chrysemys picta picta*). *Journal of Herpetology*, 39(4), 678-681. doi:10.1670/122-05N.1
- Radder, R. S., Shanbhag, B. A., & Saidapur, S. K. (2002). Pattern of yolk internalization by hatchlings is related to breeding timing in the garden lizard, *Calotes versicolor*. *Current Science*, 82 (12), 1484-1485.
- Radder, R. S., Shanbhag, B. A., & Saidapur, S. K. (2004). Yolk partitioning in embryos of the lizard, *Calotes versicolor*: maximize body size or save energy for later use? *Journal of Experimental Zoology Part A: Comparative Experimental Biology*, 301A(9), 783-785. doi:10.1002/jez.a.95
- Rome, L. C., Stevens, E. D., & John-Alder, H. B. (1992). The influence of temperature and thermal acclimation on physiological function. In *Environmental Physiology of the Amphibians* (M.E. Feder and W.W. Burggren ed., pp. 183–205 ). Chicago: University of Chicago Press.
- Santos, R. G., Pinheiro, H. T., Martins, A. S., Riul, P., Bruno, S. C., Janzen, F. J., & Ioannou, C. C. (2016). The anti-predator role of within-nest emergence synchrony in sea turtle hatchlings. *Proceedings of the Royal Society B: Biological Sciences*, 283(1834).
- Sartori, M. R., Taylor, E. W., Abe, A. S. & Crossley II, D. A. (2015). An appraisal of the use of an infrared digital monitoring system for long-term measurement of heart rate in reptilian embryos. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 188, 17-21. <https://doi.org/10.1016/j.cbpa.2015.06.012>.
- Scheiner, S. M. (1993). Genetics and Evolution of Phenotypic Plasticity. *Annual Review of Ecology and Systematics*, 24(1), 35-68. doi:10.1146/annurev.es.24.110193.000343
- Scheiner, S. M. (2001). MANOVA: multiple response variables and multispecies interactions. In *Design and analysis of ecological experiments* (Scheiner SM, Gurevitch J ed., pp. 99-115). New York, NY: Oxford University.

- Scheiner, S. M., & Lyman, R. F. (1991). The genetics of phenotypic plasticity. II. Response to selection. *Journal of Evolutionary Biology*, 4(1), 23-50. doi:10.1046/j.1420-9101.1991.4010023.x
- Sheldon, B. C., Kruuk, L. E. B., Merilä, J., & Crespi, B. (2003). Natural selection and inheritance of breeding time and clutch size in the collared flycatcher. *Evolution*, 57(2), 406-420. doi:10.1554/0014-3820(2003)057[0406:NSAIOB]2.0.CO;2
- Shepherdley, C. A., Daniels, C. B., Orgeig, S., Richardson, S. J., Evans, B. K., & Darras, V. M. (2002). Glucocorticoids, thyroid hormones, and iodothyronine deiodinases in embryonic saltwater crocodiles. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 283(5), 1155-1163. doi:10.1152/ajpregu.00015.2002
- Shine, R. (2004). Adaptive consequences of developmental plasticity. *Reptilian incubation: environment, evolution and behaviour*, 187-210.
- Shine, R., & Elphick, M. J. (2001). The effect of short-term weather fluctuations on temperatures inside lizard nests, and on the phenotypic traits of hatchling lizards. *Biological Journal of the Linnean Society*, 72(4), 555-565. doi:10.1111/j.1095-8312.2001.tb01338.x
- Sih, A., & Moore, R. D. (1993). Delayed hatching of salamander eggs in response to enhanced larval predation risk. *The American Naturalist*, 142(6), 947-960.
- Spencer, R.-J., & Janzen, F. J. (2011). Hatching behavior in turtles. *Integrative and Comparative Biology*, 51(1), 100-110. doi:10.1093/icb/icr045
- Spencer, R.-J., Thompson, M. B., & Banks, P. B. (2001). Hatch or wait? A dilemma in reptilian incubation. *Oikos*, 93(3), 401-406. doi:10.1034/j.1600-0706.2001.930305.x
- Spencer, R. J., & Janzen, F. J. (2014). A novel hypothesis for the adaptive maintenance of environmental sex determination in a turtle. *Proceedings of the Royal Society B: Biological Sciences*, 281(1789), 20140831. doi:10.1098/rspb.2014.0831
- Tazawa, H. (2005). Cardiac rhythms in avian embryos and hatchlings. *Avian and Poultry Biology Reviews*, 16(3), 123-150. doi:10.3184/147020605783437977
- Tazawa, H., Hiraguchi, T., Kuroda, O., Tullett, S. G., & Deeming, D. C. (1991). Embryonic heart rate during development of domesticated birds. *Physiological Zoology*, 64(4), 1002-1022. doi:10.2307/30157954
- Tazawa, H., Kuroda, O., & Whittow, G. C. (1991). Noninvasive determination of embryonic heart rate during hatching in the brown noddy (*Anous stolidus*). *The Auk*, 108(3), 594-601. doi:10.2307/4088100

- Telemeco, R. S., Gangloff, E. J., Cordero, G. A., Mitchell, T. S., Bodensteiner, B. L., Holden, K. G., . . . Janzen, F. J. (2016). Reptile embryos lack the opportunity to thermoregulate by moving within the egg. *The American Naturalist*, 188(1), E13-E27. doi:10.1086/686628
- Thompson, M. B. (1988). Nest temperatures in the Pleurodiran turtle, *Emydura macquarii*. *Copeia*, 1988(4), 996-1000.
- Thompson, M. B. (1989). Patterns of metabolism in embryonic reptiles. *Respiration Physiology*, 76(2), 243-255. doi:10.1016/0034-5687(89)90101-1
- Thompson, M. B. (1997). Egg physiology and biology. In *The Biology, Husbandry And Healthcare of Reptile* (pp. 88-121). New Jersey: TFH Publications Inc.
- Thompson, M. B., Packard, G. C., Packard, M. J., & Rose, B. (1996). Analysis of the nest environment of tuatara *Sphenodon punctatus*. *Journal of Zoology*, 238(2), 239-251. doi:10.1111/j.1469-7998.1996.tb05392.x
- Tucker, J. K., Paukstis, G. L., & Janzen, F. J. (2008). Does predator swamping promote synchronous emergence of turtle hatchlings among nests? *Behavioral Ecology*, 19(1), 35-40. doi:10.1093/beheco/arm097
- Vince, M. A., & Chinn, S. (1971). Effect of accelerated hatching on the initiation of standing and walking in the Japanese quail. *Animal Behaviour*, 19(1), 62-66. doi:http://dx.doi.org/10.1016/S0003-3472(71)80135-5
- Vitt, L. J. (1991). Ecology and life history of the scansorial arboreal lizard *Plica plica* (Iguanidae) in Amazonian Brazil. *Canadian Journal of Zoology*, 69(2), 504-511. doi:doi:10.1139/z91-077
- Vleck, C. M., Hoyt, D. F., & Vleck, D. (1979). Metabolism of avian embryos: patterns in altricial and precocial birds. *Physiological Zoology*, 52(3), 363-377. doi:10.2307/30155757
- Vleck, D. (1987). Measurement of O<sub>2</sub> consumption, CO<sub>2</sub> production, and water vapor production in a closed system. *Journal of Applied Physiology*, 62(5), 2103-2106.
- Vleck, D., Vleck, C. M., & Hoyt, D. F. (1980). Metabolism of avian embryos: ontogeny of oxygen consumption in the rhea and emu. *Physiological Zoology*, 53(2), 125-135. doi:10.2307/30152575
- Wallace, B. P., & Jones, T. T. (2008). What makes marine turtles go: a review of metabolic rates and their consequences. *Journal of Experimental Marine Biology and Ecology*, 356(1-2), 8-24. doi:http://dx.doi.org/10.1016/j.jembe.2007.12.023

- Walther, G.-R., Post, E., Convey, P., Menzel, A., Parmesan, C., Beebee, T. J. C., . . . Bairlein, F. (2002). Ecological responses to recent climate change. *Nature*, *416*, 389. doi:10.1038/416389a
- Warkentin, K. M. (1995). Adaptive plasticity in hatching age: a response to predation risk trade-offs. *Proceedings of the National Academy of Sciences*, *92*(8), 3507-3510. doi:10.1073/pnas.92.8.3507
- Warkentin, K. M. (2000). Wasp predation and wasp-induced hatching of red-eyed treefrog eggs. *Animal Behaviour*, *60*(4), 503-510. doi:10.1006/anbe.2000.1508
- Warkentin, K. M., & Calswell, M. S. (2009). Assessing risk: embryos, information, and escape hatching. . In R. Dukas & J. Ratcliffe (Eds.), *Cognitive ecology II. The evolutionary ecology of learning, memory, and information use*. (pp. 177-200). Chicago: University of Chicago Press.
- Webb, G. J. W., Choquenot, D., & Whitehead, P. J. (1986). Nests, eggs, and embryonic development of *Carettochelys insculpta* (Chelonia: Carettochelidae) from northern Australia. *Journal of Zoology*, *1*(3), 521-550. doi:10.1111/j.1096-3642.1986.tb00646.x
- Weisrock, D. W., & Janzen, F. J. (1999). Thermal and fitness-related consequences of nest location in painted turtles (*Chrysemys picta*). *Functional Ecology*, *13*(1), 94-101. doi:10.1046/j.1365-2435.1999.00288.x
- West-Eberhard, M. J. (1989). Phenotypic plasticity and the origins of diversity. *Annual Review of Ecology and Systematics*, *20*, 249-278.
- West-Eberhard, M. J. (2003). *Developmental Plasticity and Evolution*. Oxford: Oxford University Press.
- Wilbur, H. M. (1975). The evolutionary and mathematical demography of the turtle *Chrysemys picta*. *Ecology*, *56*(1), 64-77. doi:10.2307/1935300
- Wilhoft, D. C. (1958). The Effect of Temperature on Thyroid Histology and Survival in the Lizard, *Sceloporus occidentalis*. *Copeia*, *1958*(4), 265-276. doi:10.2307/1439957
- Willette, D. A. S., Tucker, J. K., & Janzen, F. J. (2005). Linking climate and physiology at the population level for a key life-history stage of turtles. *Canadian Journal of Zoology*, *83*(6), 845-850. doi:10.1139/z05-078

## Figure Legends

**Fig. 1:** Eggs from each clutch were divided and incubated separately for 7 d, then returned to the same containers to continue incubation at 28 °C. Asynchronous group: three more-advanced eggs (black) were incubated at 30°C and six eggs (white) were incubated at 26 °C for 7 d. Synchronous group- three not advanced eggs (black) were incubated at 26 °C and six eggs (white) were incubated at 26 °C for 7 d. All eggs were then placed in 28 °C incubators until the completion of incubation.

**Fig. 2:** Mean incubation period of eggs in asynchronous and synchronous groups,  $\pm$  SE, N = 78. Letters indicate significant pairwise differences among weeks in both asynchronous and synchronous groups, identical letters indicate no difference.

**Fig. 3:** LSMeans of oxygen consumption (mL/h) of embryos in asynchronous (dark grey) and synchronous (light grey) groups from week 4 to 8 of incubation,  $\pm$  SE, n = 10, N = 50. Letters indicate significant pairwise differences among weeks in both asynchronous and synchronous groups.

**Fig. 4:** LSMeans of heart rate (bpm) of embryos in asynchronous (dark grey) and synchronous (light grey) groups from week 3 to 8 of incubation,  $\pm$  SE, n = 10, N = 60. Letters indicate significant pairwise differences among weeks in both asynchronous and synchronous groups.

**Fig. 5:** The difference from mean heart rate of asynchronous (dark grey) and synchronous (light grey) groups over a 24 h period across weeks 3 to 8 of incubation. The x-axis represents duration of time (h) away from the peak heart rate and the y-axis represents the



difference from the mean heart rate (bpm),  $\pm$  SE, N = 62. Letters indicate significant pairwise differences among 6 h time periods in both asynchronous and synchronous groups.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

