

1993

Rapid Communication: MspI Restriction Fragment Length Polymorphism at the Swine Myogenin Locus

C. W. Ernst

Iowa State University

D. A. Vaske

Iowa State University

R. G. Larson

Iowa State University

Max F. Rothschild

Iowa State University, mfrothsc@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/ans_pubs

 Part of the [Agriculture Commons](#), [Animal Sciences Commons](#), and the [Genetics and Genomics Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/ans_pubs/292. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Animal Science at Iowa State University Digital Repository. It has been accepted for inclusion in Animal Science Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Rapid Communication: MspI Restriction Fragment Length Polymorphism at the Swine Myogenin Locus

Abstract

Source and Description of Probe. A 1,486-bp rat cDNA clone for myogenin (MYOG) was excised from the EcoRI site of the plasmid, pBluescribe M13- (Wright et al., 1989). Method of Detection. DNA was isolated from whole blood and digested with MspI. Fragments were then separated by agarose gel electrophoresis and transferred to charged nylon membranes. Hybridizations were at 65°C for 16 to 20 h (.5 M NaCl, .05 M Naphosphate buffer, pH 6.5, 5x Denhardt's reagent, 10% dextran sulfate, .5% SDS, 100 pg/mL sonicated, denatured salmon sperm DNA). Final washes were at 55 to 60°C in .7x SSC, 5% SDS for 15 to 20 min.

Keywords

Porcine, Muscle, Myogenin, RFLP

Disciplines

Agriculture | Animal Sciences | Genetics and Genomics

Comments

This is an article from *Journal of Animal Science* 71 (1993): 3479, doi:/1993.71123479x. Posted with permission.

Rapid Communication: *MspI* Restriction Fragment Length Polymorphism at the Swine Myogenin Locus^{1,2}

C. W. Ernst³, D. A. Vaske, R. G. Larson, and M. F. Rothschild⁴

Department of Animal Science, Iowa State University, Ames 50011-3135

Source and Description of Probe. A 1,486-bp rat cDNA clone for myogenin (MYOG) was excised from the *EcoRI* site of the plasmid, pBluescribe M13⁻ (Wright et al., 1989).

Method of Detection. DNA was isolated from whole blood and digested with *MspI*. Fragments were then separated by agarose gel electrophoresis and transferred to charged nylon membranes. Hybridizations were at 65°C for 16 to 20 h (.5 M NaCl, .05 M Naphosphate buffer, pH 6.5, 5× Denhardt's reagent, 10% dextran sulfate, .5% SDS, 100 µg/mL sonicated, denatured salmon sperm DNA). Final washes were at 55 to 60°C in .7× SSC, .5% SDS for 15 to 20 min.

Description of Polymorphism. Hybridization with the rat MYOG probe revealed three swine fragments (Figure 1). The 4.9- and 4.2-kb fragments were polymorphic, and the 2.3-kb fragment was monomorphic. No polymorphisms were detected for swine MYOG in unrelated pigs using *TaqI* (34 pigs), *SacI* (34 pigs), *PstI* (31 pigs), *BamHI* (16 pigs), or *PvuII* (15 pigs).

Inheritance Pattern. Autosomal Mendelian segregation of the 4.9- and 4.2-kb swine MYOG fragments was observed in 29 pigs from four, two- and three-generation families.

Frequency. Analysis of 83 unrelated pigs from nine breeds indicated overall allelic frequencies of .58 for the 4.2-kb fragment and .42 for the 4.9-kb fragment (Table 1).

Chromosomal Location. Unknown.

Comments. Myogenin is a member of a family of transcription factors that are specific to skeletal muscle (Wright et al., 1989). Myogenin is involved in regulating the expression of genes associated with the determination and differentiation of myoblasts.

Literature Cited

Wright, W. E., D. A. Sassoon, and V. K. Lin. 1989. Myogenin, a factor regulating myogenesis, has a domain homologous to MyoD. *Cell* 56:607.

Key Words: Porcine, Muscle, Myogenin, RFLP

J. Anim. Sci. 1993. 71:3479




Figure 1. Mendelian segregation of porcine myogenin *MspI* fragments in a two-generation crossbred family of Meishan × Hampshire pigs. Square denotes sire; circle denotes dam.

Table 1. Frequency of Myogenin genotypes in several breeds

Breed	no.	Frequency of pigs with indicated genotype ^a		
		4.9/4.9	4.9/4.2	4.2/4.2
Fengjing	6	100	0	0
Meishan	8	75	25	0
Minzhu	4	50	25	25
Chester White	9	56	33	11
Landrace	13	8	46	46
Yorkshire	9	0	33	67
Duroc	14	7	29	64
Hampshire	12	0	8	92
Poland China	8	25	50	25
Total	83			

^a4.9 = 4.9-kb *MspI* fragment; 4.2 = 4.2-kb *MspI* fragment.

¹Journal paper no. J-15477 of the Iowa Agric. and Home Econ. Exp. Sta., Ames. Projects no. 3083 and 3043.

²This project is supported in part by the National Pork Producers Council, Iowa Pork Producers Association, and the Ohio Pork Producers Council. The authors wish to thank J. Newton for blood collection, C. Hergenrader and S. Kenealy for DNA preparations, and F. C. Ernst, Clear Spring, MD for Poland China samples. The rat MYOG cDNA was kindly provided by Woodring Wright, The University of Texas Southwestern Medical Center.

³Present address: Dept. of Anim. Sci., The Ohio State Univ., Columbus 43210.

⁴To whom correspondence should be addressed.

Received July 29, 1993.

Accepted September 24, 1993.