

2019

Fish growth changes over time in a Midwestern USA lake

Eric D. Katzenmeyer
Iowa State University

Michael E. Colvin
Iowa State University

Timothy W. Stewart
Iowa State University, twstewar@iastate.edu

Clay L. Pierce
U.S. Geological Survey, cpierce@iastate.edu

Scott E. Grummer
Iowa Department of Natural Resources

Follow this and additional works at: https://lib.dr.iastate.edu/nrem_pubs

 Part of the [Animal Sciences Commons](#), [Natural Resources Management and Policy Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/nrem_pubs/304. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

Fish growth changes over time in a Midwestern USA lake

Abstract

Growth of Walleye *Sander vitreus*, Yellow Bass *Morone mississippiensis*, Common Carp *Cyprinus carpio*, and Black Bullhead *Ameiurus melas* was assessed in Clear Lake, Iowa, USA, over several decades and in relation to environmental variables. Growth of Common Carp was positively correlated with phytoplankton concentration. Recent Black Bullhead growth was faster than in the 1950s and 1990s, which may be a consequence of their recent decline in abundance. Growth of Common Carp and Yellow Bass was faster in the 1940s than in more recent time periods. Relative to their entire range, Common Carp first year growth was below average whereas length at later ages was above average. Walleye relative growth showed a similar pattern. The large changes in growth over several decades suggest that as the Clear Lake ecosystem continues to change, growth rates of its important fish species are also likely to continue changing.

Disciplines

Animal Sciences | Ecology and Evolutionary Biology | Natural Resources Management and Policy | Terrestrial and Aquatic Ecology

Comments

This is a manuscript of an article published as Katzenmeyer, Eric D., Michael E. Colvin, Timothy W. Stewart, Clay L. Pierce, and Scott E. Grummer. "Fish growth changes over time in a Midwestern USA lake." *Transactions of the American Fisheries Society* (2019). doi: [10.1002/tafs.10149](https://doi.org/10.1002/tafs.10149).

Rights

Works produced by employees of the U.S. Government as part of their official duties are not copyrighted within the U.S. The content of this document is not copyrighted.

DR. CLAY PIERCE (Orcid ID : 0000-0001-5088-5431)

Article type : Article

Fish growth changes over time in a Midwestern USA lake †

Eric D. Katzenmeyer^{1, a}, Michael E. Colvin^{1, b}, Timothy W. Stewart¹, Clay L. Pierce^{2, *}, and
Scott E. Grummer³

¹Department of Natural Resource Ecology and Management, Iowa State University, Ames,
Iowa 50011-1031, USA

²U.S. Geological Survey, Iowa Cooperative Fish and Wildlife Research Unit, Iowa State
University, Ames, Iowa 50011-1031, USA

³Iowa Department of Natural Resources, 1203 North Shore Drive, Clear Lake, Iowa 50428,
USA

^aPresent address: Minnesota Department of Natural Resources, 20596 Highway 7,
Hutchinson, MN 55350, USA

^bPresent address: Department of Wildlife, Fisheries & Aquaculture, Mississippi State
University, Box 9690, Mississippi State, MS 39762-9690, USA

*Corresponding author: e-mail: cpierce@iastate.edu

†This draft manuscript is distributed solely for purposes of scientific review. Its content is
deliberative and predecisional, so it must not be disclosed or released by reviewers. Because
the manuscript has not yet been approved for publication by the U.S. Geological Survey
(USGS), it does not represent any official USGS finding or policy.

This article has been accepted for publication and undergone full peer review but has not
been through the copyediting, typesetting, pagination and proofreading process, which may
lead to differences between this version and the Version of Record. Please cite this article as
doi: 10.1002/tafs.10149

This article is protected by copyright. All rights reserved.

[A]Abstract

Growth of Walleye *Sander vitreus*, Yellow Bass *Morone mississippiensis*, Common Carp *Cyprinus carpio*, and Black Bullhead *Ameiurus melas* was assessed in Clear Lake, Iowa, USA, over several decades and in relation to environmental variables. Growth of Common Carp was positively correlated with phytoplankton concentration. Recent Black Bullhead growth was faster than in the 1950s and 1990s, which may be a consequence of their recent decline in abundance. Growth of Common Carp and Yellow Bass was faster in the 1940s than in more recent time periods. Relative to their entire range, Common Carp first year growth was below average whereas length at later ages was above average. Walleye relative growth showed a similar pattern. The large changes in growth over several decades suggest that as the Clear Lake ecosystem continues to change, growth rates of its important fish species are also likely to continue changing.

Growth is an important component of fish population dynamics, and knowledge of growth rates and understanding their determinants are essential to fisheries management (Summerfelt and Hall 1987; Weatherly and Gill 1987). Growth in the first year is important in determining eventual recruitment, as growth rate determines body size, which in turn has consequences for survival via size-dependent foraging success and vulnerability to predation (Miller et al. 1988; Chambers and Trippel 1997). Growth can be an indicator of potential management problems (e.g., overfishing, lack of food) and provide feedback on management actions, so understanding what factors influence growth is an important part of successful fishery management (Van Den Avyle and Hayward 1999). Several factors have been shown to influence fish growth, including temperature (Kelso 1972; Momot et al. 1977; Pauly 1980;

Staggs and Otis 1996), food availability (Welker et al. 1994; DeVries et al. 1998; Bremigan et al. 2003), and competition (Ridenhour 1960; Welker et al. 1994).

Clear Lake is Iowa's third-largest natural lake and is a highly valued natural resource for angling, boating, and other outdoor recreation (Downing et al. 2001). Over 432,000 people visit Clear Lake annually, and economic value for vacation and recreational use exceeds US\$43 million annually (Center for Agriculture and Rural Development 2008). A recreational fishery, primarily for Yellow Bass *Morone mississippiensis* and Walleye *Sander vitreus*, is valued between US\$1-2.5 million annually (Iowa Department of Natural Resources, unpublished data). In the past 60 years, Clear Lake has experienced increases in suspended sediment, nutrients, and phytoplankton concentrations, while water clarity, lake depth, and submerged macrophytes have all decreased (Downing et al. 2001; Anthony and Downin, 2003; Egertson et al. 2004). Common Carp *Cyprinus carpio* have been present since the early 1900s (Bailey and Harrison 1945) but were not abundant until recent decades (Larscheid 2005; Colvin et al. 2010). Zebra mussels *Dreissena polymorpha* were first discovered in Clear Lake in 2005, and densities have since increased dramatically (Colvin et al. 2010). In response to long-term declines in water quality, management actions have been implemented to improve water quality and the fishery, including in-lake best management practices such as Common Carp removals (Colvin et al. 2012), dredging and shoreline stabilization, urban best management practices such as native landscaping, permeable pavement and bio-retention cells, and agricultural best management practices such as cover crops, grassed waterways, terraces and conversion of agricultural land to the Conservation Reserve Program (Iowa Department of Natural Resources 2008; Clear Lake Enhancement and Restoration Project 2016).

Two approaches were used for quantifying and comparing growth rates (size-specific) and lengths attained (age-specific) among years and time periods for four fish species in Clear Lake. Size-specific analyses relate growth to body size rather than age (Gutreuter 1987; Putman et al. 1995; Pierce et al. 2003; Shoup et al. 2007; Quist et al. 2012). Because fish of the same age can vary greatly in size, age-specific analyses can result in comparing growth rates among groups of fish that differ significantly in preferred prey, metabolic rates, habitat requirements, risk of predation, and other important factors (Larkin et al. 1956; Osenberg et al. 1988). Although they are often referred to as “growth”, age-specific analyses actually yield estimates of attained lengths, which are the sums of growth in all previous years, and thus are not comparable with annual estimates of environmental variables as size-specific analyses are. Despite containing influences of different numbers of years, the length at age estimates resulting from age-specific analyses are useful because of their relative ease in computation, their widespread application in fisheries management, and their incorporation in quantitative methods such as the von Bertalanffy model and many associated techniques (Slipke and Maceina 2014). Although biases due to unknown yearly differences are likely when using age-specific analyses to make comparisons among adjacent years over short time frames, comparing time periods separated by longer intervals such as in this analysis of historical data reduces the effect of these biases and provides a useful way to compare growth among different periods widely separated in time (Millner and Whiting 1996). Similarly, Copp et al. (2004) demonstrated the utility of this approach for examining intercontinental growth differences in pumpkinseed *Lepomis gibbosus*.

The purpose of this study was to examine growth changes of four common fish species in Clear Lake - Walleye, Yellow Bass, Common Carp, and Black Bullhead *Ameiurus melas* - over time. Common Carp, Yellow Bass, and Walleye are the three most abundant fish

species in Clear Lake (Colvin et al. 2010). Black Bullhead were recently among the most abundant but have declined in the last two decades (Larscheid 2005). The first objective was to document annual growth rates over several years of the four fish species in their first year of life and at the length at which they are maturing. The second objective was to examine relationships between fish growth and environmental variables. The third objective was to compare recent growth to historical growth documented in previous studies. The third objective provides insight into how fish growth has changed over time as habitats and water quality have changed.

[A]Methods

[C]*Study site.*—Clear Lake is a shallow, eutrophic lake located in Cerro Gordo County in north central Iowa, USA (43°08'N, 93°22'W). It has a surface area of 1474 hectares, a watershed area of 4888 hectares, a mean depth of 3.1 m, and a maximum depth of 7.3 m. Inorganic substrates cover the vast majority of the bottom, with silt accounting for 92%, sand 4%, and rock 2% of the area. Historically, Clear Lake supported abundant aquatic vegetation, but due to poor water clarity in recent years, aquatic vegetation cover has been reduced to 2% of the bottom area and is also reduced in diversity (Downing et al. 2001; Egertson et al. 2004; Colvin et al. 2010).

The predominant fish species include Yellow Bass, Walleye, and Common Carp (Colvin et al. 2010). This is in sharp contrast to the historical fish community that contained abundant Largemouth Bass *Micropterus salmoides*, Bluegill *Lepomis macrochirus*, and Northern Pike *Esox lucius* (Bailey and Harrison 1945). Walleye have been stocked nearly every year since 1915 (Bailey and Harrison 1945; Carlander and Payne 1977). Fish species diversity has declined since 1945 when 43 species were recorded (Bailey and Harrison 1945),

including several darter and cyprinid species that are no longer present. Currently, Clear Lake supports 23 fish species (Colvin et al. 2010).

[C]*Fish and aging structure collection.*—Fish sampling was conducted as part of another study in 2007 and 2008 using a variety of methods (Colvin et al. 2010). A 132 m long beach seine, 3.5 m in depth with a 3.5 m² bag in the center and 6 mm square mesh, was used to sample the nearshore fish community in September. The offshore fish community was sampled using a semi-balloon otter trawl with an 8 m head rope, 3.8 cm stretch mesh body, and 6.3 mm mesh cod end, also in September. Night electrofishing was conducted in October and November to obtain a larger sample of Walleye. Common Carp were captured by commercial fisherman using seines in May and November of 2007 and June and November of 2008. Sex of collected fish was not determined.

Dorsal spines were used for aging and back-calculation of lengths at previous ages for Common Carp and Walleye, pectoral spines for Black Bullhead, and sagittal otoliths for Yellow Bass based on criteria discussed in Quist et al. (2012). Upon capture, Common Carp, Walleye and Black Bullhead were measured for total length (mm), an aging structure was removed and the fish were released alive. Yellow Bass were captured in greater numbers and were subsampled randomly within 2.5 cm length groups and frozen or preserved in formalin for later length measurement and extraction of aging structures. Common Carp and Walleye dorsal spines were removed by cutting the spine as close to the base as possible with a side cutter. Black Bullhead pectoral spines were removed by laying the spine flat against the fish's body and rotating it dorsally until the articulating process separated from the joint. Spines were placed singly in scale envelopes and allowed to air dry. Sagittal otoliths were removed from Yellow Bass in the lab by making a dorso-ventral cut from the top of the head through

the preopercle and bending the head away from the body to expose the otoliths (Secor et al. 1991). Otoliths were rinsed in water, placed in a scale envelope and allowed to dry.

[C]*Aging structure preparation.*—Spines were sectioned for viewing with a Buehler Isomet low speed saw (Lake Bluff, Illinois, USA) at a thickness of 0.8-1.0 mm using a diamond wafering blade. Large Common Carp spines were sectioned without additional preparation beyond drying. Small Common Carp dorsal spines, Walleye dorsal spines and Black Bullhead pectoral spines were mounted in epoxy for sectioning (Koch and Quist 2007).

Large Yellow Bass otoliths were broken in half along the dorso-ventral axis and sanded, first with 400 grain sandpaper until the nucleus was reached, and then polished with 1200 grit sandpaper (Secor et al. 1991). The otolith half was pushed into black clay (polished surface up) and covered with a drop of immersion oil for viewing. Small Yellow Bass otoliths that were too small to break in half were mounted in epoxy as described above and sectioned as close to the nucleus as possible and polished, first with 400 grain sandpaper until the nucleus was reached and then with 1200 grain sandpaper (Secor et al. 1991). These were also pushed into black clay and covered with immersion oil for viewing.

[C]*Aging structure measurement and back-calculation.*—Aging structures were examined under a dissecting microscope (8x – 115x magnification) fitted with a computerized video image analysis system (Image Pro 6.2, Media Cybernetics, Rockville, Maryland, USA). A still image (2560 x 1920 pixels) of each structure was saved for aging and measurement. The structures were aged and measurements of distance between annuli were made independently by two readers without knowledge of the size of the fish. Measurements were taken along the longest axis of the structure from the origin of the

structure to the edge for dorsal and pectoral spines. Yellow Bass otoliths were measured from the nucleus to the edge along the sulcal groove. When there was a discrepancy between the two readers, the structure was viewed together and a consensus was reached. If a consensus could not be reached the structure was excluded from analysis.

Back-calculation of length at previous ages was done using the Dahl-Lee (direct proportion) method (Quist et al. 2012). The Dahl-Lee formula is

$$L_i = (S_i/S_c)L_c, \quad (1)$$

where L_i = back-calculated fish body length at age i , L_c = fish body length at capture, S_i = mean aging structure length at annulus i , and S_c = mean aging structure total length. Back-calculated lengths at age were thus the average from the two independent readers for each fish.

[C]*Size-specific analysis of growth.*—A size-specific (Quist et al. 2012) approach was used to analyze growth of the four fish species. Annual growth for a given year was calculated as the difference between two back-calculated lengths at age from consecutive years. The year in which the aging structure was collected was excluded from analysis because the fish had not yet experienced a full growing season. Annual growth was then plotted against length at the beginning of the growing season. Estimates of length at hatching were obtained from the literature (Holland-Bartels et al. 1990) and used as the length at the beginning of the growing season for fish in their first year of life. For Black Bullhead and Walleye 7 mm was used for length at hatching, for Common Carp 3.5 mm, and 3 mm for Yellow Bass. For older ages, back-calculated length at the beginning of the growing season was used. Locally weighted scatterplot smoothing (LOWESS) regression was used to fit curves to the annual growth estimates (Pierce et al. 2003). The “loess” function in Program R

(R Development Core Team 2009) was used for these analyses. LOWESS is a nonparametric regression method appropriate where the data suggest no particular parametric model form (Cleveland 1979). After testing several different smoothing parameters (span in Program R) for LOWESS, a smoothing parameter that worked best for each species individually was used. For Common Carp and Yellow Bass a span = 1 and for Walleye and Black Bullhead a span = 1.2 were used. These parameters provided accurate fits to the data that represent the growth patterns in the dataset, while not allowing overpowering influence of individual variation and outliers.

Mean annual growth at age 0 (AG-0) and mean annual growth at length of maturation (AG-m) were chosen as expressions of growth for size-specific comparisons because they represent two important stages (AG-0) or events (AG-m) in the life cycle of fishes. LOWESS regressions represent “average” annual growth of fish at all sizes in a given year, therefore allowing estimation of growth for any size fish in that year. Age of maturity estimates were obtained from the literature (Carlander 1969, 1997) and averaged back-calculated lengths at that age for each fish species were used to estimate length at maturity for each species. Ages at maturity used were age 3 for Black Bullhead and Walleye and age 2 for Common Carp and Yellow Bass. Actual age and length at maturity can vary among sexes, water bodies, with changing environmental conditions, as well as with changes in exploitation (Hutchings and Jones 1998; Stearns 2000), so in nature they are not constants as literature-derived values imply. Therefore, we henceforth refer to our estimated “adult” growth rates as growth at length of maturation, rather than growth at length of maturity. Because age 0 fish begin growing at a fairly consistent length at hatching, age-specific growth and size-specific growth are essentially identical in the first year. Thus, AG-0 was estimated by averaging length at

age 1. Estimates of AG-m were obtained from LOWESS regressions, solving for annual growth at length at maturation.

[C]*Growth relationships with environmental variables.*—Five environmental variables were analyzed for relationships with growth. The annual number of degree days where water temperature exceeded 5°C (hereafter referred to as degree days) was used as an index of temperature because it encompasses most of the open water season and the temperature range where most fish growth occurs. Walleye do not gain weight at temperatures below 6°C (Hokanson 1977); thus, it was assumed that 5°C and greater should encompass temperature conditions favorable for growth of the four species. Air temperature data from the nearest station (Mason City, Iowa, USA) were obtained from the National Oceanic and Atmospheric Administration (2009), and from these data daily mean water temperatures were estimated using an air/water relationship developed for Clear Lake (Jacobsen 1968). Using 5°C as a base temperature, for each day that the daily mean water temperature exceeded 5°C in a given year, 5 was subtracted from the daily mean temperature and these quantities were summed over days to give the annual degree days. Mean water transparency, chlorophyll *a* concentration, phytoplankton concentration and zooplankton concentration were obtained from the Iowa Lakes Information System (ILIS) (Iowa Department of Natural Resources, 2009) and were based on three sampling periods at approximately monthly intervals during the summer of each year. Water transparency is an indicator of overall water quality and lake productivity (Carlson 1977). Chlorophyll *a* concentration is an indicator of abundance of phytoplankton, which is the trophic base of aquatic ecosystems (McCauley and Kalff 1981; Kalff 2002). Zooplankton are important in the diets of many fish species, especially young of the year (Chambers and Trippel 1997).

Relationships between size-specific growth and environmental variables were examined using Pearson correlation coefficients.

[C]Age-specific growth comparisons with historical data.—Recent fish growth estimated as described above in the 2000s (1990s – 2000s for Common Carp) (hereafter referred to as “Recent”) was compared with historical growth documented in previous time periods in the literature (Carlander 1969, 1997; Larscheid 2005; see Table S1 in Supporting Information for details). Literature data consisted of mean lengths at age with no estimates of error. Where only single studies were available from a time period (Black Bullhead, Common Carp), growth curves were plotted without estimates of error. For species where multiple years of length at age data were available for a time period (Walleye, Yellow Bass), means for each age were calculated by decade and a single growth curve with 95% confidence intervals for each age was plotted. The length at age 4 estimate for Black Bullhead from 1955 was less than the age 3 estimate and was likely erroneous; this estimate was omitted from further analysis.

For each time period L_{∞} , k and t_0 were estimated from the von Bertalanffy growth model,

$$l(t) = L_{\infty}[1 - e^{-k(t - t_0)}], \quad (2)$$

where L_{∞} is the theoretical maximum length a fish will reach in the population, k is a growth constant and t_0 is the time at which the fish’s length is zero (Quist et al. 2012). In addition, ω , which is roughly equivalent to the product of L_{∞} and k and corresponds to the growth rate near t_0 , was estimated. ω is recommended for statistical comparison (Gallucci and Quinn 1979) because of the strong negative correlation of L_{∞} and k . The original von Bertalanffy growth model can be re-parameterized using ω , and the equation is,

$$l(t) = \omega/k [1 - e^{-k(t - t_0)}]. \quad (3)$$

ω was compared across time periods for each species using pairwise *t*-tests (Katzenmeyer 2010).

For Common Carp and Walleye the approach of Jackson et al. (2008) and Quist et al. (2003) was followed to convert lengths at age to the relative growth index (RGI). This allowed comparisons of lengths attained in Clear Lake to the entire species' ranges, which may be useful for interpreting results in Clear Lake and elsewhere, as well as identification of factors contributing to slow growth and helping guide management actions (Jackson et al. 2008). RGI analysis was not attempted for Black Bullhead and Yellow Bass because length at age standards have not been developed for these species.

[A]Results

[B]Environmental Variables

Annual degree days ranged from 2,372 to 2,892 (Table 1). Annual mean summer water transparency, chlorophyll *a*, phytoplankton and zooplankton concentrations were not monitored before 2000, but from 2000-2007 transparency varied over three-fold, ranging from 0.4 to 1.4 m, chlorophyll *a* concentration varied roughly four-fold, ranging from 20.7 to 82 $\mu\text{g/L}$, phytoplankton concentration varied by over two orders of magnitude, ranging from 14.6 to 3112.7 mg/L , and zooplankton concentration varied by over an order of magnitude, ranging from 23.4 to 276.1 $\mu\text{g/L}$. Although there was substantial year-to-year variability in the five measured variables, no clear temporal trend was evident for any variable during the period when measurements were recorded.

[B]Size-Specific Growth

Aging structures were examined from 408 Common Carp, 222 Walleye, 102 Yellow Bass and 61 Black Bullhead. Size-specific growth curves fit by LOWESS regression and resulting AG-0 and AG-m estimates are shown in Supporting Information (Figures S1-S5). Black Bullhead first year growth (AG-0) ranged from 92 to 150 mm and annual growth at length at maturation (AG-m) ranged from 30 to 43 mm, Common Carp AG-0 ranged from 88 to 182 mm and AG-m ranged from 84 to 143 mm, Walleye AG-0 ranged from 111 to 212 mm and AG-m ranged from 43 to 64 mm, and Yellow Bass AG-0 ranged from 81 to 100 mm and AG-m ranged from 24 to 52 mm (Figure 1). First year growth was greater than annual growth at length at maturation for Black Bullhead, Walleye and Yellow Bass, but was similar for both life cycle stages for Common Carp.

Black Bullhead, Common Carp and Yellow Bass AG-0 gradually increased during the study, with an average increase of 58% over the years for which estimates were possible (Figure 1). Walleye AG-0 increased each year from 2002 through 2005 but then declined the next two years. Common Carp and Yellow Bass AG-m mirrored their AG-0 trends, increasing an average of 50% over the years of available data. Although there was year to year variation there were no apparent trends in Black Bullhead and Walleye AG-m.

There were relatively few significant relationships for size-specific growth variables among species, and small sample sizes (i.e., years with corresponding annual growth estimates) for correlation analysis suggest interpreting these relationships with caution. Black Bullhead AG-0 was positively correlated with Walleye and Yellow Bass AG-0 (Table 2). Although not significant at $\alpha=0.05$, Common Carp AG-0 and AG-m values appeared to be positively associated with corresponding Yellow Bass values, and Common Carp AG-m

appeared to be negatively associated with Walleye AG-0 ($0.1 > p > 0.05$). AG-0 and AG-m were not significantly correlated within any of the four species.

There were also relatively few significant relationships between growth variables and the environmental variables that were quantified, and caution due to small samples sizes applies here as well. Common Carp AG-0 and AG-m were positively correlated with phytoplankton concentration (Table 3). Although not significant at $\alpha=0.05$, Walleye AG-0 appeared to be positively associated with chlorophyll a, Yellow Bass AG-0 was positively associated with phytoplankton concentration, and Walleye AG-m was negatively associated with transparency ($0.1 > p > 0.05$; Table 3).

[B]Age-Specific Growth Comparisons with Historical Data

Differences in lengths at age between study periods were evident to varying degrees for all four species (Figures 2, 3). Black Bullhead length at age was greater in the present study than in earlier studies (Figure 2), and this was supported by significantly larger ω in the present study than in previous studies (Table 4). Common Carp length at age was much greater in the 1952 study than in the late 1990s or the present study, and this was supported by significantly larger ω (Table 4) and consistently higher RGI values (Figure 3) in the 1952 study than the two later studies. RGI values for age 2 and older fish were above 100% of “average” for Common Carp throughout its range. RGI values for age 1 Common Carp were 30 to 50% lower than for older ages in all three time periods. Walleye lengths at age were broadly similar across the three time periods, although lengths increased fastest in years 2 and 3 in the present study, and lengths beyond age 6 in the 1948-1974 time period were slightly lower than in the other time periods (Figure 2). As expressed by ω , growth was significantly faster in the present study than in previous studies (Table 4), but this was largely due to faster

early growth and a larger k in the present study and did not result in substantially greater lengths at age, especially compared to the 1940s (Figure 2). There was a general trend in all three time periods for the Walleye RGI to increase with age, with younger ages at or below average lengths and older ages above average lengths for the species (Figure 3). Lengths of age 1 Walleye in the present study were lower than in the earlier time periods (Figure 3). Yellow Bass attained significantly greater lengths at older ages in the 1940s than in the two more recent time periods (Figure 2). In the present study growth in the first 2 years was rapid, resulting in the greatest lengths at age in the first two years among the three time periods, but thereafter it slowed considerably, resulting in the shortest lengths at age among the three time periods in years 4 through 6 (Figure 2). As expressed by ω , Yellow Bass growth was significantly lower during the 1960s than during the 1940s and the present study, which did not differ (Table 4).

[A]Discussion

Patterns of Black Bullhead growth evident in both the age-specific and size-specific analyses suggest a potential role for density dependence. Black Bullhead grew slower in the 1950s and 1999 than was documented recently. Black Bullhead were described as being very abundant in the 1940s (Bailey and Harrison 1945) and 1950s (Forney 1955), and a quantitative study showed them to be the lake's most abundant species in 1959 at a biomass of 49 kg/ha (McCann 1960). In 1999, biomass of Black Bullhead was estimated to be 336 kg/ha (Larscheid 2005). In sharp contrast, biomass of Black Bullhead averaged only 10 kg/ha recently (Colvin *et al.*, 2010). Lengths at age and ω from the three time periods were negatively related to biomass, suggesting the possibility of intraspecific competition. The recent precipitous decline in Black Bullhead biomass in Clear Lake could have released them from intraspecific competition for food, and is a plausible explanation for the recent increase

in their growth. The steady increase in first-year growth of Black Bullhead from the early to middle first decade of the 2000s is consistent with this interpretation. Previous studies in Iowa (Rose and Moen 1950), Oklahoma (Houser and Collins 1962), and South Dakota (Hanchin et al. 2002) have likewise concluded that Black Bullhead growth is strongly regulated by intraspecific competition, so it is possible that density dependence may frequently be an important determinant of Black Bullhead growth. Numerous other factors have changed in Clear Lake since the 1950s (Downing et al. 2001; Egertson et al. 2004) that may have also played a role in the changes in Black Bullhead growth documented here.

Common Carp attained markedly greater lengths at age in the 1950s than in the two more recent time periods examined in this study. Unfortunately, no comparable data on Common Carp abundance and few data on environmental conditions exist across the three time periods, making interpretation difficult. Common Carp were described as “common” in the early 1940s (Bailey and Harrison 1945) and as “abundant” in the late 1940s and early 1950s (Parsons 1950; English 1951). Recent studies documented a highly abundant Common Carp population in the early 2000s (Larscheid 2005; Colvin et al. 2012). The largest annual removal of Common Carp since nuisance fish removals began in 1929 was the 111,427 kg (76 kg/ha) removed in 1949 (Iowa Department of Natural Resources, unpublished data), which is roughly half of the average population biomass estimated during 2007 to 2010 (Colvin et al. 2012), and would most likely have reduced biomass of the remaining population significantly for a few years. The studies published in the 1950s, during the time that corresponded with the 1952 length at age estimates, speculated that Common Carp played a significant role in the declines of submerged vegetation and water clarity during that time period (Parsons 1950; English 1951). It is reasonable to further speculate that the more abundant vegetation and likely lower population density and biomass in the 1950s may have

enabled faster growth of Common Carp than in more recent time periods when reduced plant abundance and water transparency were the norm.

Clear Lake presents distinctly different growing conditions to juvenile and adult Common Carp relative to the rest of the species' range, as evidenced by the distinctly lower RGI values for age 1 Common Carp in all three time periods, implying slower first year growth relative to populations elsewhere, and contrasting faster relative growth experienced at older ages. Extensive areas of shallow, vegetated, marsh-like habitat, shown in other studies (Koehn 2004; Stuart and Jones 2006; Bajer and Sorensen 2009) to be important nursery habitat for juvenile Common Carp, are scarce in Clear Lake (Egertson et al. 2004; Penne and Pierce 2008), although juvenile Common Carp seek these limited shallow, vegetated areas (Penne and Pierce 2008). Relative to their growth elsewhere, the slow growth of Common Carp in their first year may indicate that the environment of Clear Lake, currently supporting limited shallow, vegetated habitat, is unfavorable for juvenile growth. In turn, the greater than average RGI values in subsequent years implies very fast growth in the second year and at least average growth in succeeding years. This dichotomy in growth conditions relative to the species' range between the first and subsequent years in the life history of Common Carp suggests that in addition to continuing efforts to reduce adult Common Carp biomass by commercial-scale netting (Colvin et al. 2012), we suggest that future comprehensive control strategies seek to exploit the poor growing conditions for juveniles.

Although our sampling and analytical methods had the potential to reveal patterns of annual growth with environmental variables, small sample sizes limited the power of many of these correlations and suggests that all should be interpreted with caution. The significant positive correlations between growth of both Common Carp life cycle stages and phytoplankton abundance suggests a relationship with algal production in the water column. In addition to contributing to Common Carp diets in the form of settled organic detritus (Panek, 1987), phytoplankton abundance has been shown in a number of studies to be positively correlated with benthic invertebrate abundance (Dermott et al. 1977; Davies 1980; Blumenshine et al. 1997), which typically make up the majority of Common Carp diets. Increased phytoplankton concentration is frequently observed when Common Carp invade new water bodies and become established (Weber and Brown 2009), and thus the initial ecosystem impacts of Common Carp invasion may serve as a positive feedback mechanism through enhanced growth.

Although the length at age plots for Walleye in the three time periods examined are similar, growth as reflected by ω was faster in the present study than in the previous periods. The primary difference was much faster growth in the second year of life and a larger k in the present study. Explanation for this pattern is elusive, but in addition to natural influences, early growth phenomena exhibited by Walleye in Clear Lake could also be consequences of stocking. Natural reproduction of Walleye in Clear Lake is believed to be poor, and therefore Walleye have been stocked nearly every year since 1915 (Bailey and Harrison 1945; Bulkley et al. 1976; Carlander and Payne 1977). Historically all Walleye were stocked as fry (Carlander and Payne 1977), but in the last two decades Clear Lake has been stocked with large fingerlings (150-200 mm) as well as fry. Fish stocked at a larger size typically have higher survival and recruitment because they are less susceptible to predation and other

stresses (Santucci and Wahl 1993). In Clear Lake, Walleye fry are stocked in the early spring and fingerlings are stocked in September and October. Fingerlings attain a much larger size (approximately 175-200 mm) at the end of the first growing season than fry (approximately 100 mm). Pratt and Fox (2003) found that when two sizes of fingerlings were stocked, the two size groups converged four years later. Other studies have shown that the size difference persisted throughout life or that the small fingerlings grew faster (Olson et al. 2000; Brooks et al. 2002). Changes in stocking practices over time complicate comparisons of Walleye growth across time periods, especially in early age classes. The results of this study suggest that young Walleye are currently growing as fast or faster than in earlier times and are attaining adult sizes similar to or larger than in earlier times, although changes in stocking practices, differential survival of fry- and fingerling-stocked Walleye between years, and differential growth of surviving fry- and fingerling-stocked Walleye may account for some of these apparent differences.

Conditions for Walleye growth improve with age in Clear Lake relative to Walleye throughout their range. This pattern was evident in all three time periods, suggesting that factors that enable Clear Lake Walleye to grow increasingly better later in life than in many other locations have been at work for decades. Walleye are maintained in Clear Lake through yearly, intensive stocking (Carlander and Payne 1977), and the fishery and its associated harvest of adults, occurring in both winter and summer, have been popular for over a century (Wahl 2001). Regardless of current food web structure and in-lake nutrient dynamics, Clear Lake is located in one of the most nutrient-rich landscapes in the world (Downing et al. 2001). Our results suggest that future Walleye management strategies seek to enhance survival of juvenile and early adult Walleye to allow more fish to survive to older ages and capitalize on the inherently faster growth of older, larger fish.

The age-specific analysis showed that Yellow Bass growth has slowed since the 1940s in Clear Lake. Yellow Bass are native to the Mississippi River and its drainage (Harlan and Speaker 1987) but were likely introduced to Clear Lake in the 1920s and first showed up in anglers' catches in 1932 (Bailey and Harrison 1945). As is the case with many introduced species, Yellow Bass may have experienced their most rapid growth shortly after introduction. Ridenhour (1960) found evidence of intraspecific competition among Yellow Bass in Clear Lake, showing that in years when abundance was greater, growth tended to be slower than in years when abundance was low. Presently, Yellow Bass are the most abundant fish species in Clear Lake (Colvin et al. 2010) and intraspecific competition likely contributes to their reduction in growth since the 1940s.

The potential value of the increased temporal resolution of size-specific analysis relative to age-specific analysis is illustrated by the recent trend in Yellow Bass annual growth. Although sufficient data to rigorously examine relationships of fish growth with zebra mussel abundance were lacking, the doubling observed in the annual growth at length at maturation in Yellow Bass during the first three years (2004-2007) of zebra mussel colonization in Clear Lake may have been related to the coinciding increase in zebra mussels (Colvin et al. 2010). Stewart et al. (1998) showed that increased benthic habitat provided by zebra mussel shells led to increases in several benthic macroinvertebrate taxa, including hydrids, turbellarians, gastropods, amphipods and chironomids. Many of these taxa, especially amphipods and chironomids, commonly make up large proportions of the diet of Yellow Bass in Clear Lake (Welker 1963; Atchison 1967). Thus, increased zebra mussel abundance could have positively facilitated faster adult Yellow Bass growth through increased food availability, as has been demonstrated in Yellow Perch (Thayer et al. 1997) and in several species in the Hudson River (Strayer et al. 2004).

The multiple species and life stages represented in the size-specific analyses provided the opportunity to test whether the species and life stages responded similarly to the unique set of growth conditions presented each year. Perhaps not surprisingly, growth in the first year and as fish were reaching maturation was not correlated in any of the four species. Ontogenetic changes in diet, preferred habitat and numerous other characteristics occur in all fish species (Mittelbach and Osenberg 1993; Jobling 1995), and as conditions changed in Clear Lake from year to year the growth of the two life stages studied in each species responded asynchronously. Similarly, there were no significant correlations of adult growth among the four species studied, again suggesting that as conditions changed over time the adult growth of the four species responded uniquely. There were some positive relationships in first year growth among species, and this is consistent with the general observation that diets are similar among fish species when they are young and diverge as they become older (Osenberg et al. 1988). As researchers seek to predict growth responses and managers seek to improve growth responses of fish populations to future environmental changes and management actions, it would be prudent to remember that responses will likely differ among species. Further complicating predictions of impacts of invasive species is the growing evidence that responses of both invaders and resident species may be population-specific rather than species-specific due to genetic and phenotypic variation among populations (Reichard et al. 2015).

In addition to previously mentioned cautions regarding small sample sizes and stocking of walleyes, three reasons for caution when interpreting our results and their implications deserve mention. First, sex of collected fish was not determined in either the recent or the historical studies (see Table S1 for one minor exception), which means that growth differences associated with sex, if any, were not accounted for and thus increased the

variation in our analyses. The consequence of this is that differences evident in our analyses were probably conservative estimates, although it is conceivable that unknown gender bias in fish collections could have occurred and influenced apparent differences in unknown ways. Second, our use of literature derived age of maturity for establishing common lengths within species for the purpose of estimating size-specific growth corresponding to the approximate onset of “adulthood” could have resulted in so-called “adult” growth that was in fact occurring during the transition to true adulthood or even prior to adulthood in some individual fish. Although it is important to be aware of this potential bias, it is not critical to interpretation of results of this study. The more important consideration is that the size-specific approach we used ensured that growth rates occurring at consistent lengths were compared within each species, which in turn implies consistency of the comparison with respect to ecological and physiological characteristics such as preferred prey, metabolic rate, habitat requirements, risk of predation, as well as probable state of sexual maturation. Third, fish sampling methods and ageing structures used differed between the historical and recent time periods. With up to a six-decade time span between the time periods from which we obtained data, advances in technology and changes in practices were inevitable. The methods used during each period were the best available at the time, and although their existence allows important temporal comparison, caution due to potential biases resulting from different methods is warranted.

Clear Lake has undergone significant changes in the past and is poised to change rapidly in the future. A 4-year food web modeling study highlighted large changes in biomass and trophic pathways during 2007-2010 associated with the roles of non-native Common Carp, invading zebra mussels, and phytoplankton dynamics, all in the context of multiple management actions (Colvin et al. 2015). The most recent (2005-2007) of the annual growth

Accepted Article

rates measured in the present study occurred during the first few years of the zebra mussel invasion in Clear Lake, which may have been sufficient to allow glimpses of the eventual effects zebra mussels will have, but was not sufficient to permit a rigorous analysis of zebra mussel impacts. The large changes in growth documented over several decades and more recently suggest that as the Clear Lake ecosystem continues to change, growth rates of its important fish species are also likely to continue changing.

[A]Acknowledgments

Thanks to Jon Christensen, Jason Euchner, Dan Fjeld, David Knoll, Jake Norlin, and Ed Thelen for assistance with field data collection, Jake Norlin for assistance with lab work, Kim Bogenschutz, Joe Larscheid, Mike McGhee, and Jim Wahl for agency support, and Jonathan Spurgeon and two anonymous reviewers for constructive criticism that improved the manuscript. Funding was provided by the Iowa Department of Natural Resources. This study was performed under the auspices of Iowa State University protocol #6-07-6387-1. Reference to trade names does not imply endorsement by the U.S. Government.

[A]References

- Anthony, J. L., and J. A. Downing. 2003. Physical impacts of wind and boat traffic on Clear Lake, Iowa. *Lake and Reservoir Management* 19: 1-14.
- Atchison, G. J. 1967. Contributions to the life history of the yellow bass, *Roccus mississippiensis* (Jordan and Eigenmann), in Clear Lake, Iowa. Master's thesis. Iowa State University, Ames. <http://lib.dr.iastate.edu/rtd/16858>.
- Bailey, R., and H. Harrison. 1945. The fishes of Clear Lake, Iowa. *Iowa State College Journal of Science* 20: 57-77.
- Bajer, P. G., and P. W. Sorensen. 2009. Recruitment and abundance of an invasive fish, the common carp, is driven by its propensity to invade and reproduce in basins that experience wintertime hypoxia in interconnected lakes. *Biological Invasions* 12: 1101-1112.
- Blumenshine, S. C., Y. Vadeboncoeur, D. M. Lodge, K. L. Cottingham, and S. E. Knight. 1997. Benthic-pelagic links: responses of benthos to water-column nutrient enrichment. *Journal of the North American Benthological Society* 16(3): 466-479.
- Bremigan, M. T., J. M. Dettmers, and A. L. Mahan. 2003. Zooplankton selectivity by larval yellow perch in Green Bay, Lake Michigan. *Journal of Great Lakes Research* 29: 501-510.
- Brooks, R. C., R. C. Heidinger, R. J. H. Hoxmeier, and D. H. Wahl. 2002. Relative survival of three sizes of walleyes stocked into Illinois lakes. *North American Journal of Fisheries Management* 22: 995-1006.
- Bulkley, R. V., V. L. Spykermann, and L. E. Inmon. 1976. Food of the pelagic young of walleyes and five cohabiting fish species in Clear Lake, Iowa. *Transactions of the American Fisheries Society* 105: 77-83.

- Accepted Article
- Carlander, K. D. 1969. Handbook of freshwater fishery biology: volume one. Iowa State University Press, Ames.
- Carlander, K. D. 1997. Handbook of freshwater fishery biology: volume three. Iowa State University Press, Ames.
- Carlander, K. D., and P. M. Payne. 1977. Year-class abundance, population, and production of walleye (*Stizostedion vitreum vitreum*) in Clear Lake, Iowa, 1948-74, with varied fry stocking rates. *Journal of the Fisheries Research Board of Canada* 34: 1792-179.
- Carlson, R. E. 1977. A trophic state index for lakes. *Limnology and Oceanography* 22: 361-369.
- Center for Agricultural and Rural Development. 2008. Iowa lakes valuation project. Iowa State University. <http://www.card.iastate.edu/lakes>.
- Chambers, R. C., and E. A. Trippel, editors. 1997. Early life history and recruitment in fish populations. Chapman and Hall, London, UK.
- Clear Lake Enhancement and Restoration Project. 2016. <http://www.clearproject.net/>.
- Cleveland, W. S. 1979. Robust locally weighted regression and smoothing scatterplots. *Journal of the American Statistical Association* 74: 829-836.
- Colvin, M. E., E. D. Katzenmeyer, T. W. Stewart, and C. L. Pierce. 2010. The Clear Lake ecosystem simulation model (CLESM) project. Report to the Iowa Department of Natural Resources. Iowa State University, Ames. http://mec685.cfr.msstate.edu/pdfs/2010_annualReport_FY2009_FV.pdf.
- Colvin, M. E., C. L. Pierce, T. W. Stewart, and S. E. Grummer. 2012. Strategies to control a common carp population by pulsed commercial harvest. *North American Journal of Fisheries Management* 32: 1251-1264.

- Colvin, M. E., C. L. Pierce, and T. W. Stewart. 2015. A food web modeling analysis of a Midwestern, USA eutrophic lake dominated by non-native common carp and zebra mussels. *Ecological Modeling* 312: 26-40.
- Copp, G. H., M. G. Fox, M. Przybylski, F. N. Godinho, and A. Villa-Gispert. 2004. Life-time growth patterns of pumpkinseed *Lepomis gibbosus* introduced to Europe, relative to North American populations. *Folia Zoologica Praha* 53(3): 237-254.
- Crivelli, A. J. 1983. The destruction of aquatic vegetation by carp. *Hydrobiologia* 106: 37-41.
- Davies, I. J. 1980. Relationships between dipteran emergence and phytoplankton production in the Experimental Lakes Area, Northwestern Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* 37: 523-533.
- Dermott, R. M., J. Kalff, W. C. Leggett, and J. Spence. 1977. Production of *Chironomus*, *Procladius* and *Chaoborus* at different levels of phytoplankton biomass in Lake Memphremagog, Quebec-Vermont. *Journal of the Fisheries Research Board of Canada* 34: 2001-2007.
- DeVries, D. R., M. T. Bremigan, and R. A. Stein. 1998. Prey selection by larval fishes as influenced by available zooplankton and gape limitation. *Transactions of the American Fisheries Society* 127: 1040–1050.
- Downing, J. A., J. Kopaska, and D. Bonneau, editors. 2001. Clear Lake Diagnostic and Feasibility Study. Iowa Department of Natural Resources, Des Moines, IA.
<http://limnology.eeob.iastate.edu/reportsandstudies/ClearLake/ClearLakeDiagnosticAndFeasibilityStudy.htm>.
- Egertson, C. J., J. A. Kopaska, and J. A. Downing. 2004. A century of change in macrophyte abundance and composition in response to agricultural eutrophication. *Hydrobiologia* 524: 145-156.

- English, T. S. 1951. Growth studies of the carp *Cyprinus carpio* Linneaus in Clear Lake, Iowa. Master's thesis. Iowa State University, Ames.
- Forney, J. L. 1955. Life history of the black bullhead, (*Ameiurus melas* Rafinesque), of Clear Lake, Iowa. Iowa State College Journal of Science 30: 145–162.
- Gallucci, V.F., and T. J. Quinn II. 1979. Reparameterizing, fitting and testing a simple growth model. Transactions of the American Fisheries Society 108: 14-25.
- Gutreuter, S. 1987. Considerations for estimation and interpretation of annual growth rates. Pages 115-126 in R. C. Summerfelt and G. E. Hall, editors. Age and Growth of Fish. Iowa State University Press, Ames.
- Hanchin, P. A., D. W. Willis, and M. J. Hubers. 2002. Black bullhead growth in South Dakota waters: limnological and community influences. Journal of Freshwater Ecology 17: 65-73.
- Harlan, J. R, and E. B. Speaker. 1987. Iowa Fish and Fishing. Iowa Department of Natural Resources, Des Moines.
- Hokanson, K. E. F. 1977. Temperature requirements of some percids and adaptations to the seasonal temperature cycle. Journal of the Fisheries Research Board of Canada 34: 1524-1550.
- Holland-Bartels, L. E., S. K. Littlejohn, and M. L. Huston. 1990. A guide to larval fishes of the upper Mississippi River. U. S. Fish and Wildlife Service, La Crosse, WI.
- Houser, A., and C. Collins. 1962. Growth of black bullhead catfish in Oklahoma. Oklahoma Fisheries Research Lab Report No. 79, Norman.
- Hutchings, J. A., and M. E. B. Jones. 1998. Life-history variation and growth rate thresholds for maturity in Atlantic salmon, *Salmo salar*. Canadian Journal of Fisheries and Aquatic Sciences 55(suppl. 1) 22-47.

Iowa Department of Natural Resources. 2008. Lake restoration 2008 report and 2009 plan.

Iowa Legislative Services Agency. Des Moines.

Iowa Department of Natural Resources. 2009. Iowa lakes information system. Iowa State University, Ames. <http://limnology.eeob.iastate.edu/lakereport>.

Jackson, Z. J., M. C. Quist and J. G. Larscheid. 2008. Growth standards for nine North American fish species. *Fisheries Management and Ecology* 15: 107-118.

Jacobsen, T. V. 1968. Air-water temperature relationship at Clear Lake, Iowa. Master's thesis. Iowa State University, Ames.

Jobling, M. 1995. *Environmental Biology of Fishes*. Chapman and Hall, London, UK.

Kalff, J. 2002. *Limnology*. Prentice-Hall, Englewood Cliffs, NJ.

Katzenmeyer, E. D. 2010. Fish growth responses to a changing environment: effects of aquatic nuisance species and environmental conditions in a shallow, eutrophic lake. Master's thesis. Iowa State University, Ames.

Kelso, J. R. M. 1972. Conversion, maintenance, and assimilation for walleye *Stizostedion vitreum vitreum*, as affected by size, diet, and temperature. *Journal of the Fisheries Research Board of Canada* 29: 1181-1192.

Koch, J. D., and M. C. Quist. 2007. A technique for preparing fin rays and spines for age and growth analysis. *North American Journal of Fisheries Management* 27: 782-784.

Koehn, J. D. 2004. Carp (*Cyprinus carpio*) as a powerful invader in Australian waters. *Freshwater Biology* 49: 882-894.

Larkin, P. A., J. G. Terpenning, and R. R. Parker. 1956. Size as a determinant of growth rate in rainbow trout *Salmo gairdneri*. *Transactions of the American Fisheries Society* 86: 84-96.

- Larscheid, J. G. 2005. Population densities, biomass and age-growth of common carp and black bullheads in Clear Lake and Ventura Marsh. Iowa Department of Natural Resources, Des Moines.
- McCann, J. A. 1960. Estimates of the fish populations of Clear Lake, Iowa. Master's thesis, Iowa State University, Ames.
- McCauley, E., and J. Kalff. 1981. Empirical relationships between phytoplankton and zooplankton biomass in lakes. *Canadian Journal of Fisheries and Aquatic Sciences* 38: 458-463.
- Miller, T. J., L. B. Crowder, J. A. Rice, and E. A. Marschall. 1988. Larval size and recruitment mechanisms in fishes: toward a conceptual framework. *Canadian Journal of Fisheries and Aquatic Sciences* 45: 1657-1670.
- Millner, R. S., and C. L. Whiting. 1996. Long-term changes in growth and population abundance of sole in the North Sea from 1940 to the present. *ICES Journal of Marine Science* 53:1185-1195.
- Mittelbach, G. G., and C. W. Osenberg. 1993. Stage-structures interactions in bluegill: consequences of adult resource variation. *Ecology* 74: 2381-2394.
- Momot, W. T., J. Erickson, and R. Stevenson. 1977. Maintenance of a walleye (*Stizostedion vitreum vitreum*) fishery in a eutrophic reservoir. *Journal of the Fisheries Research Board of Canada* 34: 1725-1733.
- National Oceanic and Atmospheric Administration. 2009. NOAA satellite and information service. National Climate Data Center. <http://www.ncdc.noaa.gov/oa/ncdc.html>.
- Olson, M. H., T. E. Brooking, D. M. Green, A. J. VanDeValk, and L. G. Rudstam. 2000. Survival and growth of intensively reared large walleye fingerlings and extensively reared small fingerlings stocked concurrently in small lakes. *North American Journal of Fisheries Management* 20: 337-348.

- Osenberg, C. W., E. E. Werner, G. G. Mittelbach, and D. J. Hall. 1988. Growth patterns in bluegill (*Lepomis macrochirus*) and pumpkinseed (*L. gibbosus*) sunfish: environmental variation and the importance of ontogenetic niche shifts. *Canadian Journal of Fisheries and Aquatic Sciences* 45: 17-26.
- Panek, F. M. 1987. Biology and Ecology of Carp. Pages 1-15 in E. L. Cooper. editor. *Carp in North America*. American Fisheries Society, Bethesda, MD.
- Parsons, J. W. 1950. The life history of the yellow perch *Perca flavescens* (Mitchell) of Clear Lake, Iowa. Master's thesis. Iowa State University, Ames.
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters and mean environmental temperature in 175 fish stocks. *Journal du Conseil International pour l'Exploration de la Mer* 39: 175-192.
- Penne, C. R., and C. L. Pierce. 2008. Seasonal distribution, aggregation, and habitat selection of Common Carp in Clear Lake, Iowa. *Transactions of the American Fisheries Society* 137: 1050-1062.
- Pierce, C. L., C. S. Guy, M. A. Pegg, and P. J. Braaten. 2003. Fish growth, mortality, recruitment, condition and size structure. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone rivers. Volume 4. U.S. Geological Survey, Cooperative Research Units, Ames, IA.
- Pratt, T. C., and M. G. Fox. 2003. Contribution and relative survival of age-0 walleyes stocked at two sizes in six eastern Ontario lakes. *North American Journal of Fisheries Management* 23: 997-1005.
- Putman, J. H., C. L. Pierce, and D. M. Day. 1995. Relationships between environmental variables and size-specific growth rates of Illinois stream fishes. *Transactions of the American Fisheries Society* 124: 252-261.

Quist, M. C., C. S. Guy, R. D. Schultz, and J. L. Stephen. 2003. Latitudinal comparison of walleye growth in North America and factors influencing growth of walleyes in Kansas reservoirs. *North American Journal of Fisheries Management* 23: 677-692.

Quist, M. C., M. A. Pegg, and D. R. DeVries. 2012. Age and growth. Pages 677-732 in A. V. Zale, D. L. Parrish, and T. M. Sutton, editors. *Fisheries techniques*, 3rd edition. American Fisheries Society, Bethesda, MD.

R Development Core Team. 2009. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.

Reichard, M., K. Douda, M. Przybylski, O. P. Popa, E. Karbanova, K. Matasova, K. Rylkova, M. Polacik, R. Blazek, and C. Smith. 2015. Population-specific responses to an invasive species. *Proceedings of the Royal Society of London, B* 282: 20151063.

Ridenhour, R. L. 1960. Abundance, growth and food of young game fish in Clear Lake, Iowa, 1949 to 1957. *Iowa State Journal of Science* 35: 1-23.

Rose, E. T., and T. Moen. 1950. Results of increased fish harvest in Lost Island Lake. *Transactions of the American Fisheries Society* 80: 50-55.

Santucci, V.J., Jr., and D. H. Wahl. 1993. Factors influencing survival and growth of stocked walleye in a centrarchid-dominated impoundment. *Canadian Journal of Fisheries and Aquatic Sciences* 50: 1548-1558.

Secor, D. H., J. M. Dean, and E. H. Laban. 1991. Manual for otolith removal and preparation for microchemical examination. Technical Publication No. 1991-01. Electric Power Research Institute, and the Belle W. Baruch Institute for Marine Biology and Coastal Research, Palo Alto, CA.

Shoup, D. E., S. P. Callahan, D. H. Wahl, and C. L. Pierce. 2007. Size-specific growth of bluegill, largemouth bass and channel catfish in relation to prey availability and limnological variables. *Journal of Fish Biology* 70: 21-34.

Slipke, J. W., and M. Maceina. 2014. Fishery Analysis and Modeling Simulator (FAMS).
American Fisheries Society, Bethesda, MD.

Staggs, M. S., and K. J. Otis, K. 1996. Factors affecting first-year growth of fishes in Lake
Winnebago, Wisconsin. *North American Journal of Fisheries Management* 16: 608-618.

Stearns, S. C. 2000. Life history evolution: successes, limitations, and prospects.
Naturwissenschaften 87:476-486.

Stewart, T. W., J. G. Miner, and R. L. Lowe. 1998. Quantifying mechanisms for zebra mussel
effects on benthic macroinvertebrates: organic matter production and shell-generated
habitat. *Journal of the North American Benthological Society* 17: 81-94.

Strayer, D. L., K. A. Hattala, and A. W. Kahnle. 2004. Effects of an invasive bivalve
(*Dreissena polymorpha*) on fish in the Hudson River estuary. *Canadian Journal of
Fisheries and Aquatic Sciences* 61: 924-941.

Stuart, I. G., and M. Jones. 2006. Large, regulated forest floodplain is an ideal recruitment
zone for non-native common carp (*Cyprinus carpio* L.). *Marine and Freshwater
Research* 57: 333-347.

Summerfelt, R. C., and G. E. Hall, editors. 1987. Age and growth of fish. Iowa State
University Press, Ames.

Thayer, S. A., R. C. Haas, R. D. Hunter, and R. H. Kushler. 1997. Zebra mussel (*Dreissena
polymorpha*) effects on sediment, other zoobenthos, and the diet and growth of adult
yellow perch in pond enclosures. *Canadian Journal of Fisheries and Aquatic Sciences*
54(8): 1903-1915.

Van Den Avyle, M. J., and R. S. Hayward. 1999. Dynamics of exploited fish populations.
Pages 127-166 in C. C. Kohler & W. A. Hubert, editors. *Inland fisheries management in
North America*, 2nd edition. American Fisheries Society, Bethesda, MD.

Accepted Article

Wahl, J. 2001. An analysis of the fishery of Clear Lake, Iowa. In Downing, J. A., J. Kopaska, and D. Bonneau, editors. Clear Lake Diagnostic and Feasibility Study. Iowa Department of Natural Resources, Des Moines.
<http://limnology.eeob.iastate.edu/reports/studies/ClearLake/ClearLakeDiagnosticAndFeasibilityStudy.htm>.

Weatherly, A. H., and H. S. Gill. 1987. The biology of fish growth. Academic Press, London, UK.

Weber, M. J., and M. L. Brown. 2009. Effects of common carp on aquatic ecosystems 80 years after “Carp as a Dominant”: ecological insights for fisheries management. *Reviews in Fisheries Science* 17: 524-537.

Welker, B. D. 1963. Summer food habits of yellow bass and black bullheads in Clear Lake. *Proceedings of the Iowa Academy of Science* 69: 286-295.

Welker, M. T., C. L. Pierce, and D. H. Wahl. 1994. Growth and survival of larval fishes: roles of competition and zooplankton abundance. *Transactions of the American Fisheries Society*, 123, 703–717.

[A]Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article.

Table 1. Environmental variables in Clear Lake, Iowa, USA, 1994 - 2007. Degree days are annual totals; other variables are annual summertime averages. Empty cells indicate no data.

Year	Degree days (°C)	Transparency (m)	Chlorophyll <i>a</i> (µg/L)	Phytoplankton (mg/L)	Zooplankton (µg/L)
1994	2,655				
1995	2,584				
1996	2,372				
1997	2,486				
1998	2,892				
1999	2,602				
2000	2,729	0.4	30.8	37.1	23.4
2001	2,688	0.7	70.7	14.6	124.3
2002	2,569	0.8	30.2	142.6	223.7
2003	2,518	1.4	20.7	66.6	101.9
2004	2,465	0.5	81.5	543.6	276.1
2005	2,836	0.5	82.0	305.3	39.3
2006	2,584	0.5	53.1	3112.7	119.8
2007	2,878	1.2	22.4	40.9	66.5

Table 2. Correlations of annual growth (mm) at age 0 (AG-0) and at length at maturation (AG-m) of Black Bullhead, Common Carp, Walleye and Yellow Bass in Clear Lake, Iowa, USA. In each cell the top value is the Pearson correlation coefficient, the middle value is the *p*-value, and the bottom value is the sample size. Dashes indicate sample size < 3.

	Black Bullhead	Common Carp		Walleye		Yellow Bass	
	AG-m	AG-0	AG-m	AG-0	AG-m	AG-0	AG-m
Black Bullhead		0.88	-0.16	0.97	-0.03	0.97	-0.05
AG-0	-	0.122	0.843	0.027	0.962	0.030	0.948
		4	4	4	5	4	4
Black Bullhead				-0.55	0.49		0.01
AG-m		-	-	0.626	0.671	-	0.991
				3	3		3
Common Carp			0.55	-0.31	0.27	0.93	0.55
AG-0			0.101	0.688	0.652	0.070	0.633
			10	4	5	4	3
Common Carp				-0.82	0.29	0.81	0.93
AG-m				0.087	0.584	0.100	0.071
				5	6	5	4
Walleye					-0.19	0.07	-0.24
AG-0					0.723	0.931	0.657
					6	4	6
Walleye						0.48	-0.32
AG-m						0.331	0.533
						6	6
Yellow Bass							0.53
AG-0							0.466
							4

Table 3. Correlations of annual growth (mm) at age 0 (AG-0) and at length at maturation (AG-m) of Black Bullhead, Common Carp, Walleye and Yellow Bass with environmental variables in Clear Lake, Iowa, USA. In each cell the top value is the Pearson correlation coefficient, the middle value is the *p*-value, and the bottom value is the sample size.

	Black Bullhead		Common Carp		Walleye		Yellow Bass	
	AG-0	AG-m	AG-0	AG-m	AG-0	AG-m	AG-0	AG-m
Degree days (°C)	0.44	-0.95	0.29	0.10	-0.24	-0.11	-0.11	0.72
	0.463	0.202	0.380	0.759	0.598	0.791	0.841	0.110
	5	3	11	12	7	8	6	6
Transparency (m)	-0.50	-0.33	-0.16	0.08	-0.41	-0.69	-0.18	0.44
	0.394	0.784	0.798	0.877	0.419	0.057	0.738	0.383
	5	3	5	6	6	8	6	6
Chlorophyll <i>a</i> (µg/L)	0.71	-0.17	0.40	-0.18	0.79	0.19	0.39	-0.54
	0.179	0.894	0.505	0.728	0.061	0.651	0.449	0.2733
	5	3	5	6	6	8	6	6
Phytoplankton (mg/L)	0.66	0.97	0.94	0.89	0.03	0.45	0.75	0.12
	0.222	0.167	0.018	0.018	0.953	0.266	0.085	0.826
	5	3	5	6	6	8	6	6
Zooplankton (µg/L)	-0.34	0.99	-0.39	-0.22	-0.24	0.10	0.20	-0.71
	0.573	0.099	0.513	0.671	0.647	0.818	0.706	0.116
	5	3	5	6	6	8	6	6

Table 4. Growth parameter estimates and standard errors (S.E.) for Black Bullhead, Common Carp, Walleye and Yellow Bass from studies conducted in different time periods in Clear Lake, Iowa, USA. N is the sample size of fish examined in the recent time period and n is the number of studies used to calculate each parameter and N . L_{∞} , k , t_0 and ω are parameters from the von Bertalanffy growth model. Significant differences among ω values within species are indicated by superscripts; values with the same superscripts are not significantly different.

Species	Time period	N	n	L_{∞}		k		t_0		ω	
				Estimate	S.E.	Estimate	S.E.	Estimate	S.E.	Estimate	S.E.
Black Bullhead	1955		1	343.7	26.61	0.28	0.07	-0.22	0.33	96.9 ¹	17.95
	1999		1	331.2	16.47	0.26	0.02	-0.12	0.05	84.9 ¹	3.71
	Recent	61	1	313.3	11.00	0.43	0.04	-0.02	0.05	136.0 ²	7.16
Common Carp	1952		1	844.1	13.02	0.49	0.03	0.39	0.04	416.8 ¹	17.16
	1999		1	758.2	21.97	0.31	0.04	0.25	0.09	237.5 ²	24.04
	Recent	40	1	728.4	3.21	0.36	0.01	0.48	0.02	258.9 ²	2.92
Walleye	1940s		2	779.0	13.50	0.19	0.01	-0.49	0.10	144.7 ¹	5.43
	1948-1974		1	694.4	7.40	0.20	0.01	-0.46	0.08	140.8 ¹	3.99
	Recent	22	1	688.7	16.14	0.26	0.01	0.05	0.02	178.8 ²	4.08
Yellow Bass	1940s		9	262.8	6.10	0.54	0.05	0.28	0.08	140.8 ¹	10.51
	1960s		9	245.3	10.95	0.38	0.06	0.01	0.04	94.0 ²	9.71
	Recent	10	1	229.2	2.91	0.54	0.02	-0.01	0.02	123.1 ¹	2.97

Figure Captions

Figure 1. Annual growth (mm) at age 0 (AG-0) and at length at maturation (AG-m) (\pm 1 S.E.) for Black Bullhead, Common Carp, Walleye and Yellow Bass in Clear Lake, Iowa, USA, 1994-2007. AG-0 values are mean back-calculated lengths at age 1; AG-m values were estimated from LOWESS regressions (Supporting Information Figures S1-S5).

Figure 2. Mean back-calculated length at age (mm) (\pm 95% C. I.) for Black Bullhead, Common Carp, Walleye and Yellow Bass from studies conducted in different time periods in Clear Lake, Iowa, USA.

Figure 3. Relative growth index (RGI) (%) for Common Carp and Walleye from studies conducted in different time periods in Clear Lake, Iowa, USA. Dotted horizontal lines represent the “average” length at age as predicted by the standardized growth model for each species.





