

12-2000

Osajin and Pomiferin, Two Isoflavones Purified from Osage Orange Fruits, Tested for Repellency to the Maize Weevil (Coleoptera: Curculionidae)

Chris Peterson
Iowa State University

Anne Fristad
Iowa State University

Rong Tsao
Iowa State University

Joel R. Coats
Iowa State University, jcoats@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/ent_pubs

 Part of the [Entomology Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/ent_pubs/328. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Entomology at Iowa State University Digital Repository. It has been accepted for inclusion in Entomology Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Osajin and Pomiferin, Two Isoflavones Purified from Osage Orange Fruits, Tested for Repellency to the Maize Weevil (Coleoptera: Curculionidae)

Abstract

The fruit of the osage orange tree, *Maclura pomifera* (Raf.) Schneid (Moraceae), has long been thought to be repellent to insects. A preliminary study reported here confirmed repellency of fruit extracts to the maize weevil, *Sitophilus zeamais* Motschulsky. Two isoflavones, osajin and pomiferin, were isolated from the mature fruit of *M. pomifera* in high purity ($\geq 95\%$). Testing of purified osajin and pomiferin failed to show repellency. Repellency is likely caused by factors other than isoflavones in the fruit.

Keywords

Maclura pomifera, *Sitophilus zeamais*, osajin, pomiferin, isoflavone, repellency

Disciplines

Entomology

Comments

This article is from *Environmental Entomology* 29 (2000): 1133, doi:[10.1603/0046-225X-29.6.1133](https://doi.org/10.1603/0046-225X-29.6.1133). Posted with permission.

Rights

This article is the copyright property of the Entomological Society of America and may not be used for any commercial or other private purpose without specific permission of the Entomological Society of America.

Osajin and Pomiferin, Two Isoflavones Purified from Osage Orange Fruits, Tested for Repellency to the Maize Weevil (Coleoptera: Curculionidae)

CHRIS PETERSON, ANNE FRISTAD,¹ RONG TSAO,² AND JOEL R. COATS³

Pesticide Toxicology Laboratory, 116 Insectary, Department of Entomology, Iowa State University, Ames, IA 50011

Environ. Entomol. 29(6): 1133–1137 (2000)

ABSTRACT The fruit of the osage orange tree, *Maclura pomifera* (Raf.) Schneid (Moraceae), has long been thought to be repellent to insects. A preliminary study reported here confirmed repellency of fruit extracts to the maize weevil, *Sitophilus zeamais* Motschulsky. Two isoflavones, osajin and pomiferin, were isolated from the mature fruit of *M. pomifera* in high purity ($\geq 95\%$). Testing of purified osajin and pomiferin failed to show repellency. Repellency is likely caused by factors other than isoflavones in the fruit.

KEY WORDS *Maclura pomifera*, *Sitophilus zeamais*, osajin, pomiferin, isoflavone, repellency

THE FRUIT OF the osage orange, *Maclura pomifera* (Raf.) Schneid (Moraceae), has been used as an insect repellent for many years. Pioneers placed the ripe fruit of this tree in cupboards to repel roaches and other insects (Sand 1991, Brandies 1979). Past research indicates that there may be scientific justification for this well-popularized use. Karr and Coats (1991) found fragments of the fruit, as well as its hexane and methanol extracts, to be significantly repellent to the German cockroach *Blattella germanica*. The wood of the osage orange tree resisted termites (Esenther 1977; Wolcott 1953, 1955, 1957) and had antifungal properties (Smith and Perino 1981). Antimicrobial activity has been found in extracts containing osajin or pomiferin (Mahmoud 1981), two of the major angular isoflavonoid components present in osage orange fruits (Fig. 1A) (Walter et al. 1938, Wolfrom and Mahan 1942). Osage orange fruit was also found to contain small amounts of the linear isoflavonoids scandenone and auricularin, structural isomers of osajin and pomiferin, respectively (Fig. 1B) (Delle Monache et al. 1994).

In this study we report repellency from the extracts of *M. pomifera* fruit. Because osajin and pomiferin are well known to occur in this fruit, the extraction and purification of osajin and pomiferin is reported. The results of an assay that examined the behavioral response of the maize weevil *Sitophilus zeamais* Motschulsky to highly purified ($\geq 95\%$) osajin and pomiferin are reported as well.

Materials and Methods

Extraction of Fruits for Preliminary Assay. Mature *M. pomifera* fruit was collected in Auburn, NE, in October 1995, and stored at -20°C for 8 mo. For both extraction types reported here, the fruits were divided into "rind" and "core" portions. The rind was defined as the portion of the fruit from the outer surface inward to the outer perimeter of the seed layer. The core was defined as the seed layer and extending inward. After thawing to room temperature, the fruit was cut into 250-g rind or 250-g core pieces and placed into separate soxhlet extraction apparatuses. For each fruit portion, rind or core, one soxhlet apparatus contained 500 ml hexane, another contained 500 ml dichloromethane, and a third contained 500 ml methanol. Each apparatus was heated to the boiling point of its respective solvent and allowed to cycle for 24 h. The solvent was collected, filtered to remove particulates by using Whatman (Hillsboro, OR) G8 glass fiber filters, and passed through anhydrous sodium sulfate to remove water. The solvents were removed by rotary evaporation at 500 mmHg vacuum and 30°C . Alternatively, 250 g of rind and core pieces were placed in 500 ml of either hexane, dichloromethane, or methanol and allowed to soak for 24 h. The particulates were removed by vacuum filtration, the water was removed with anhydrous sodium sulfate and the solvents were removed by rotary evaporation. The residues were redissolved in a minimum amount of solvent and used in the bioassays described later in this section. The extracts obtained from each method were tested in a preliminary round of testing (data not shown). Those extracts obtained by soaking extraction of the cores and rinds with hexane, by soxhlet extraction of the rind and core with dichloromethane, and by soxhlet extraction of the rind with methanol were retested

¹ Current address: Luther College, Decorah, IA, 52101.

² Current address: Agriculture Canada, Vineland, ON, Canada.

³ To whom correspondence should be addressed.




Fig. 1. Structures of (A) osajin (R = H) and pomiferin (R = OH) and (B) scandenone (R = H) and auricularin (R = OH).

because they showed potential activity in the initial testing. The results of the second round of testing are those reported here. Repellency was measured at 0.5, 18, and 24 h.

Extraction of Osajin and Pomiferin. After thawing to room temperature, 12 fruits were cut into 2-cm³ pieces and pressed by hand to obtain the juice, which contained high amounts of the components of interest. The collected juice was centrifuged for 30 min. Thin-layer chromatography detected no osajin or pomiferin in the water layer. The water layer was discarded, and the particulates were extracted three times with 200 ml dichloromethane (ACS certified grade, Fisher, Pittsburgh, PA). The dichloromethane solution was passed through anhydrous sodium sulfate, collected, and vacuum filtered. The pressed fruit pieces were extracted also with three 500-ml portions of dichloromethane, which were passed through anhydrous sodium sulfate to remove water and then vacuum filtered. The collected dichloromethane was combined with the dichloromethane extract of the juice. The dichloromethane was removed from the combined extracts via rotary evaporation. The residue was dissolved in acetone and passed through a large (5.5 by 40 cm) open silica gel column wet-packed with acetone. Acetone (100%, ACS certified grade, Fisher, Pittsburgh, PA) was used as the eluent, and washing continued until products were no longer observed in the eluate, as determined by spotting the eluate on thin-layer chromatography (TLC) plates, using 1:1 hexane:ethyl acetate as a solvent system.

The acetone was removed by rotary evaporation. The resultant mixture, containing osajin and pomiferin, as well as other components, was run through an open silica gel column with 9:1 hexane:ethyl acetate as an eluent. Fractions were collected in 25-ml portions until the fractions containing osajin began to contain pomiferin as well, evidenced by TLC. The solvent system was then changed to 4:1 hexane:ethyl acetate until the fractions began to contain pomiferin only. At that time, the solvent system was changed to 7:3 hexane:ethyl acetate until pomiferin was no longer detected in the fractions. Fractions containing a mixture

of osajin and pomiferin were purified either on an identical column, or by preparatory thin-layer chromatography in 1:1 hexane:ethyl acetate. The purity of the resulting osajin and pomiferin samples was determined by high-pressure liquid chromatography (HPLC). The identities of osajin and pomiferin were confirmed by using solid-probe electron-impact mass spectrometry (EIMS) and proton nuclear magnetic resonance spectroscopy (¹H-NMR).

Analysis and Identification. The melting range of each isolated compound was determined by using a Fisher-Johns Melting Point Apparatus (Pittsburgh, PA). Infrared spectra (IR) of these compounds were obtained by using a Beckman Acculab 2 (Fullerton, CA). High-pressure liquid chromatography was conducted by using a Hewlett-Packard Series 1100 HPLC (Palo Alto, CA) and a Waters μ Bondupak C₁₈ column (Milford, MA) with a flow rate of 2 ml/min of 1:1 acetonitrile:water initially and ramped to 100% acetonitrile over 10 min. Peaks were detected at 254 nm by using a Kratos Spectraflow 757 variable-wavelength detector (Chestnut Ridge, NY). Proton nuclear magnetic resonance spectra (¹H-NMR) in deuterioacetone and carbon nuclear magnetic resonance spectra (¹³C-NMR) in deuteriochloroform (tetramethylsilane internal standard) were obtained on a Varian VXR-300 NMR (Palo Alto, CA). A Finnigan TSQ700 mass spectrometer (San Jose, CA) was used to obtain the mass spectra, with solid probe introduction of the sample, and electron impact of 70 eV.

Behavioral Bioassay of Osajin and Pomiferin. The repellency assay is based on that used for German cockroaches by Karr and Coats (1991). Maize weevil (*S. zeamais*) adults from a colony reared in the laboratory for over five years were used. The repellency tests were conducted in a choice apparatus, which consisted of two 35-ml jars connected at their mouths by transparent plastic tubing (7 cm by 1.5 cm i.d.). For the preliminary test, the residue from the extraction process was dissolved in a minimum amount of solvent, and 1 ml was applied to the filter paper. Appropriate solvent controls were conducted. Highly pure ($\geq 95\%$) osajin or pomiferin was diluted in acetone to

Table 1. Repellency of five osage orange extracts to the maize weevil

Extraction method	% repelled from treated chamber (\pm SEM)		
	0.5 h	18 h	24 h
Soak: Hexane core	56.0 \pm 6.0	41.0 \pm 3.0	28.0 \pm 3.0
Soak: Hexane rind	62.5 \pm 12.5	59.5 \pm 9.5	41.0 \pm 3.0
Soxhlet: Dichloromethane core	40.5 \pm 21.5	34.5 \pm 3.5	72.0 \pm 3.0
Soxhlet: Dichloromethane rind	90.5 \pm 9.5	97.0 \pm 3.0	84.5 \pm 3.5
Soxhlet: Methanol rind	56.5 \pm 12.5	50.0 \pm 0.0	59.5 \pm 15.5

50 mg/ml (approximately saturated), then serially diluted to 5 and 0.5 mg/ml. One milliliter of one of these solutions (treated side) or 1 ml of acetone (untreated side) was applied to a filter paper, and the acetone allowed to evaporate in a fume hood before the papers were placed in the choice-jar units. The set-up for the control groups was the same but both sides contained papers previously treated with 1 ml acetone. Each assay was conducted with 20 insects, starting with 10 per jar. The test had five replicates. Two series of randomizations were carried out; one for each replicate to determine the position of the choice-jar units in relation to one another, and a second to determine for each treatment unit the position of the treated side, either to the right or to the left. The distribution of the insects, in the treated side, the untreated side, and in the tube connecting the two sides was determined at 1, 17, 24, and 48 h. Any repellency of the test compounds would likely occur on contact because of the compounds' low volatility, therefore the assumption was made that the chemical repelled the insects in the connecting tube. The data were analyzed by using analysis of variance (ANOVA) to detect significance. The four time points were compared with each other by using least significant difference (LSD) (lsd, $\alpha = 0.05$).

Results and Discussion

Extraction of Fruit and Preliminary Assay. Upon removal of the solvent, 250-g fruit pieces yielded 1.14 g in hexane soaking extraction of the core, 0.77 g in hexane soaking extraction of the rind, 1.93 g in dichloromethane soxhlet extraction of the core, 1.84 g in dichloromethane soxhlet extraction of the rind, and 28.0 g in methanol soxhlet extraction of the rind. The methanol extraction resulted in a syrup, because

starch, sugars, and water are likely to be soluble in hot methanol.

Table 1 reports the percentage of weevils repelled from the treated chamber (\pm SEM). ANOVA detected that significance caused by extraction method was not found at 0.5 h, but was found at 18 ($F = 25.06$; $df = 4, 5$; $P < 0.002$) and 24 h ($F = 9.32$; $df = 4, 5$; $P < 0.02$). In cases where significance was observed, the dichloromethane extraction of the rind was the most effective. Because osajin and pomiferin are well known to occur in the extracts of this plant, we decided to focus on these two compounds as the potential active ingredients.

Analysis and Identification. The purity of osajin was found by HPLC to be 94.6%, and the purity of pomiferin was 98.2%.

The melting point range for osajin was observed to be from 186 to 191°C, and past studies report a melting point of 190–192°C (Jain and Sharma 1974). Scandone has a reported melting point range of 164–165°C (Delle Monache et al. 1994). Pomiferin melted at 172–176°C, at which temperature a color change occurred and the substance resolidified. This resolidified substance melted at 204–209°C. The product was recovered from the melting point apparatus with acetone, and the solution was run on TLC with 1:1 hexane:ethyl acetate. Visualization of the plate with potassium permanganate (0.1 M) spray revealed a spot identical in R_f value to the unheated sample ($R_f = 0.55$), and a lower spot that streaked from the origin to $R_f = 0.30$. Pomiferin has a reported melting point of 200.5°C at maximum purity, whereas auricularin melts at 176–178°C (Delle Monache et al. 1994).

Infrared spectroscopy of osajin showed peaks at the following frequencies: 3,610 cm^{-1} , 3,060 cm^{-1} , 1,650 cm^{-1} , 1,620 cm^{-1} , 1,590 cm^{-1} , and 1,520 cm^{-1} , 1,435 cm^{-1} , 1,380 cm^{-1} , 1,275 cm^{-1} , 1,230 cm^{-1} , and 1,180

Table 2. Repellency of osajin and pomiferin to the maize weevil

Compound	Dose (mg/chamber)	% repelled from treated chamber (\pm SEM)			
		1 h	17 h	24 h	48 h
Osajin	50	76 (3.3)a	59 (3.3)a	45 (3.9)a	47 (4.1)a
	5.0	68 (9.8)a	41 (4.8)b	48 (4.1)ab	48 (4.9)ab
	0.5	51 (8.0)a	38 (3.7)b	48 (2.6)ab	47 (4.6)ab
	0 (control)	61 (4.8)a	54 (11.7)a	42 (12.1)a	40 (4.7)a
Pomiferin	50	76 (7.0)a	55 (7.6)a	53 (4.6)a	58 (11.4)a
	5.0	70 (6.1)a	53 (4.6)a	57 (9.6)a	55 (8.5)a
	0.5	75 (4.2)a	62 (8.5)ab	57 (4.6)b	64 (4.0)ab
	0 (control)	61 (4.8)a	54 (11.7)a	42 (12.1)a	40 (4.7)a

Values followed by the same letter within a row are not significantly different according to LSD ($\alpha = 0.05$).

cm⁻¹. The reported structures of osajin (Wolfrom et al. 1946) and scandenone (Pelter and Stainton 1966) both contain bonds that may give rise to these observed peaks.

The mass spectra of osajin and pomiferin agreed with that reported by Delle Monache et al. (1994). Osajin and its isomer scandenone are indistinguishable from one another by mass spectrometry, as are pomiferin and auricularin.

The spectrum of osajin obtained by ¹³C-NMR in deuteriochloroform has peaks corresponding to the carbons of osajin, as displayed in Fig. 1A. These chemical shifts should differ from those of Delle Monache et al. (1994), who report chemical shifts in deuterioacetone. The spectrum displayed the following peaks: δ 181.14 (C at 4), δ 159.26 (C at 5), δ 157.31 (C at 4'), δ 155.98 (C at 7), δ 152.37 (C at 2), δ 150.55 (C at 8a), δ 131.64 (C at 3'''), δ 130.34 (C at 2' and 6'), δ 127.17 (C at 3''), δ 123.57 (C at 3), δ 122.98 (C at 1'), δ 121.90 (C at 2'''), δ 115.70 (C at 3' and 5'), δ 114.94 (C at 4''), δ 112.90 (C at 6), δ 105.58 (C at 4a), δ 100.76 (C at 8), δ 77.09 (C at 2''), δ 28.12 (the two 2'' methyls), δ 25.80 (*trans*-3''' methyl), δ 21.30 (C at 1'''), and δ 17.92 (*cis*-3''' methyl). See Fig. 1A for designation of carbon atoms. Determination of carbon positions was achieved through comparison with Delle Monache et al. (1994) and published ¹³C-NMR spectra from similar compounds (Silverstein et al. 1991).

Delle Monache et al. (1994) reported ¹H-NMR spectrum for osajin and pomiferin in deuterioacetone. Our spectrum of osajin agreed with theirs to within δ 0.03 for all peaks, except for δ 8.50, which they report as δ 8.60. This proton is on a hydroxyl group, the chemical shift of which can vary more so than other protons. Delle Monache et al. (1994) reported diagnostic peaks occurring at δ 8.17 and 3.31 for osajin, and at δ 8.26 and 3.41 for scandenone. Our spectrum agreed with osajin. Our analysis of pomiferin by ¹H-NMR in deuterioacetone agreed with theirs to within δ 0.02, except for δ 8.00, which varied from theirs by δ 0.07. As was the case with osajin, this proton occurred on a hydroxyl group. Delle Monache et al. (1994) reported diagnostic peaks at δ 8.16 and 3.31 for pomiferin, and δ 8.24 and 3.40 for auricularin. Our spectrum was consistent with pomiferin.

Scandenone is present in *M. pomifera* fruit at around 15% of the concentration of osajin, and auricularin is present at 13% of pomiferin's concentration (Delle Monache et al. 1994). Though our MS, IR, and ¹³C-NMR data are not conclusive, the observed melting point, ¹H-NMR, and the relative abundance in the plant reported in Delle Monache et al. (1994) indicate that we have isolated the angular isoflavones osajin and pomiferin by the processes outlined here. Small amounts of scandenone and auricularin are present as minor contaminants, evidenced by minute peaks on ¹H-NMR corresponding to the diagnostic peaks of the linear isomers.

Behavioral Assay of Osajin and Pomiferin. The results of the behavioral assay are summarized in Table 2. The data are expressed as the mean percentage of insects repelled from the treated chamber (±SEM).

ANOVA detected significance caused by time for osajin ($F = 7.73$; $df = 3, 64$; $P < 0.001$) and pomiferin ($F = 4.72$; $df = 3, 64$; $P < 0.005$). Significance caused by concentration was not seen for osajin or pomiferin, and the interaction between concentration and time was not significant for either compound. Although the number of insects repelled in 1 h was significantly more at 50 than at 0.5 mg osajin per chamber, it was not significantly different from the control, because there was no significance caused by concentration in the model. When significance was seen for a concentration (at $\alpha = 0.05$) among the various time intervals, the number of insects repelled decreased with increasing time in the apparatus. Because repellency caused by concentration was not significant, we concluded that factors other than osajin or pomiferin, unknown to us at this time, affected the distribution of the insects within the apparatus.

No significant repellency was observed when the weevils were exposed to purified osajin and pomiferin. We have thus concluded that neither osajin nor pomiferin is singularly responsible for *M. pomifera* repellency to *S. zeamais*. Many of the constituents in this plant are volatile, but were not examined in this study. Volatility is important for repellency over large areas, such as in homes or workplaces. Efforts are currently underway in our laboratory to identify volatile components from this fruit and to test the repellency of these volatile components in behavioral assays.

Acknowledgments

Charles Baker and Kamel Harrata (Iowa State University Instrument Services, Department of Chemistry, Iowa State University, Ames, IA); Chris Schmidt and Mickolyn Funne-mark-Clapper for their assistance; and the James and Nancy Kite family for the oranges. We thank the Program for Women in Science and Engineering at Iowa State University, Ames, IA. This is journal paper J-18001 of the Iowa Agriculture and Home Economics Experiment Station, Iowa State University, Ames, IA, project number 3187.

References Cited

- Brandies, M. 1979. The 'apple' that chases bugs. *Org. Garden*. 26: 138-140.
- Delle Monache, G., R. Scurria, A. Vitali, B. Botta, B. Monacelli, G. Pasqua, C. Palocci, and E. Cernia. 1994. Two isoflavones and a flavone from the fruits of *Maclura pomifera*. *Phytochemistry* 37: 893-898.
- Esenher, G. 1977. Nutritive supplement method to evaluate resistance of natural or preservative-treated wood to subterranean termites. *J. Econ. Entomol.* 70: 341-346.
- Jain, A. C., and B. N. Sharma. 1974. Synthesis of alpinum isoflavone, osajin, and warangalone. *J. Org. Chem.* 39: 2215-2217.
- Karr, L. L., and J. R. Coats. 1991. Repellency of dried bay leaves (*Laurus nobilis*), Wrigley's[®] spearmint gum, raw osage orange fruit (*Maclura pomifera*), and extracts of osage orange fruit to the German cockroach, 1989. *Insectic. Acaric. Tests* 17: 393.
- Mahmoud, Z. 1981. Antimicrobial components from *Maclura pomifera* fruit. *J. Med. Plant Res.* 42: 299-301.

- Pelter, A., and P. Stainton. 1966. The extractives from *Derris scandens*, part II: The isolation of osajin and two new isoflavones, scandenone and scandinone. *J. Chem. Soc. (C)* 7: 701-704.
- Sand, S. 1991. A tree history, the osage orange. *Am. Horticult.* 37-39.
- Silverstein, R. M., G. C. Bassler, and T. C., Morill [eds.]. 1991. Spectrometric identification of organic compounds 5th ed. Wiley, New York.
- Smith, J. L., and J. V. Perino. 1981. Osage orange (*Maclura pomifera*): history and economic uses. *Econ. Bot.* 35: 24-41.
- Walter, E. D., M. L. Wolfrom, and W. M. Hess. 1938. A yellow pigment from the osage orange. *J. Am. Chem. Soc.* 60: 574-577.
- Wolcott, G. N. 1953. Stilbene and comparable materials for dry-wood termite control. *J. Econ. Entomol.* 46: 374-375.
- Wolcott, G. N. 1955. Organic termite repellents tested against *Cryptotermes brevis* Walker. *J. Agric. Univ. P.R.* 39: 115-149.
- Wolcott, G. N. 1957. Inherent natural resistance of woods to the attack of the west Indian dry-wood termite, *Cryptotermes brevis* Walker. *J. Agric. Univ. P.R.* 41: 259-311.
- Wolfrom, M. L., and J. Mahan. 1942. Osage orange pigments, IX: Improved separation; establishment of the isopropylidene group. *J. Am. Chem. Soc.* 64: 308-311.
- Wolfrom, M. L., W. D. Harris, G. F. Johnson, F. E. Mahan, S. M. Moffett, and B. Wildi. 1946. Osage orange pigments, XI: complete structures of osajin and pomiferin. *J. Am. Chem. Soc.* 68: 406-418.

Received for publication 14 February 2000; accepted 15 August 2000.
