

IOWA STATE UNIVERSITY
Digital Repository
Farm Progress Reports

2016 Report
Issue 1 2016 *Farm Progress Reports*

Number RFR-A1606

2017

Iowa Acoustic Bat Surveys

Julie Blanchong
Iowa State University, julieb@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/farmprogressreports>



Part of the [Agriculture Commons](#), and the [Population Biology Commons](#)

Recommended Citation

Blanchong, Julie (2017) "Iowa Acoustic Bat Surveys," *Farm Progress Reports*: Vol. 2016 : Iss. 1 , Article 160.
DOI: <https://doi.org/10.31274/farmprogressreports-180814-1725>
Available at: <https://lib.dr.iastate.edu/farmprogressreports/vol2016/iss1/160>

This Ag Engineering/Agronomy, Central Iowa and BioCentury Research Farms is brought to you for free and open access by the Extension and Experiment Station Publications at Iowa State University Digital Repository. It has been accepted for inclusion in Farm Progress Reports by an authorized editor of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Iowa Acoustic Bat Surveys

RFR-A1606

Julie Blanchong, associate professor
Department of Natural Resource Ecology
and Management

Introduction

To better understand and assess the bat population of Iowa, a survey is needed. One approach to surveying bats is to use acoustic detection equipment.

Materials and Methods

The Iowa Acoustic Bat Surveys covered 10 counties with a total of 19 driving transects, as well as six walking transects conducted in central Iowa parks and farms. The surveys were designed to use Anabat SD2 acoustic monitors to detect and record bat activity. We were able to complete all transects at least two times over the course of our two-month data collection period.

Results and Discussion

Across the state, seven different bats species were recorded at least once (Figure 1). Bat activity levels for individual transects ranged from a minimum of 1.9 percent to a maximum

of 19.7 percent of 15s intervals containing bat activity with a statewide mean of 7.3 percent. Bat activity was consistently higher in eastern Iowa compared with central and southern Iowa.

Central Iowa Farms. Identifying the species that produced a bat call is difficult and most calls could not be narrowed down to a single species. For those calls that could be identified, big brown, hoary, and silver-haired bats were the most frequently identified species at the farms. This is consistent with the results of our driving transects indicating these are the most common species in central Iowa. Other bat species including the tri-colored, evening, and *Myotis* spp. bats were detected. These tend to be more common in eastern Iowa (Figure 1).

For the farms surveyed in central Iowa, the ISU Curtiss Farm had the highest amount of bat activity with 16 percent of 15s intervals containing bat activity, followed closely by the ISU Horticulture Station with 13 percent. The ISU Ag Engineering/Agronomy Farm had the least bat activity with only 3 percent of 15s intervals containing bat calls (Figure 2).




Figure 1. Regional breakdown of bat species by location.




Figure 2. Breakdown of bat species detected in central Iowa by farm location.