

2017

Corn Yield Response to Nitrogen Fertilizer Application Timing in Central Iowa

John Lundvall
Iowa State University

Dan Barker
Iowa State University, dbarker@iastate.edu

John Sawyer
Iowa State University, jesawyer@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/farmprogressreports>

 Part of the [Agriculture Commons](#)

Recommended Citation

Lundvall, John; Barker, Dan; and Sawyer, John (2017) "Corn Yield Response to Nitrogen Fertilizer Application Timing in Central Iowa," *Farm Progress Reports*: Vol. 2016 : Iss. 1 , Article 171.

DOI: <https://doi.org/10.31274/farmprogressreports-180814-1734>

Available at: <https://lib.dr.iastate.edu/farmprogressreports/vol2016/iss1/171>

This Ag Engineering/Agromony, Central Iowa and BioCentury Research Farms is brought to you for free and open access by the Extension and Experiment Station Publications at Iowa State University Digital Repository. It has been accepted for inclusion in Farm Progress Reports by an authorized editor of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Corn Yield Response to Nitrogen Fertilizer Application Timing in Central Iowa

RFR-A16130

John Lundvall, research affiliate
Dan Barker, assistant scientist
John Sawyer, professor
Department of Agronomy

Introduction

The objective of this project was to measure corn yield response to fertilizer nitrogen (N) application timing (spring pre-plant or at-planting vs. split/side-dress) across multiple N rates (0 to 250 lb total-N/acre). Results were determined through fitting yield response to N-rate regression equations and calculation of economic optimum nitrogen rate (EONR) based on a 0.10 N fertilizer price-to-corn price ratio.

Materials and Methods

The project was conducted in 2015 to 2016 at ISU Research and Demonstration Farms near Sutherland, Kanawha, Nashua, Ames, Lewis, and Crawfordsville. At all sites, corn was in rotation with soybean.

Multiple total-N rates were applied preplant (PRE) or preplant + side-dress at approximately V5 to V6 corn growth stage (SPLIT). Treatments were arranged in a randomized complete block design, with four replications. Preplant fertilizer-N application rates included 0, 50, 100, 150, 200, and 250 lb N/acre. Split fertilizer-N application rates included the same total in 0/0, 0/50, 50/50, 50/100, 50/150, and 50/200 lb N/acre splits.

Corn was grown with either no-till management or spring disk-field cultivator tillage for seedbed preparation. Fertilizer-N sources were injected urea-ammonium nitrate solution or surface broadcast/incorporated

urea. Adapted corn hybrids were planted in 30-in. row spacing.

Results and Discussion

At Ames, urea-ammonium nitrate (UAN) solution (32-0-0) fertilizer was coulter-injected as the PRE (5/6/16) and side-dress (6/7/16) N application. Corn was planted into a tilled seedbed May 12, 2016. Results from the Ames site underscored the effect of growing season precipitation (and timing of precipitation) on corn yield and optimum N fertilizer rate. In 2016, Ames received about 33 in. of precipitation during the March to October growing season. Rainfall totals included over 8 inches in April and May and 21.5 in. from July 1 to September 30. These rainfall totals indicate potential for greater-than-normal yield response to N application and higher-than-normal N fertilization requirement in 2016. There were no excessive rainfall events immediately following PRE N application on May 6 and SPLIT N application June 7. However, rainfall totals in the week after PRE N and SPLIT N applications afforded sufficient moisture to move injected UAN into the soil profile and root zone.

Averaged across PRE and SPLIT N application timings with no N applied, corn yielded 118 bushels/acre in 2016. The EONR averaged 146 lb N/acre in 2016 (171 lb N/acre PRE and 121 lb N/acre SPLIT). Corn yield at the EONR in 2016 was 206 bushels/acre PRE and 210 bushels/acre SPLIT.

Across site-years from multiple studies, the application timing results were grouped according to each site's results for timing within ± 10 lb N/acre of the EONR (Table 1). Response to application timing was mixed,

and there was no consistent corn yield or fertilizer rate difference associated with the PRE or SPLIT application. Over all sites, the SPLIT application had a lower EONR (only 6 lb N/acre), with no difference in corn yield at the EONR. For four sites, the calculated EONR for the SPLIT averaged 29 lb N/acre less than the PRE application; however, corn yield was unchanged. For three sites, the EONR for the PRE averaged 18 lb N/acre less than the SPLIT application, however, corn yield was only 3 bushels/acre different and higher with the SPLIT application. For seven

sites, the EONR was within 10 lb N/acre for the PRE and SPLIT applications, and the yield at the EONR the same. These results indicate a combination of weather and soil properties can significantly influence corn response to springtime N application timing. One would not expect one or the other timing to always be the best.

Acknowledgements

Appreciation is extended to the farm superintendents and their staff for assistance with this project.

Table 1. Effect of application timing, spring preplant (Pre) or split Pre/side-dress (Split), on calculated economic optimum nitrogen fertilizer rate (EONR) and corresponding corn yield (YEONR) across 14 site-years in 2014 to 2016.

Timing response category	Sites	Mean EONR		Mean YEONR	
		Pre	Split	Pre	Split
		--- lb N/ac ---		--- bu/ac ---	
Split EONR at least 10 lb N/acre lower than Preplant	4	167	138	202	201
Preplant EONR at least 10 lb N/acre lower than Split	3	108	126	203	206
Preplant and Split EONR within 10 lb N/acre	7	151	147	221	221
Overall Mean	14	146	140	212	212

Based on N response equations and 0.10 N/corn price ratio. The table includes additional sites from 2014 not discussed for individual research farms (Lundvall, Barker, Sawyer, and Hall, 2014-2016).