

7-19-2010

## Late Summer Seeding of Forage Crops

Stephen K. Barnhart  
*Iowa State University*

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

---

### Recommended Citation

Barnhart, Stephen K., "Late Summer Seeding of Forage Crops" (2010). *Integrated Crop Management News*. 397.  
<http://lib.dr.iastate.edu/cropnews/397>

**The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.**

---

# Late Summer Seeding of Forage Crops

## **Abstract**

Late summer can be an excellent time to establish forage crops, provided there is sufficient moisture for germination and good seedling growth. It is also a good time to seed bare or thin spots in spring established forage stands. The following steps will improve the chances for successful late summer forage stand establishment.

## **Keywords**

Agronomy

## **Disciplines**

Agricultural Science | Agriculture | Agronomy and Crop Sciences

[Subscribe to Crop News](#)



#### Archives

[2015](#)[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)

#### ISU Crop Resources

[Extension Field Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology Extension](#)[Ag and Biosystems Engineering Extension](#)[Agribusiness Education Program](#)[Iowa Grain Quality Initiative](#)[College of Agriculture and Life Sciences](#)[ISU Extension](#)

# Integrated Crop Management NEWS

 PRINT STORY  
 EMAIL STORY  
 ADD TO DELICIOUS  
 ATOM FEED  
 FOLLOW ON TWITTER

## Late Summer Seeding of Forage Crops

By Stephen K. Barnhart, Department of Agronomy

Late summer can be an excellent time to establish forage crops, provided there is sufficient moisture for germination and good seedling growth. It is also a good time to seed bare or thin spots in spring established forage stands. The following steps will improve the chances for successful late summer forage stand establishment.

### Plan-Ahead

For best success, taking care of some preparatory steps a few months, or years, ahead of the actual seeding may be necessary.

- Test soils and apply needed, corrective lime and/or fertilizer during previous cropping seasons; corrective fertilizer can be incorporated during forage seedbed preparation.
- Begin to control problem perennial weeds a year or more ahead of seeding.
- Be careful with herbicide selection in crops grown in the field before the forage seeding because some may have residual soil activity and will harm new forage seedings if proper waiting periods are not observed. Read the labels for details.

### This is late-summer seeding, not fall seeding

- Seed as early as possible. Seedlings require six to eight weeks of growth after emergence to have adequate vigor to survive the winter. Seed by Aug. 10 in the northern third of Iowa, by Aug. 20 in central parts of the state and by Sept. 1 in southern Iowa.
- Forage legumes such as red clover and alfalfa can be seeded up to the dates listed above if moisture is present. Slow establishing species like birdsfoot trefoil or reed canarygrass should be planted in early August. Most forage grasses such as Kentucky bluegrass, smooth brome grass, orchardgrass, tall fescue and timothy can actually be seeded 15 days or more later than the dates listed above.
- Don't plant warm-season prairie grasses as a late-summer seeding.
- Keep in mind that the above dates assume sufficient moisture to establish the crop. Planting later than the dates mentioned above is sometimes successful depending on fall and winter weather patterns, but there is increased risk of failure and reduced yield potential for the stand as planting is delayed.
- A good rule of thumb for alfalfa is to have six to eight inches of growth before a killing frost.

### Prepare a firm seedbed if using tillage

Loose seedbeds dry out very quickly. Deep tillage should be completed several weeks ahead of seeding so rains can settle the soil before final seedbed preparation. A cultipacker or roller is an excellent last-pass tillage tool. The soil should be firm enough for a footprint to sink no deeper than 3/8 to 0.5 inch.

**Late-summer pasture interseeding and no-till forage seeding is an excellent way to conserve moisture**, provided weeds are controlled prior to seeding. Remove all straw after small grain harvest. Any remaining stubble

should either be left standing, or clipped and removed. Do not leave clipped stubble in fields as it forms a dense mat that prevents good emergence.

**Don't plant alfalfa immediately after older established alfalfa.**

Autotoxic compounds are released by old alfalfa plants that inhibit growth and productivity of new alfalfa seedlings. It is best to rotate to another crop for a year or more before going back to alfalfa; however, thickening up seedings within 12 to 15 months of the original planting date is considered to be a low-risk practice because autotoxicity concerns are greatest with older alfalfa fields.

**Seed when soil moisture is adequate or a good rain system is in the forecast.** There is a higher risk of seedling failure when planting seeds into dry soil, as there may be just enough moisture to germinate the seed but not enough for seedling establishment.

**Plant seed shallow and in firm contact with the soil.**

Carefully check seeding depth, especially when no-tilling. Drills with press wheels usually provide the greatest success in the summer. Broadcasting seed on the surface without good soil coverage and without firm packing is usually a recipe for failure in the summer.

**Use high quality seed of known varieties.**

Cheap seed often results in big disappointments and shorter stand life. Make sure legume seed has fresh inoculum of the proper rhizobium.

**Don't harvest new summer seedings this fall.**

Allow them to establish well and develop winter hardiness.

*Stephen K. Barnhart is a professor of agronomy and the Iowa State University Extension forage agronomist. He can be reached at 515-294-7835 or by emailing [sbarnhar@iastate.edu](mailto:sbarnhar@iastate.edu).*

---

This article was published originally on 7/19/2010. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.