

5-7-2010

Temperature Fluctuations Have and May Continue to Inhibit Corn Emergence

Roger Elmore
Iowa State University

Lori Abendroth
Iowa State University, labend@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Climate Commons](#)

Recommended Citation

Elmore, Roger and Abendroth, Lori, "Temperature Fluctuations Have and May Continue to Inhibit Corn Emergence" (2010). *Integrated Crop Management News*. 451.
<http://lib.dr.iastate.edu/cropnews/451>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Temperature Fluctuations Have and May Continue to Inhibit Corn Emergence

Abstract

Corn planting proceeded at an all-time record this spring. We've experienced one of the best planting seasons ever in Iowa. According to the most recent USDA-NASS report (2 May 2010), 84 percent of Iowa's corn was planted with cropping districts ranging from 69 percent complete in south central Iowa to 92 percent complete in north central Iowa. Overall, corn has been planted within the recommended window which should serve to maximize yield potential relative to planting date.

Keywords

Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Climate

[Subscribe to Crop News](#)

Archives

[2015](#)[2014](#)[2013](#)[2012](#)[2011](#)[2010](#)[2009](#)[2008](#)[Previous Years](#)

ISU Crop Resources

[Extension Field Agronomists](#)[Crop & Soils Info](#)[Pesticide Applicator Training](#)[Agronomy Extension](#)[Entomology Extension](#)[Plant Pathology Extension](#)[Ag and Biosystems Engineering Extension](#)[Agribusiness Education Program](#)[Iowa Grain Quality Initiative](#)[College of Agriculture and Life Sciences](#)[ISU Extension](#)

Integrated Crop Management NEWS

-  PRINT STORY
-  EMAIL STORY
-  ADD TO DELICIOUS
-  ATOM FEED
-  FOLLOW ON TWITTER

Temperature Fluctuations Have and May Continue to Inhibit Corn Emergence

By Roger Elmore and Lori Abendroth, Department of Agronomy

Corn planting proceeded at an all-time record this spring. We've experienced one of the best planting seasons ever in Iowa. According to the most recent [USDA-NASS report](#) (2 May 2010), 84 percent of Iowa's corn was planted with cropping districts ranging from 69 percent complete in south central Iowa to 92 percent complete in north central Iowa. Overall, corn has been planted within the [recommended window](#) which should serve to maximize yield potential relative to planting date.

Soil and weather conditions at planting and emergence were excellent for most of the state. Alison Robertson recently emphasized the need to assess seedling health when doing stand counts in a [May 4, 2010 ICM article](#) in response to some of the problems occurring in southeast Iowa this spring. Seedling emergence problems there are correlated with swings in April soil temperature.

Cooler soil temperatures slow the germination process and predispose seedlings to fungal infection. We have also observed, or heard reports of, seedling growth problems in some parts of Iowa already this year, including:

- [Imbibitional chilling](#) damage, which is the chilling effect seeds may experience when they imbibe, or absorb, water when soil temperatures are less than 55 F for an extended time. Seedlings may "corkscrew" or not emerge when exposed to these cool soil temperatures. This may happen also when there are rapid swings in air temperatures of nearly 30 F.
- Soil crusting due to wet soils at planting or heavy rains after planting reducing plant stands. Significant [stand reductions](#) lower yield potential.
- "Leafing out" underground - see Photo 1. This occurs most often in crusted soils and also appears associated with imbibitional chilling, mentioned above.
- [Variable plant emergence](#) and reduced plant population - see Photo 2. Variable emergence and growth will reduce yield.

Cool Temperatures Soon?

The forecast for the next several days is for air temperatures dropping into the high 30's at night; soil temperatures will also lower respectively. Growers and agronomists should pay particular attention to corn fields planted recently as they are the most likely to exhibit seedling rot or poorer emergence associated with some of the issues mentioned here. [Frost](#) is likely to occur in low lying areas causing leaf necrosis and delayed growth of sprouted seeds.

Any of the situations mentioned above may result in a need to replant. [Assess](#)

[stands](#) well before making this decision.

Irrespective of all of these issues, 2010 has provided one of the best planting seasons to date. Most of Iowa's corn was planted within a timeframe that should serve to maximize yield potential.



Photo 1. Seedling leaved out under soil surface. Story Co. IA, 4 May 2010. Roger Elmore.



Photo 2. Variable seedling emergence dates can reduce yields. Story Co. IA, 4 May 2010. Roger Elmore. See [additional photos](#) for leaving out underground and variable seedling emergence.

Roger Elmore is a professor of agronomy with research and extension responsibilities in corn production. Lori Abendroth is an agronomy specialist with research and extension responsibilities in corn production. Elmore can be contacted by email at relmore@iastate.edu or (515) 294-6655; Abendroth can be contacted by email at labend@iastate.edu or (515) 294-5692.

This article was published originally on 5/6/2010. The information contained within the article may or may not be up to date depending on when you are accessing the information.

Links to this material are strongly encouraged. This article may be republished without further permission if it is published as written and includes credit to the author, Integrated Crop Management News and Iowa State University Extension. Prior permission from the author is required if this article is republished in any other manner.