


10-20-2006

Use of Allylic Strain To Enforce Stereochemistry. Direct Syntheses of 7,8-Dihydroxycalamenene and Mansonone C

George A. Kraus
Iowa State University, gakraus@iastate.edu

Insik Jeon
Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/chem_pubs

 Part of the [Environmental Chemistry Commons](#), [Organic Chemistry Commons](#), [Other Chemistry Commons](#), and the [Polymer Chemistry Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/chem_pubs/467. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Chemistry at Iowa State University Digital Repository. It has been accepted for inclusion in Chemistry Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Use of Allylic Strain To Enforce Stereochemistry. Direct Syntheses of 7,8-Dihydroxycalamenene and Mansonone C

Abstract

Direct syntheses of 7,8-dihydroxycalamenene and mansonone C were achieved. The *cis*-stereochemistry required for the synthesis of 7,8-dihydroxycalamenene was introduced by an intramolecular cyclization directed by allylic strain.

Disciplines

Chemistry | Environmental Chemistry | Organic Chemistry | Other Chemistry | Polymer Chemistry

Comments

Reprinted (adapted) with permission from *Organic Letters*, 8(23); 5315-5316. Doi: [10.1021/ol0621194](https://doi.org/10.1021/ol0621194).
Copyright 2006 American Chemical Society.

Use of Allylic Strain To Enforce Stereochemistry. Direct Syntheses of 7,8-Dihydroxycalamenene and Mansonone C


George A. Kraus* and Insik Jeon

Department of Chemistry, Iowa State University, Ames, Iowa 50011

gakraus@iastate.edu

Received August 28, 2006

ABSTRACT



Direct syntheses of 7,8-dihydroxycalamenene and mansonone C were achieved. The *cis*-stereochemistry required for the synthesis of 7,8-dihydroxycalamenene was introduced by an intramolecular cyclization directed by allylic strain.

Allylic 1,3-strain has been used in acyclic systems to direct the introduction of new stereogenic centers.¹ Notable examples include the work of Kishi,² Adam,³ and Giese.⁴ We are not aware of any application of allylic strain to control the relative stereochemistry in disubstituted tetralins. Tetralins such as **1**, **3**, and **4** have attracted considerable synthetic attention (Figure 1).⁵ 7,8-Dihydroxycalamenene (**1a**) exhibits

extracted from the heartwood of *Mansonia altissima*,⁸ was found to possess promising antifungal, larvicidal, and anti-oxidant properties.⁹ Schmalz recorded a synthesis of **1a** using arenechromium complexes to introduce the relative stereochemistry.¹⁰

Many synthetic approaches to these compounds begin with natural products such as menthone wherein the relative stereochemistry has already been established.¹¹

Several researchers have noted that attempts to install stereochemistry by epimerization or by cyclization onto the aromatic ring have led to mixtures.^{12,13} We report herein that a system such as **5**, wherein allylic strain between G and




Figure 1. Tetralin-derived products and mansonone C.

useful anti-infective activity.⁶ Hydroxycalamenene (**1b**) was isolated from *Hypericum elodeoides*.⁷ Mansonone C (**2**),

(1) Hoffmann, R. W. *Chem. Rev.* **1989**, *89*, 1841. Johnson, F. *Chem. Rev.* **1968**, *68*, 375.

(2) Johnson, M. R.; Nakata, T.; Kishi, Y. *Tetrahedron Lett.* **1979**, 4343.

(3) Adam, W.; Wirth, T. *Acc. Chem. Res.* **1999**, *32*, 703.

(4) Giese, B.; Bulliard, M.; Zeitz, H. G. *Synlett* **1991**, 425.

(5) Erogorgiaene (**3**): Rodriguez, A. D.; Ramirez, C. *J. Nat. Prod.* **2001**, *64*, 100. Elisapterosin B (**4**): Rodriguez, A. D.; Ramirez, C.; Rodriguez, I. I.; Barnes, C. L. *J. Org. Chem.* **2000**, *65*, 1390.

(6) Wahyouno, S.; Hoffmann, J. J.; Bates, R. B.; McLaughlin, S. P. *Phytochemistry* **1991**, *30*, 2175.

(7) Mathela, D. K.; Mathela, C. S.; Dev, V. *J. Indian Chem. Soc.* **1984**, *61*, 792.

(8) Marini Bettolo, G. B.; Casinovi, C. G.; Galeffi, C. *Tetrahedron Lett.* **1965**, 4857.

(9) Tiew, P.; Ioset, J.-R.; Kokpol, U.; Chavasiri, W.; Hostettmann, K. *Phytother. Res.* **2003**, *17*, 190.


(10) Schmalz, H. G.; Hollander, J.; Arnold, M.; Duerner, G. *Tetrahedron Lett.* **1993**, *34*, 6259.

(11) Serra, S.; Fuganti, C. *Tetrahedron Lett.* **2005**, *46*, 4769.


(12) Ramaiah, P.; Rao, A. S. *Synth. Commun.* **1989**, *19*, 931.

(13) Beckwith, P. L. M.; Ghisalberti, E. L.; Jefferies, P. R.; Raston, C. L.; White, A. H. *Aust. J. Chem.* **1988**, *41*, 351.

the methyl group forces the methyl group to be axial as the six-membered ring is being formed, affords exclusively the *cis*-stereoisomer **7** (Scheme 1).




In order to evaluate the directing effect, we first synthesized allylic acetate **8** (Scheme 2). Allylic acetate **8** was synthesized starting from 6-methyl-5-hepten-2-one and the anion of 1,4-dimethoxybenzene. Dehydroxylation of the resulting benzylic alcohol using Li/NH₃ followed by allylic oxidation by the method of Sharpless¹⁴ yielded an allylic alcohol which, upon acetylation, afforded **8**. Surprisingly, cyclization of allylic acetate **8** using the conditions of Ma and Zheng¹⁵ afforded tetralin **10** and naphthalene **11** in 51% and 37% yields, respectively. Compound **9**, the expected product, was not isolated.




We believe that compounds **10** and **11** result from a novel cation-mediated disproportionation reaction. Comparison of the NMR spectrum of **10** with that of the literature compound¹³ showed that the *cis*-stereoisomer was exclusively formed.

In support of this allylic strain assisted stereoselective cyclization, we have found that cyclization of allylic acetate **12**, which does not contain the directing group G, affords tetralin **13** as a 1.3:1 mixture of diastereomers (Scheme 3).

With the stereochemistry of **10** established, we began the synthesis of **1a** by the reaction of 5-methyl-4-hexen-1-ol¹⁶



with the anion of 1,2-bis(methoxymethoxy)-3-methylbenzene (Scheme 4). The resulting alkoxide was in situ acetylated by the addition of acetic anhydride. The displacement of the acetate to the corresponding methyl group was achieved using Me₃Al to afford **15**. In order to enhance the stability for the acid-mediated cyclization, MOM protecting groups were converted to the more stable methyl ether **16**. It was then oxidized and acetylated by the same methods used to generate **8**. Cyclization of **17** using trifluoroacetic acid in acetic acid at 70 °C for 12 h followed by deprotection by the use of BBr₃ generated 7,8-dihydroxycalamenene (**1a**) in two steps (30% yield). Interestingly enough, mansonone C (**2**) was also obtained from the same reaction in two steps (46% yield). The direct synthesis of 7,8-dihydroxycalamenene (**1a**) demonstrates the advantage of employing the concept of allylic strain in organic synthesis.



Acknowledgment. We thank the National Institutes of Health (Grant No. P01 ES12020) and the Office of Dietary Supplements for partial financial support through the Center for Research on Botanical Dietary Supplements at Iowa State University.

Supporting Information Available: Experimental procedures, characterization data, and NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0621194

(15) Ma, S.; Zhang, J. *Tetrahedron* **2003**, *59*, 6273.
(16) Kraus, G. A.; Kim, J. *Org. Lett.* **2004**, *6*, 3115.

(14) Umbreit, M. A.; Sharpless, K. B. *J. Am. Chem. Soc.* **1977**, *99*, 5526