

2019

Photon radiation in hot nuclear matter by means of chiral anomalies

Kirill Tuchin

Iowa State University, tuchin@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/physastro_pubs



Part of the [Plasma and Beam Physics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/physastro_pubs/488. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Physics and Astronomy at Iowa State University Digital Repository. It has been accepted for inclusion in Physics and Astronomy Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Photon radiation in hot nuclear matter by means of chiral anomalies

Abstract

A new mechanism of photon emission in the quark-gluon plasma is proposed. Photon dispersion relation in the presence of the CP-odd topological regions generated by the chiral anomaly acquires an imaginary mass. It allows photon radiation through the decay $q \rightarrow q\gamma$ and annihilation $qq^{\bar{}} \rightarrow \gamma$ processes closely related to the chiral Cherenkov radiation. Unlike previous proposals this mechanism does not require an external magnetic field. The differential photon emission rate per unit volume is computed and shown to be comparable to the rate of photon emission in conventional processes.

Disciplines

Physics | Plasma and Beam Physics

Comments

This is a pre-print of the article Tuchin, Kirill. "Photon radiation in hot nuclear matter by means of chiral anomalies." *arXiv preprint arXiv:1903.02629* (2019). Posted with permission.

Photon radiation in hot nuclear matter by means of chiral anomalies

Kirill Tuchin¹

¹*Department of Physics and Astronomy, Iowa State University, Ames, Iowa, 50011, USA*

(Dated: March 8, 2019)

A new mechanism of photon emission in the quark-gluon plasma is proposed. Photon dispersion relation in the presence of the CP -odd topological regions generated by the chiral anomaly acquires an imaginary mass. It allows photon radiation through the decay $q \rightarrow q\gamma$ and annihilation $q\bar{q} \rightarrow \gamma$ processes closely related to the chiral Cherenkov radiation. Unlike previous proposals this mechanism does not require an external magnetic field. The differential photon emission rate per unit volume is computed and shown to be comparable to the rate of photon emission in conventional processes.

I. INTRODUCTION

Photon radiation by hot nuclear matter has been a focus of experimental and theoretical studies for many decades. However, in spite of considerable progress, there are still unresolved problems concerning the photon spectrum produced in relativistic heavy ion collisions such as the puzzling enhancement of the direct photon production [1, 2]. The major contributors to the photon spectrum are the quark–antiquark annihilation and the QCD Compton scattering processes in the quark-gluon plasma and the inelastic reactions in the hot hadronic gas [3–18]. In addition to these “conventional” processes, other contributors have been proposed such as photon emission by the nuclear matter before the QGP formation [19, 20], the synchrotron radiation [21, 22], radiation via the conformal anomaly [23] and through the chiral anomaly [24] as well as the modification of the conventional processes due to the axial charge fluctuations [25, 26]. The mechanisms suggested in Refs. [21, 23, 24] rely on existence of intense magnetic field produced in heavy-ion collisions. In this paper we argue that there is a different unconventional mechanism of photon production via the chiral anomalies of QED and QCD which *does not* involve the external magnetic field.

The hot nuclear matter, or quark-gluon plasma (QGP), is believed to contain the topological CP -odd domains created by the random sphaleron-mediated transitions between different QCD vacua. Interaction of the electromagnetic field with these domains can be described by adding to the QED Lagrangian the axion-photon coupling term [27]

$$\mathcal{L}_A = -\frac{c_A}{4}\theta F_{\mu\nu}\tilde{F}^{\mu\nu}, \quad (1)$$

where $c_A = N_c \sum_f q_f^2 e^2 / 2\pi^2$ is the QED anomaly coefficient and the field θ is sourced by the topological charge density

$$q(x) = \frac{g^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a\mu\nu}(x) \quad (2)$$

which varies in space and time across a CP -odd domain. As a result (1) cannot be rewritten as a total derivative and removed from the Lagrangian. Instead, it appears in the modified Maxwell equations as the spatial and the temporal derivatives of θ .

It has been known since the pioneering article by Carroll, Feld and Jackiw [28] that in QED coupled to the axion field, photons acquire an imaginary mass m_A making possible their spontaneous emission by fermions. This phenomenon is referred to as the vacuum Cherenkov radiation [29, 30]. Since the electromagnetic field in QGP is coupled to the axion field θ , it is natural to expect that a similar mechanism of photon radiation exists in hot nuclear medium as well. This idea was developed in [31, 32] where it was argued that ultrarelativistic fermions moving in a finite- θ domain radiate photons, which we referred to as the *chiral* Cherenkov radiation. Additionally, fermions in QGP radiate the *chiral* transition radiation as they cross the boundary between the plasma and vacuum due to the difference in the photon wave function inside and outside the plasma. The spectra of both processes are proportional to the average values of the spatial and the temporal θ -derivatives. Since the chiral Cherenkov radiation scales with the system volume, whereas the chiral transition radiation scales with its area, the former is dominant when the contribution of the entire QGP (as opposed to a single fast quark) is considered. Thus, the present work focuses on the chiral Cherenkov radiation by QGP.

The analysis of [31, 32] relied on two basic assumptions: (i) $\theta(x)$ is a slowly varying adiabatic function of its arguments and (ii) the absolute value of the photon mass generated by the anomaly $|m_A|$ is much larger than the plasma frequency ω_{pl} . The first assumption is the simplest model that captures the essential dynamics of the chiral magnetic effect [33–36]. It is supported by the results obtained by Zhitnitsky [37, 38]. The second assumption is justified for large enough photon energy ω because m_A^2 is proportional to ω , see (10), whereas the plasma frequency is ω -independent. These are the assumptions that are carried over to the present study as well. However, unlike the radiation by a single quark discussed in [31, 32] where one is free to choose the quark energy high enough so that most of the photon spectrum satisfy $|m_A| \gg \omega_{\text{pl}}$, in the case of QGP the bulk of the photon radiation occurs at $\omega \lesssim T$, where T is the QGP temperature. Still, it is argued in the next section that at high enough temperatures, the photon mass satisfies the second assumption since the plasma frequency is proportional to T , see (8), whereas the absolute value of m_A^2 is proportional

to the sphaleron transition rate Γ which rises at high temperatures as T^4 .

The paper is organized as follows. Sec. II deals with the qualitative discussion of the electromagnetic fields in presence of the CP -odd domains. The mean value of the θ -field in a domain is related to the sphaleron transition rate and hence scales as T^4 at high temperatures. This indicates that at high enough temperatures the chiral Cherenkov radiation becomes possible. In Sec. III, the photon dispersion relation at finite θ is discussed. The main section is Sec. IV where the photon radiation rate is computed. In order to simplify the derivations and emphasize the main physics points, I am going to consider the relativistic limit $\omega \gg |m_A|$; generalization beyond this limit is straightforward. In fact, such a generalization for a single quark has been recently obtained in [39]. The discussion and summary is presented in Sec. V.

II. ELECTRODYNAMICS IN QUARK-GLUON PLASMA WITH CP -ODD DOMAINS

The CP -odd domains in the chiral matter can be described by the axion – a pseudo-scalar field θ whose interactions with the electromagnetic $F_{\mu\nu}$ and color $G_{\mu\nu}^a$ fields are governed by the Lagrangian [28, 40–42]

$$\mathcal{L} = \mathcal{L}_{\text{QED}} + \mathcal{L}_{\text{QCD}} - \frac{c_A}{4}\theta F_{\mu\nu}\tilde{F}^{\mu\nu} - \frac{c'_A}{4}\theta G_{\mu\nu}^a\tilde{G}^{a\mu\nu} + f^2 \left[\frac{1}{2}(\partial_\mu\theta)^2 - \frac{1}{2}m_{\text{ax}}^2\theta^2 \right], \quad (3)$$

where $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\lambda\rho}F^{\lambda\rho}$ is the dual field tensor, c_A, c'_A are the QED and QCD anomaly coefficients respectively and f, m_{ax} are constants with mass dimension one. It follows that the axion equation of motion is

$$(\partial^2 + m_{\text{ax}}^2)\theta = -\frac{1}{4f^2} \left(c'_A G_{\mu\nu}^a \tilde{G}^{a\mu\nu} + c_A F_{\mu\nu} \tilde{F}^{\mu\nu} \right). \quad (4)$$

In the quark-gluon plasma the electromagnetic contribution to the topological charge density is presumed to be negligible so that the axion dynamics is driven primarily by the topologically non-trivial gluon configurations. Assuming that θ is slowly varying inside a CP -odd domain one can express it in terms of the topological number density (2) as

$$\theta(x) = -\frac{N_f q(x)}{f^2 m_{\text{ax}}^2}. \quad (5)$$

The equations of motion of electromagnetic field coupled to the axion field read

$$\partial_\mu F^{\mu\nu} = j^\nu - c_A \tilde{F}^{\mu\nu} \partial_\mu \theta, \quad (6)$$

$$\partial_\mu \tilde{F}^{\mu\nu} = 0. \quad (7)$$

In a slowly varying field θ , its first derivatives $\partial^\mu\theta$ can be replaced by their constant domain-average values denoted by $\sigma_\chi = c_A\dot{\theta}$ [35, 36, 43], referred to as the chiral conductivity, and $\mathbf{b} = c_A\nabla\theta$. In this approximation photon and axion dynamics decouple and one can consider electrodynamics in the topologically non-trivial background [44].

The average of the axion field over an ensemble of CP -odd domains vanishes. However, its value in a single domain can be finite due to the fluctuations of the topological number N_{CS} . In the context of this work we need to know the temperature dependence of the axion field in a domain because it determines the temperature dependence of the effective photon mass m_A . In particular, if its T -dependence is steeper than linear, then one expects that there is a range of temperatures where the plasma becomes radioactive as explained at the end of Sec. I. The topological number density can be estimated as $q \sim N_{CS}/V_{\text{dom}}$, where $V_{\text{dom}} \sim 1/m_{\text{ax}}^4$ is the domain 4-volume. Since the sphaleron size is inversely proportional to T , the domain volume decreases as $V_{\text{dom}} \sim 1/T^4$. Fluctuations of N_{CS} are related to the sphaleron transition rate Γ as $\langle N_{CS}^2 \rangle = 2\Gamma V_{\text{pl}}$ [45] for large enough 4-volume V_{pl} of plasma. Therefore, the variance of the topological number density is $\langle q^2 \rangle \sim m_{\text{ax}}^8 \Gamma V_{\text{pl}}$. Employing (5) it is seen that the typical variance of the axion field strength is $\langle \theta^2 \rangle \sim m_{\text{ax}}^4 \Gamma V_{\text{pl}} / f^4$. Γ is exponentially suppressed at low temperatures, but increases as T^4 at high temperatures [46–49]. It follows, using (10) of the next section, that $m_A \sim \langle \theta \rangle^{1/2} \sim T^4$. Thus, $|m_A|$ exceeds ω_{pl} at high T making the chiral Cherenkov radiation possible.

III. PHOTON DISPERSION RELATION

Now that the model parameters have been outlined, it is instructive to review the photon dispersion relation. In the case $\theta = 0$ the photon dispersion relation at finite temperature T and finite chemical potentials of the right and left-handed fermions $\mu_{R,L}$ was computed in [50]. In the high-energy limit, when the photon is near the mass-shell and transversely polarized, its dispersion relation is $\omega^2 - k^2 = \omega_{\text{pl}}^2$, where

$$\omega_{\text{pl}}^2 = \frac{m_D^2}{2} = \frac{e^2}{2} \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} \right), \quad (8)$$

and $\mu^2 = \mu_R^2 + \mu_L^2$.

At finite θ the photon dispersion relation acquires an extra term due its interaction with the CP -odd domains

$$\omega^2 - k^2 = \omega_{\text{pl}}^2 + m_A^2 + \mathcal{O}(\omega - k), \quad (9)$$

where m_A^2 is given by

$$m_A^2 = -\lambda\sigma_\chi\omega, \quad \text{or} \quad m_A^2 = -\lambda\mathbf{k} \cdot \mathbf{b}, \quad (10)$$

depending on which of the parameters σ_χ or b is largest [32]* and $\lambda = \pm 1$ is the right and left-handed photon polarization. Note that m_A can be real or imaginary. As explained in the previous two sections, at high enough photon energies and plasma temperatures ω_{pl} is but a small correction compared to m_A and will be neglected in the following sections.

IV. PHOTON RADIATION RATE

Photon emission by means of the chiral Cherenkov radiation mechanism can proceed via two channels: (i) the decay channel $q \rightarrow q\gamma$ and (ii) the annihilation channel $q\bar{q} \rightarrow \gamma$.[†] The total photon radiation rate is the sum of rates of these two processes.

A. Decay channel

The scattering matrix element for photon radiation in the decay channel $q(p) \rightarrow q(p') + \gamma(k)$ is given by $S_D = (2\pi)^4\delta^{(4)}(p' + k - p)i\mathcal{M}_D$ where

$$i\mathcal{M}_D = -ieQ \frac{\bar{u}_{\mathbf{p}'s'} \not{\epsilon}_{\mathbf{k}\lambda}^* u_{\mathbf{p}s}}{\sqrt{8\varepsilon\varepsilon'\omega V^3}}. \quad (11)$$

The components of the 4-vectors are $p = (\varepsilon, \mathbf{p})$, $p' = (\varepsilon', \mathbf{p}')$ and $k = (\omega, \mathbf{k})$, Q is quark charge and $m = gT/\sqrt{3}$ its thermal mass [11]. We retained the relativistic normalization factors $(2p^0V)^{-1/2}$ for each of the three fields, where V is the normalization volume. The radiation probability can be computed as

$$dw_D = 2N_c \frac{1}{2} \sum_{\lambda s s'} |S_D|^2 f(\varepsilon) [1 - f(\varepsilon')] \frac{V d^3 p'}{(2\pi)^3} \frac{V d^3 k}{(2\pi)^3} \frac{V d^3 p}{(2\pi)^3}, \quad (12)$$

where $2N_c$ accounts for the number quarks and antiquarks of different color, $1/2$ comes from the incident quark spin average and $f(\varepsilon)$ is the quark equilibrium distribution function, which reads

$$f(\varepsilon) = \frac{1}{e^{\varepsilon/T} + 1}. \quad (13)$$

* In [32] m_A was denoted as μ . The dispersion relations for arbitrary σ_χ and b can be found in [28].

[†] I am using the term ‘the chiral Cherenkov radiation’ with respect to both channels.

The small chemical potentials of quarks is neglected. The rate of photon production per unit volume can be computed as

$$d\Gamma_D = \frac{dw_D}{VT} = 2N_c \frac{\delta(\omega + \varepsilon' - \varepsilon)}{16(2\pi)^5 \varepsilon \varepsilon' \omega} \sum_{\lambda, s, s'} |i\mathcal{M}_D|^2 f(\varepsilon) [1 - f(\varepsilon')] d^3k d^3p. \quad (14)$$

Performing the summation over the transverse photon polarizations using

$$\sum_{\lambda} \epsilon_{\mathbf{k}\lambda}^{\mu} \epsilon_{\mathbf{k}\lambda}^{\nu*} = \begin{cases} 0, & \nu\mu = 0, \\ \delta^{ij} - \frac{k^i k^j}{k^2}, & \nu = i, \mu = j. \end{cases} \quad (15)$$

yields the result

$$\sum_{ss'} |\mathcal{M}_D|^2 = 4 \left[\varepsilon \varepsilon' - m^2 - \frac{(\mathbf{k} \cdot \mathbf{p})(\mathbf{k} \cdot \mathbf{p}')}{k^2} \right]. \quad (16)$$

In the high energy limit the momenta of the initial and final quarks and the photon have a large component, say along the z -direction, that allows one to approximate

$$p_z \approx \varepsilon \left(1 - \frac{p_{\perp}^2 + m^2}{2\varepsilon^2} \right), \quad k_z \approx \omega \left(1 - \frac{k_{\perp}^2 + m_A^2}{2\omega^2} \right), \quad p'_z \approx \varepsilon' \left(1 - \frac{p'_{\perp}{}^2 + m^2}{2\varepsilon'^2} \right). \quad (17)$$

Denoting by $x = \omega/\varepsilon$ the fraction of the incident quark energy carried away by the photon and substituting (17) into (15) one derives

$$\sum_{ss'} |\mathcal{M}_D|^2 = \frac{2}{x^2(1-x)} [q_{\perp}^2(2-2x+x^2) + m^2 x^4], \quad (18)$$

where $\mathbf{q}_{\perp} = x\mathbf{p}_{\perp} - \mathbf{k}_{\perp}$. In the same approximation the energy delta-function can be written as

$$\delta(\omega + \varepsilon' - \varepsilon) \approx 2x(1-x)\varepsilon \delta(q_{\perp}^2 + m_A^2(1-x) + m^2 x^2), \quad (19)$$

Substituting (18) and (19) into (14) and integrating over q_{\perp} instead of p_{\perp} one finds

$$\omega \frac{d\Gamma_D}{d^3k} = 2N_c \frac{e^2 Q^2 \pi}{4(2\pi)^5} \int_0^1 \frac{dx}{x^4} f\left(\frac{\omega}{x}\right) \left[1 - f\left(\frac{\omega(1-x)}{x}\right) \right] \sum_{\lambda} \left\{ -m_A^2 [(1-x)^2 + 1] - 2m^2 x^2 \right\} \theta(-\kappa_{\lambda}), \quad (20)$$

where it is denoted

$$\kappa_{\lambda} = m_A^2(1-x) + m^2 x^2. \quad (21)$$

Evidently, since $m^2 > 0$ the non-vanishing contribution to the photon production rate in this channel exists only if $m_A^2 < 0$. Moreover, κ_{λ} is negative only if $|m_A^2|(1-x) > x^2 m^2$ which occurs when

$$0 \leq x < \frac{|m_A^2|}{2m^2} \left(\sqrt{1 + \frac{4m^2}{|m_A^2|}} - 1 \right). \quad (22)$$

One can perform the integration of x explicitly in the limit $m \ll |m_A|$. It is convenient to introduce a new variable $\xi = 1/x - 1$ in place of x and rewrite (20) as

$$\omega \frac{d\Gamma_D}{d^3k} = 2N_c \frac{e^2 Q^2}{8(2\pi)^4} \int_0^\infty d\xi \{ -m_A^2 [\xi^2 + (1-\xi)^2] - 2m^2 \} f(\omega(1+\xi)) [1 - f(\omega\xi)] \times \theta(-m_A^2 \xi(1+\xi) - m^2), \quad (23)$$

where only the photon polarization that gives $m_A^2 < 0$ contributes. Neglecting m one obtains

$$\omega \frac{d\Gamma_D}{d^3k} = 2N_c \frac{e^2 Q^2}{8(2\pi)^4} |m_A^2| \int_0^\infty d\xi [\xi^2 + (1-\xi)^2] f(\omega(1+\xi)) [1 - f(\omega\xi)]. \quad (24)$$

Note that the condition (22) is now trivial $0 < x < 1$. Also, since $e \ll g$, $\omega_{\text{pl}} \ll m$ implying that $m_A \approx m_A$ in this approximation. We also approximate $1 - f(\omega\xi) \approx 1 - (e+1)^{-1} = 0.73$ since the argument of f is typically on the order of unity, for otherwise the distribution $f(\omega(1+\xi))$ of the incident quark is exponentially suppressed. Thus we derive

$$\omega \frac{d\Gamma_D}{d^3k} = 0.73 \cdot 2N_c \frac{e^2 Q^2}{8(2\pi)^4} |m_A^2| \left[\frac{\ln(1 + e^{-\beta\omega})}{\beta\omega} + \frac{2\text{Li}_2(-e^{-\beta\omega})}{(\beta\omega)^2} - \frac{4\text{Li}_3(-e^{-\beta\omega})}{(\beta\omega)^3} \right]. \quad (25)$$

The low and high energy regions of the spectrum read

$$\omega \frac{d\Gamma_D}{d^3k} = 0.73 \cdot 2N_c \frac{e^2 Q^2}{8(2\pi)^4} |m_A^2| \begin{cases} \frac{3\zeta(3)}{(\beta\omega)^3}, & \omega \ll T \\ \frac{1}{\beta\omega} e^{-\beta\omega}, & \omega \gg T. \end{cases} \quad (26)$$

Taking into account that m_A^2 is proportional to ω , we find that at $\omega \ll T$, the photon of spectrum scales as $1/\omega^2$. Thus the total photon rate Γ_D is dominated by soft photons $\omega \ll T$ that produce the large logarithm $\ln(T/m)$.

B. Annihilation channel

The scattering matrix element for photon radiation in the annihilation channel $q(p) + \bar{q}(p_1) \rightarrow \gamma(k)$ is given by $S_A = (2\pi)^4 \delta^{(4)}(p + p_1 - k) i\mathcal{M}_A$ where

$$i\mathcal{M}_A = -ieQ \frac{\bar{v}_{\mathbf{p}_1 s_1} \not{\epsilon}_{\mathbf{k}\lambda}^* u_{\mathbf{p}s}}{\sqrt{8\varepsilon_1 \omega V^3}}. \quad (27)$$

The corresponding radiation probability can be computed as

$$dw_A = N_c \frac{1}{4} \sum_{\lambda s s'} |S_A|^2 f(\varepsilon) f(\varepsilon_1) \frac{V d^3 p_1}{(2\pi)^3} \frac{V d^3 k}{(2\pi)^3} \frac{V d^3 p}{(2\pi)^3} \quad (28)$$

where N_c accounts for different colors and $1/4$ stems from the incident quark and antiquark spin average. The rate of photon production per unit volume reads

$$d\Gamma_A = \frac{dw_A}{VT} = N_c \frac{\delta(\omega - \varepsilon_1 - \varepsilon)}{32(2\pi)^5 \varepsilon \varepsilon_1 \omega} \sum_{\lambda s s_1} |i\mathcal{M}_A|^2 f(\varepsilon) f(\varepsilon_1) d^3 k d^3 p. \quad (29)$$

Summation over the transverse photon polarizations using (15) yields

$$\sum_{ss_1} |\mathcal{M}_A|^2 = 4 \left[\varepsilon \varepsilon_1 + m^2 - \frac{(\mathbf{k} \cdot \mathbf{p})(\mathbf{k} \cdot \mathbf{p}_1)}{\mathbf{k}^2} \right]. \quad (30)$$

Employing the high energy limit (17) and denoting by $y = \varepsilon/\omega$ the energy fraction that the incident quark contributed to the photon energy and $\ell_\perp = y\mathbf{k}_\perp - \mathbf{p}_\perp$ one derives

$$\sum_{ss_1} |\mathcal{M}_A|^2 = \frac{2}{y(1-y)} [\ell_\perp^2 (y^2 + (1-y)^2) + m^2], \quad (31)$$

and

$$\delta(\omega - \varepsilon_1 - \varepsilon) \approx 2y(1-y)\omega \delta(\ell_\perp^2 - m_A^2 y(1-y) + m^2). \quad (32)$$

These formulas can also be obtained from the results of the previous subsection using the crossing-symmetry. Substituting (31) and (32) into (29) and integrating over ℓ_\perp instead of p_\perp one finds

$$\omega \frac{d\Gamma_A}{d^3k} = N_c \frac{e^2 Q^2 \pi}{8(2\pi)^5} \int_0^1 dy f(y\omega) f((1-y)\omega) \sum_\lambda [m_A^2 (2y^2 - 2y + 1) + 2m^2] \theta(-\varkappa_\lambda), \quad (33)$$

where it is denoted

$$\varkappa_\lambda = -m_A^2 y(1-y) + m^2. \quad (34)$$

In the annihilation channel m_A^2 must be positive in order that \varkappa_λ be negative. Additionally, the energy fraction y is restricted to the interval

$$\frac{1}{2} \left(1 - \sqrt{1 - \frac{4m^2}{|m_A^2|}} \right) < y < \frac{1}{2} \left(1 + \sqrt{1 - \frac{4m^2}{|m_A^2|}} \right). \quad (35)$$

Clearly, the radiation is possible only if $|m_A| > 2m$.

In the limit $|m_A| \gg m$, (33) simplifies

$$\omega \frac{d\Gamma_A}{d^3k} = N_c \frac{e^2 Q^2}{16(2\pi)^4} |m_A^2| \int_0^1 dy f(y\omega) f((1-y)\omega) (2y^2 - 2y + 1), \quad (36)$$

where only the photon polarization that gives $m_A^2 > 0$ contributes. The integral can be taken exactly:

$$\begin{aligned} & \int_0^1 dy f(y\omega) f((1-y)\omega) (2y^2 - 2y + 1) \\ &= \frac{1}{e^{\beta\omega} - 1} \left[-\frac{8\text{Li}_3(-e^{\beta\omega}) + 6\zeta(3)}{(\beta\omega)^3} + \frac{4\text{Li}_2(-e^{\beta\omega}) - \pi^2/3}{(\beta\omega)^2} + \frac{2\ln(1 + e^{\beta\omega}) - \ln 4}{\beta\omega} - \frac{2}{3} \right]. \end{aligned} \quad (37)$$

At low and high photon energy the spectrum reads

$$\omega \frac{d\Gamma_A}{d^3k} = N_c \frac{e^2 Q^2}{16(2\pi)^4} |m_A^2| \begin{cases} \frac{1}{6}, & \omega \ll T \\ \frac{2}{3} e^{-\beta\omega}, & \omega \gg T. \end{cases} \quad (38)$$

Comparing with (26) one can see that the decay channel dominates the low energy part of the spectrum, whereas the annihilation channel dominates the high energy tail, see Fig. 1. It is remarkable that since the photon polarization in the two channels is opposite, the total photon spectrum has different polarization direction at low and high energies with respect to T .




FIG. 1. Differential photon emission rate (solid line) and its two contributions from the decay (dashed line) and annihilation (dotted line) processes. Plasma temperature $T = 400$ MeV, chiral conductivity $\sigma_\chi = 1$ MeV and $\sum_f Q_f^2 = 5/9$ (for the two lightest flavors).

V. DISCUSSION AND SUMMARY

The main result of this paper are Eqs. (20) and (33) that represent the differential rates of photon emission rate by means of the chiral Cherenkov radiation in the decay and annihilation channels. Their sum gives the total photon emission rate (per unit volume). The magnitude of this contribution to the total photon yield by QGP shown in Fig. 1 is comparable with the conventional contributions as one can glean from Fig. 3 of [17].

An important phenomenological question is the value of the photon emission threshold at a given QGP temperature. Electromagnetic radiation by means of the mechanism described in this paper is possible if $\omega_{\text{pl}} < |m_A|$. The plasma frequency (8) of QGP at temperature $T = 400$ MeV is $\omega_{\text{pl}} \approx 35$ MeV. The chiral conductivity is unknown, but is often estimated to be of the order of a fraction of MeV. Importantly, it rapidly increases as T^4 . Thus, for example, if $\sigma_\chi = 1 - 10$ MeV then using the first of the equations (10), the infrared photon emission threshold is $\omega_0 \sim 0.1 - 1$ GeV. This is certainly within the range of phenomenologically interesting photon energies. A more precise knowledge of σ_χ may be extracted from the measurements of the charge separation effect in relativistic heavy-ion collisions because this effect is generated by the anomalous electric current proportional to σ_χ [34].

The calculations performed in this paper relied on the high energy approximation in which $\omega \gg m$. Since the quark thermal mass m is of the order of a hundred MeV, this approximation is not sufficiently reliable for the phenomenological applications to the QGP at realistic temperatures. Still, considering this limit has a great advantage of emphasizing the physics mechanism of photon radiation with the least mathematical and numerical complications possible. A comprehensive phenomenological approach would of course require going beyond the high-energy approximation.

ACKNOWLEDGMENTS

This work was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-87ER40371.

-
- [1] A. Adare *et al.* [PHENIX Collaboration], “Enhanced production of direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV and implications for the initial temperature,” *Phys. Rev. Lett.* **104**, 132301 (2010)
 - [2] J. Adam *et al.* [ALICE Collaboration], “Direct photon production in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Lett. B* **754**, 235 (2016)
 - [3] R. Baier, H. Nakkagawa, A. Niegawa and K. Redlich, “Production rate of hard thermal photons and screening of quark mass singularity,” *Z. Phys. C* **53**, 433 (1992).
 - [4] J. I. Kapusta, P. Lichard and D. Seibert, “High-energy photons from quark - gluon plasma versus hot hadronic gas,” *Phys. Rev. D* **44**, 2774 (1991) Erratum: [*Phys. Rev. D* **47**, 4171 (1993)].
 - [5] C. M. Hung and E. V. Shuryak, “Dilepton/photon production in heavy ion collisions, and the QCD phase transition,” *Phys. Rev. C* **56**, 453 (1997)
 - [6] J. V. Steele, H. Yamagishi and I. Zahed, “Dilepton and photon emission rates from a hadronic gas,” *Phys. Lett. B* **384**, 255 (1996)
 - [7] J. V. Steele, H. Yamagishi and I. Zahed, “Dilepton and photon emission rates from a hadronic gas. 2.,” *Phys. Rev. D* **56**, 5605 (1997)
 - [8] K. Dusling and I. Zahed, “Thermal photons from heavy ion collisions: A spectral function approach,” *Phys. Rev. C* **82**, 054909 (2010)
 - [9] C. H. Lee, J. Wirstam, I. Zahed and T. H. Hansson, “Thermal dileptons from a nonperturbative quark - gluon phase,” *Phys. Lett. B* **448**, 168 (1999)
 - [10] P. Aurenche, F. Gelis and H. Zaraket, “Landau-Pomeranchuk-Migdal effect in thermal field theory,” *Phys. Rev. D* **62**, 096012 (2000)
 - [11] P. B. Arnold, G. D. Moore and L. G. Yaffe, “Photon emission from quark gluon plasma: Complete leading order results,” *JHEP* **0112**, 009 (2001)

- [12] T. Peitzmann and M. H. Thoma, “Direct photons from relativistic heavy ion collisions,” *Phys. Rept.* **364**, 175 (2002)
- [13] S. Turbide, R. Rapp and C. Gale, “Hadronic production of thermal photons,” *Phys. Rev. C* **69**, 014903 (2004)
- [14] E. L. Bratkovskaya, S. M. Kiselev and G. B. Sharkov, “Direct photon production from hadronic sources in high-energy heavy-ion collisions,” *Phys. Rev. C* **78**, 034905 (2008)
- [15] I. Vitev and B. W. Zhang, “A Systematic study of direct photon production in heavy ion collisions,” *Phys. Lett. B* **669**, 337 (2008)
- [16] H. van Hees, C. Gale and R. Rapp, “Thermal Photons and Collective Flow at the Relativistic Heavy-Ion Collider,” *Phys. Rev. C* **84**, 054906 (2011)
- [17] J. F. Paquet, C. Shen, G. S. Denicol, M. Luzum, B. Schenke, S. Jeon and C. Gale, “Production of photons in relativistic heavy-ion collisions,” *Phys. Rev. C* **93**, no. 4, 044906 (2016)
- [18] O. Linnyk, V. Konchakovski, T. Steinert, W. Cassing and E. L. Bratkovskaya, “Hadronic and partonic sources of direct photons in relativistic heavy-ion collisions,” *Phys. Rev. C* **92**, no. 5, 054914 (2015)
- [19] M. Chiu, T. K. Hemmick, V. Khachatryan, A. Leonidov, J. Liao and L. McLerran, “Production of Photons and Dileptons in the Glasma,” *Nucl. Phys. A* **900**, 16 (2013)
- [20] L. McLerran and B. Schenke, “The Glasma, Photons and the Implications of Anisotropy,” *Nucl. Phys. A* **929**, 71 (2014)
- [21] K. Tuchin, “Electromagnetic radiation by quark-gluon plasma in a magnetic field,” *Phys. Rev. C* **87**, no. 2, 024912 (2013)
- [22] H. U. Yee, “Flows and polarization of early photons with magnetic field at strong coupling,” *Phys. Rev. D* **88**, no. 2, 026001 (2013)
- [23] G. Basar, D. Kharzeev, D. Kharzeev and V. Skokov, “Conformal anomaly as a source of soft photons in heavy ion collisions,” *Phys. Rev. Lett.* **109**, 202303 (2012)
- [24] K. Fukushima and K. Mameda, “Wess-Zumino-Witten action and photons from the Chiral Magnetic Effect,” *Phys. Rev. D* **86**, 071501 (2012)
- [25] K. A. Mamo and H. U. Yee, “Spin polarized photons and dileptons from axially charged plasma,” *Phys. Rev. D* **88**, no. 11, 114029 (2013)
- [26] K. A. Mamo and H. U. Yee, “Spin polarized photons from an axially charged plasma at weak coupling: Complete leading order,” *Phys. Rev. D* **93**, no. 6, 065053 (2016)
- [27] K. Fujikawa and H. Suzuki, “Path integrals and quantum anomalies,” Oxford, UK: Clarendon (2004) 284 p
- [28] S. M. Carroll, G. B. Field and R. Jackiw, “Limits on a Lorentz and Parity Violating Modification of Electrodynamics,” *Phys. Rev. D* **41**, 1231 (1990).
- [29] R. Lehnert and R. Potting, “Vacuum Cherenkov radiation,” *Phys. Rev. Lett.* **93**, 110402 (2004)
- [30] R. Lehnert and R. Potting, “The Cerenkov effect in Lorentz-violating vacua,” *Phys. Rev. D* **70**, 125010 (2004) Erratum: [*Phys. Rev. D* **70**, 129906 (2004)]

- [31] K. Tuchin, “Radiative instability of quantum electrodynamics in chiral matter,” *Phys. Lett. B* **786**, 249 (2018),
- [32] X. G. Huang and K. Tuchin, “Transition Radiation as a Probe of the Chiral Anomaly,” *Phys. Rev. Lett.* **121**, no. 18, 182301 (2018),
- [33] D. Kharzeev, “Parity violation in hot QCD: Why it can happen, and how to look for it,” *Phys. Lett. B* **633**, 260 (2006)
- [34] D. Kharzeev and A. Zhitnitsky, “Charge separation induced by P-odd bubbles in QCD matter,” *Nucl. Phys. A* **797** (2007) 67
- [35] K. Fukushima, D. E. Kharzeev and H. J. Warringa, “The Chiral Magnetic Effect,” *Phys. Rev. D* **78** (2008) 074033
- [36] D. E. Kharzeev, “Topologically induced local P and CP violation in QCD \times QED,” *Annals Phys.* **325**, 205 (2010)
- [37] A. R. Zhitnitsky, “The topological long range order in QCD. Applications to heavy ion collisions and cosmology,” arXiv:1411.2606 [hep-ph].
- [38] A. R. Zhitnitsky, “Topological order and Berry connection for the Maxwell Vacuum on a four-torus,” *Phys. Rev. D* **90**, no. 10, 105007 (2014)
- [39] K. Tuchin, “Chiral Cherenkov and chiral transition radiation in anisotropic matter,” *Phys. Rev. D* **98**, no. 11, 114026 (2018)
- [40] F. Wilczek, “Two Applications of Axion Electrodynamics,” *Phys. Rev. Lett.* **58**, 1799 (1987).
- [41] P. Sikivie, “On the Interaction of Magnetic Monopoles With Axionic Domain Walls,” *Phys. Lett. B* **137**, 353 (1984).
- [42] T. Kalaydzhyan, “Chiral superfluidity of the quark-gluon plasma,” *Nucl. Phys. A* **913**, 243 (2013)
- [43] D. E. Kharzeev and H. J. Warringa, “Chiral Magnetic conductivity,” *Phys. Rev. D* **80**, 034028 (2009)
- [44] D. E. Kharzeev, “The Chiral Magnetic Effect and Anomaly-Induced Transport,” *Prog. Part. Nucl. Phys.* **75**, 133 (2014)
- [45] V. A. Rubakov and M. E. Shaposhnikov, “Electroweak baryon number nonconservation in the early universe and in high-energy collisions,” *Usp. Fiz. Nauk* **166**, 493 (1996) [*Phys. Usp.* **39**, 461 (1996)]
- [46] G. D. Moore, C. r. Hu and B. Muller, “Chern-Simons number diffusion with hard thermal loops,” *Phys. Rev. D* **58**, 045001 (1998)
- [47] D. Bodeker, G. D. Moore and K. Rummukainen, “Chern-Simons number diffusion and hard thermal loops on the lattice,” *Phys. Rev. D* **61**, 056003 (2000)
- [48] G. D. Moore and M. Tassler, “The Sphaleron Rate in SU(N) Gauge Theory,” *JHEP* **1102**, 105 (2011)
- [49] D. T. Son and A. O. Starinets, “Minkowski space correlators in AdS/CFT correspondence: Recipe and applications,” *JHEP* **0209**, 042 (2002)
- [50] Y. Akamatsu and N. Yamamoto, “Chiral Plasma Instabilities,” *Phys. Rev. Lett.* **111**, 052002 (2013),