

12-2014

Case 3664. *Tipula contaminata* Linnaeus, 1758 (currently *Ptychoptera contaminata*; Insecta, Diptera): proposed conservation of prevailing usage through designation of a neotype

Andrew Fasbender
Iowa State University

Gregory W. Courtney
Iowa State University, gwcourt@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/ent_pubs



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/ent_pubs/569. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Entomology at Iowa State University Digital Repository. It has been accepted for inclusion in Entomology Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Case 3664. *Tipula contaminata* Linnaeus, 1758 (currently *Ptychoptera contaminata*; Insecta, Diptera): proposed conservation of prevailing usage through designation of a neotype

Abstract

The purpose of this application, under Article 75.6 of the Code, is to conserve the universal usage of *Tipula contaminata* Linnaeus, 1758 by setting aside all previous type fixations and designating a neotype. *Tipula contaminata* is the type species of the genus *Ptychoptera* Meigen, 1803, itself the type genus of the family PTYCHOPTERIDAE Osten Sacken, 1862. This species is found over much of Europe, and all authors subsequent to Meigen (1803) have utilized his concept of the species. However, the holotype of *Tipula contaminata* Linnaeus, 1758 represents a species of TIPULIDAE. It is proposed that a neotype be designated for *Tipula contaminata* to preserve two hundred years of common usage and ensure nomenclatural stability at the genus and family rank.

Keywords

Nomenclature, taxonomy, Insecta, Diptera, PTYCHOPTERIDAE, *Ptychoptera*, *Tipula*, *Ptychoptera contaminata*, Phantom Crane Fly, Palearctic, Europe

Disciplines

Ecology and Evolutionary Biology | Entomology

Comments

This article is published as Andrew Fasbender and Gregory W. Courtney, "Case 3664 *Tipula contaminata* Linnaeus, 1758 (currently *Ptychoptera contaminata*; Insecta, Diptera): proposed conservation of prevailing usage through designation of a neotype," *The Bulletin of Zoological Nomenclature* 71(4), 237-243, (1 December 2014). <https://doi.org/10.21805/bzn.v71i4.a14>. Posted with permission.

Case 3664***Tipula contaminata* Linnaeus, 1758 (currently *Ptychoptera contaminata*; Insecta, Diptera): proposed conservation of prevailing usage through designation of a neotype**

Andrew Fasnender & Gregory W. Courtney

Department of Entomology, Iowa State University, 3222 Science II, Ames, Iowa 50011–3222, United States of America

(e-mails: afsbndr@iastate.edu, gwcourt@iastate.edu)

Abstract. The purpose of this application, under Article 75.6 of the Code, is to conserve the universal usage of *Tipula contaminata* Linnaeus, 1758 by setting aside all previous type fixations and designating a neotype. *Tipula contaminata* is the type species of the genus *Ptychoptera* Meigen, 1803, itself the type genus of the family PTYCHOPTERIDAE Osten Sacken, 1862. This species is found over much of Europe, and all authors subsequent to Meigen (1803) have utilized his concept of the species. However, the holotype of *Tipula contaminata* Linnaeus, 1758 represents a species of TIPULIDAE. It is proposed that a neotype be designated for *Tipula contaminata* to preserve two hundred years of common usage and ensure nomenclatural stability at the genus and family rank.

Keywords. Nomenclature; taxonomy; Insecta; Diptera; PTYCHOPTERIDAE; *Ptychoptera*; *Tipula*; *Ptychoptera contaminata*; Phantom Crane Fly; Palaearctic; Europe.

1. Linnaeus (1758, p. 586) described the species *Tipula contaminata*. The only available syntype (Linnaeus did not designate a holotype) is a member of the family TIPULIDAE Latreille, 1802. The specimen is an adult male mounted by a pin through the dorsal surface of the thorax. The complete head and abdomen remain attached to the thorax, as does one wing. All legs are detached, with only one leg glued to the determination label, on which is written '*T. contaminata*' in cursive script. No other labels are present. The material is housed at the Linnean Society of London, and the authors examined photographic images of the specimen. In his description of this species, Linnaeus (1758) referred to entry 1134 in his *Fauna Svecica* (Linnaeus, 1746, p. 333), but did not indicate in either of these works how many specimens he examined. The current syntype does not match Linnaeus's (1746) description or (1758) diagnosis, 'alis nigro maculatis, corpore nigro,' [wing maculated black, body black]. The syntype thorax has the scutum and pleurites russet brown with dark patterning. It is possible that this specimen was part of a larger series assigned to *Tipula contaminata* and this is the only specimen still existing. It is currently unknown who designated the syntype. The examination of this specimen was undertaken by the first author based on photomicrographs provided by the Linnean Society of London. The specimen is in good diagnostic condition, retaining the male genitalia and is thus distinguishable to species level with dissection, though the authors feel that a tipulid specialist should undertake this examination.



Fig. 1. Habitus of the proposed neotype of *Tipula contaminata* Linnaeus, 1758. Scale bar 1mm.

2. Fabricius (1775, pp. 749–750; 1781, p. 402) provided a diagnosis of *Tipula contaminata*, correctly referencing Linnaeus as the originator of the name. The species was also diagnosed by Fabricius (1787, p. 322), but no citation to Linnaeus was provided.

3. Meigen (1800) proposed several dozen new genera of Diptera with the description of the genus *Liriope* Meigen, 1800 closely corresponding to his later (Meigen, 1803) description of the genus *Ptychoptera*. Meigen's (1800) publication was suppressed in 1960 by the Commission in Opinion 678 (BZN 20: 339–342, 1963).

4. Meigen (1803, pp. 262–263), listed '*Tipula contaminata* Fabr.' and *Tipula albimana* Fabricius, 1787 as members of the genus *Ptychoptera* Meigen, 1803. '*Tipula contaminata* Fabr.' clearly refers to *Tipula contaminata* Linnaeus, 1758 as presented



Fig. 2. Locality label of the proposed neotype of *Tipula contaminata*. Scale bar 1mm.

in Fabricius's works. Meigen's generic description (diagnosis based on the elongate first antennal flagellomere/third antennal segment) does not fit the holotype of *Tipula contaminata* Linnaeus, 1758. Specifically, 'PTYCHOPTERA. Die Fühlhörner vorgestreckt, sechszehngliedrig: das erste Glied walzenförmig, kurz; das zweite becherförmig; das dritte walzenförmig, lang; die folgenden länglicht, dünnhaarig Die Flügel halb offen.' [PTYCHOPTERA. Antennae stretched forward, sixteen segmented: the first segment cylindrical, short; the second cup-shaped; the third cylindrical, long; the following oblong, fine setae. The wings half open.] In the Linnaean type specimen the first flagellomere/third antennal segment is round, and not significantly longer than the succeeding flagellomeres.

5. Latreille (1810, p. 442) subsequently designated 'Ptychoptera contaminata, Fab.' as the type species of *Ptychoptera* as a First Reviser action (Latreille). According to Article 67.7 the incorrect citation of '*Ptychoptera contaminata* Fabr.' is considered to refer to *Tipula contaminata* Linnaeus, 1758, and is a valid designation. Based on Article 69.1 and the Opinions 11 (Smithsonian Miscellaneous Publications 1938: 17–18, 1910) and 136 (Opinions and Declarations 2: 13–19, August 1939) Latreille's (1810) citation is to be accepted as a designation of *Tipula contaminata* Linnaeus, 1758 as the type of the genus.

6. Osten Sacken (1862, p. 12) proposed the family-group taxon PTYCHOPTERINA (later emended to PTYCHOPTERINAE by Schiner, 1863) within TIPULIDAE in 1862. The genera *Bittacomorpha* Westwood, 1835, *Macrochile* Loew, 1850, *Protoplasa* Osten Sacken, 1860 and *Ptychoptera*, were included; the type genus to be inferred is *Ptychoptera*.

7. Brauer (1869) and Hart (1895, pp. 189–190) considered PTYCHOPTERIDAE as a full family.

8. Meigen's concept of *Tipula contaminata* has been universally accepted (Handlirsch, 1909, p. 269, 271–272; Tonnoir, 1919, pp. 115, 119; Grünberg, 1920, pp. 76–77; Reidel, 1921, p. 147; Séguy, 1925, pp. 8–9, 11–14; Audcent, 1934, pp. 106–109, 111, 116–119; Peus, 1958, pp. 11, 15–21, 26, 28, 30, 32, 34; Brindle, 1962, p. 212–216; Tjeder, 1968, pp. 73–75; Zitek-Zwyrtek, 1971, pp. 416, 418, 420–423; Zwyrtek, 1971, pp. 36–38; Hansen, 1981, pp. 59–63; Theischinger, 1978, pp. 26; Draskovits, 1983, pp. 80, 82–83, 85; Krzemiński, 1986, pp. 105, 107–108, 117–119; Zwick, 1988, pp. 123, 128–129; Podenas, 1991, p. 155; Krzemiński & Zwick, 1993, pp. 80, 85–86; Stubbs,




Fig. 3. Male genitalia of the proposed neotype of *Tipula contaminata*: **a.** Overall lateral view; **b.** Epandrium lateral view; **c.** Epandrium dorsal view; **d.** Aedeagus anterior view; **e.** Aedeagus lateral view; **f.** Hypandrium posterior view; **h.** Gonopod dorsal view; **i.** Gonopod dorsal view; **j.** Paramere posterior view; **k.** Paramere dorsal view. Scale bars **a–i** 0.5 mm, **j & k** 0.25 mm.

1993, pp. 7, 9, 12, 25, 28; Rozkošný, 1997, pp. 294–295; Pârvu, 2004, p. 190; Ujvárosi et al., 2011, pp. 40, 42–43, etc.). This species (*Ptychoptera* sp.) has black thoracic

sclerites as well as wings with maculated infuscation, matching the description by Linnaeus (1746, 1758).

9. The authors cannot find a single instance of the usage of *Ptychoptera contaminata* after 1800 which is definitively referable to the species represented by the Linnaean type material, and the descriptions provided by Linnaeus and Fabricius do not match the coloration of the syntype.

10. The consequences of designating the single surviving syntype as a lectotype would be placing *Ptychoptera* as a synonym of an undetermined tipulid genus and PTYCHOPTERIDAE as a subjective junior synonym of TIPULIDAE. The current concept of the taxa *Ptychoptera* and PTYCHOPTERIDAE would require new names, and *Ptychoptera* sp. would be assigned a different name than that to which it has historically been referred. There are no accepted junior synonyms of *Ptychoptera* sp., and Peus (1958 p. 40) did not refer any of the nomina dubia within the genus to *Ptychoptera contaminata*.

11. The authors propose that the Commission, under Article 75.6, set aside all previous type fixations (including the name-bearing specimen of *Ptychoptera contaminata* in the Linnean Society of London, and designate a specimen of the currently unnamed species from the collection of the Natural History Museum of Denmark as the neotype. The neotype was chosen because it is representative of the prevailing usage of the name *Ptychoptera contaminata*, for the state of preservation of the specimen and the locality in northern Europe where Linnaeus' specimen was probably collected. This neotype designation would preserve over two centuries of common usage of *Ptychoptera contaminata*, and preserve the established usage of the genus- and family-rank names *Ptychoptera* Meigen, 1803 and PTYCHOPTERIDAE Osten Sacken, 1862.

12. The authors do not anticipate any opposition among ptychopterid or tipulid taxonomists, as this measure is essentially conservative and avoids any nomenclatural changes in either taxon.

13. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary power to set aside all type fixations for the nominal species *contaminata* Linnaeus, 1758, as published in the binomen *Tipula contaminata*, and to designate the specimen from the Natural History Museum of Denmark (ZMUC) with the locality label data 'Dania, Lolland, Saxkøbing Sómose 20–6-1966 Bo Vest Pedersen,' and a second label reading 'NEOTYPE: *Tipula contaminata* Linnaeus 1758, det. A Fasbender' as the neotype;
- (2) to place on the Official List of Specific Names in Zoology the name *contaminata* Linnaeus 1758, as published in the binomen *Tipula contaminata* and as defined by the neotype designated in (1) above.

Acknowledgements

F. Christian Thompson provided valuable comments and advice at the outset of this work. The Linnean Society of London provided access to images of the type specimen of Linnean *Tipula contaminata*. Wayne Mathis (*United States National Museum of Natural History*), Thomas Pape (*Natural History Museum of Denmark*), and Jennifer Thomas (*University of Kansas Snow Entomological Museum*) facilitated loans of specimens which were used in this work. Three anonymous Commission members are thanked for suggestions that improved this manuscript. This project

was supported in part by National Science Foundation grants (DEB-0933218 and EF-1115112) to G.W. Courtney and the National Institute of Food and Agriculture, Project No. 6693.

References

- Alexander, C.P.** 1919. The crane-flies of New York. Part I. Distribution and taxonomy of the adult flies. *Cornell University Agricultural Experimental Station Memoirs*, **25**: 767–1133.
- Audcent, H.** 1934. British Liriopeidae (Diptera, Nematocera). *Transactions of the Society for British Entomology*, **1**(2): 103–123.
- Brauer, F.** 1869. Kurze Charakteristik der Dipteren-Larven zur Bekräftigung des neuer von Dr. Schiner entworfenen Dipteren-Systems. *Abhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, **19**: 843–852.
- Brindle, A.** 1962. Taxonomic notes on the larvae of British Diptera: 9. The family Ptychopteridae. *The Entomologist*, **95**: 212–216.
- Draskovits, A.** 1983. 11. család. Ptychopteridae – Redős Szúnyogok. *Fauna Hungariae*, **14**: 79–88.
- Evenhuis, N.L.** 1997. *Litteratura taxonomica dipteorum (1758–1930)*, vol. 2, 427–871 pp. Backhuys Publishers, Leiden.
- Fabricius, J.C.** 1775. *systema entomologiae sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. 832 pp. Kortii, Flensburgi & Lipsiae.
- Fabricius, J.C.** 1781. *species insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca Natalia, metamorphosin, adiectis observationibus, descriptionibus*, vol. 2. 494 pp. E. Bohnii, Hamburgi & Kilonii.
- Fabricius, J.C.** 1787. *mantissa insectorum sistens nuper detectas adiectis synonymis, observationibus, descriptionibus, emendationibus*, vol. 2. 382 pp. Proft, Halfnia.
- Grünberg, K.** 1920. Diptera. In Brauer, A. (Ed.), *Die Süßwasserfauna Deutschlands*, vol. 2a. Fischer, Jena.
- Handlirsch, A.** 1909. Zur Phylogenie und Flügelmorphologie der Ptychopteriden (Dipteren). *Annalen des K.K. Naturhistorischen Hofmuseums*, **23**: 263–272.
- Hansen, S.B.** 1981. Bestemmelsesnøgle til larver af danske Ptychopteridae (Diptera: Nematocera), med noter om arternes habitatpræferenser. *Entomologiske Meddelelser*, **49**: 59–64.
- Hart, C.A.** 1895. On the entomology of the Illinois River and adjacent waters – First paper. *Bulletin of the Illinois State Laboratory of Natural History*, **4**: 149–273.
- Krzemiński, W.** 1986. Ptychopteridae of Poland (Diptera: Nematocera). *Polskie Pismo Entomologiczne*, **56**: 105–131.
- Krzemiński, W. & Zwick, P.** 1993. New and little known Ptychopteridae (Diptera) from the Palaearctic region. *Aquatic Insects*, **15**(2): 65–87.
- Latreille, P.A.** 1810. *Considérations générales sur l'ordre naturel des animaux composant les classes des Crustacés, des Arachnides et des Insectes avec un tableau méthodique de leurs genres disposés en familles*. 444 pp. Schoell, Paris.
- Linnaeus, C.** 1746. *Fauna Suecica sistens animalia suecica regni: quadrupedia, aves, amphibian, pisces, insecta, vermes, distributa per classes & ordines, genera & species*. Pp. 1–28, 1–411, 1 pl. Wishoff, Lugduni Batavorum (Leiden).
- Linnaeus, C.** 1758. *Systema Naturae*, Ed. 10, vol. 1. 824 pp. Salvii, Holmiae.
- Meigen, J.W.** 1800. *Nouvelle Classification des Mouches à Deux Ailes*. 40 pp. Fuchs & Perroneau, Paris.
- Meigen, J.W.** 1803. Versuch einer neuen Gattungs-Eintheilung der europäischen zweiflügeligen Insekten. *Magazin für Insektenkunde (Illiger)*, **2**: 259–281.
- Osten Sacken, C.R.** 1862. Fam. VI. Tipulidae. In Loew, H. *Monographs of the Diptera of North America*, Part I. *Smithsonian Miscellaneous Collections*, **6**: 1–221.
- Pârvu, C.** 2004. Nematocerans (Diptera: Nematocera) from Piatra Craiului National Park (Romania). *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, **46**: 181–192.

- Peus, F.** 1958. Liriopidae. In Lindner, E. (Ed.), *Die Fliegen der Palaearktischen Region*, vol. 10. 44. pp Schweizerbart'sche Verlagsbuchhandlung, Stuttgart.
- Podenas, S.** 1992. Tipulidae, Limoniidae, Cylindrotomidae, Trichoceridae and Ptychopteridae of Lithuania. *Acta Zoologica Cracoviensia*, **35**: 147–156.
- Reidel, M.P.** 1921. Altes und Neues von den Faltenmücken. *Entomologisches Jahrbuch*, **30**: 144–148.
- Rozkošný, R.** 1997. Family Ptychopteridae. Pp. 291–297 in Papp, L., Darvas, B. (Eds.), *Contributions to a manual of Palaearctic Diptera*, vol. 2, Science Herald, Budapest.
- Schiner, I.R.** 1863. *Fauna Austriaca. Die Fliegen (Diptera). Nach der analytischen Methode bearbeitet*, vol. 2, Heft 11/13 "1864". C. Gerold's Sohn, Wien [publication date after Evenhuis, 1997].
- Séguy, E.** 1925. Diptères (Nématocères piqueurs): Ptychopteridae, Orphnephilidae, Simuliidae, Culicidae, Psychodidae Phlebotominae. *Faune de France*, **12**: 1–109.
- Stubbs, A.E.** 1993. *Provisional atlas of the ptychopterid craneflies (Diptera: Ptychopteridae) of Britain and Ireland*. Huntingdon: Biological Records Centre. Dixon Printing, United Kingdom.
- Theischinger, G.** 1978. Faltenmücken aus Oberösterreich. *Naturkunde Jahrbuch Stadt Linz*, **23**: 25–28.
- Tjeder, B.** 1986. Notes on the Scandinavian Ptychopteridae, with description of a new species (Diptera). *Opuscula Entomologica*, **33**: 73–79.
- Tonnair, A.** 1919. Notes sur les Ptychopteridae (Dipt.). *Annales Societe Entomologique de Belgique*, **59**: 115–122.
- Ujvárosi, L., Kolcsár, P.L. & Török, E.** 2011. An annotated list of Ptychopteridae (Insecta, Diptera) from Romania, with notes on the individual variability of *Ptychoptera albimana* (Fabricius, 1787). *Entomologica Romanica*, **16**: 39–45.
- Zitek-Zwyrtek, K.** 1971. Czechoslovak species of the family Ptychopteridae (Diptera). *Acta Entomologica Bohemoslovaca*, **68**: 416–426.
- Zwick, P.** 1988. Contribution to the Turkish Blephariceridae and Ptychopteridae (Diptera). *Mitteilungen der Schweizerische Entomologische Gesellschaft* **61**: 123–129.
- Zwyrtek, K.** 1971. Revize slídilkovitých (Ptychopteridae, Diptera) ze sbírek Národního Muzea v Praze. *Casopis Národního Muzea*, **140**: 36–38.

Acknowledgement of receipt of this application was published in BZN 71: 145.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the I.C.Z.N., Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).