

6-2013

Using biodiversity to link agricultural productivity with environmental quality: Results from three field experiments in Iowa

Matthew Z. Liebman
Iowa State University, mliebman@iastate.edu

Matthew J. Helmers
Iowa State University, mhelmers@iastate.edu

Lisa A. Schulte-Moore
Iowa State University, lschulte@iastate.edu

Craig A. Chase
Iowa State University, cchase@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/abe_eng_pubs

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Bioresource and Agricultural Engineering Commons](#), [Natural Resources Management and Policy Commons](#), and the [Sustainability Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/abe_eng_pubs/373. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

Using biodiversity to link agricultural productivity with environmental quality: Results from three field experiments in Iowa

Abstract

Agriculture in the US Corn Belt is under increasing pressure to produce greater quantities of food, feed and fuel, while better protecting environmental quality. Key environmental problems in this region include water contamination by nutrients and herbicides emitted from cropland, a lack of non-agricultural habitat to support diverse communities of native plants and animals, and a high level of dependence on petrochemical energy in the dominant cropping systems. In addition, projected changes in climate for this region, which include increases in the proportion of precipitation coming from extreme events could make soil and water conservation in existing cropping systems more difficult. To address these challenges we have conducted three cropping systems projects in central Iowa: the Marsden Farm Cropping Systems experiment, the Science-based Trials of Row-crops Integrated with Prairies (STRIPs) experiment, and the Comparison of Biofuel Systems (COBS) experiment. Results from these experiments indicate that (1) diversification of the dominant corn–soybean rotation with small grains and forage legumes can permit substantial reductions in agrichemical and fossil hydrocarbon use without compromising yields or profitability; (2) conversion of small amounts of cropland to prairie buffer strips can provide disproportionately large improvements in soil and water conservation, nutrient retention, and densities of native plants and birds; and (3) native perennial species can generate large amounts of biofuel feedstocks and offer environmental benefits relative to corn- and soybean-based systems, including greater carbon inputs to soil and large reductions in nitrogen emissions to drainage water. Increasing biodiversity through the strategic integration of perennial plant species can be a viable strategy for reducing reliance on purchased inputs and for increasing agroecosystem health and resilience in the US Corn Belt.

Keywords

biodiversity, multifunctionality, conservation, cropping systems, Agronomy, Natural Resource Ecology and Management, Leopold Center

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Bioresource and Agricultural Engineering | Natural Resources Management and Policy | Sustainability

Comments

This article is from *Renewable Agriculture and Food Systems* 28, no. 2 (June 2013): 115–128, doi:[10.1017/S1742170512000300](https://doi.org/10.1017/S1742170512000300).

Using biodiversity to link agricultural productivity with environmental quality: Results from three field experiments in Iowa

Matt Liebman^{1,*}, Matthew J. Helmers², Lisa A. Schulte³ and Craig A. Chase⁴

¹Department of Agronomy, Iowa State University, Ames, IA 50011-1010, USA.

²Department of Agricultural and Biosystems Engineering, Iowa State University, Ames, IA 50011-3080, USA.

³Department of Natural Resource Ecology and Management, Iowa State University, Ames, IA 50011-3221, USA.

⁴Leopold Center for Sustainable Agriculture, Iowa State University, Ames, IA 50010-1050, USA.

*Corresponding author: mliebman@iastate.edu

Accepted 20 April 2012; First published online 11 March 2013

Research Paper

Abstract

Agriculture in the US Corn Belt is under increasing pressure to produce greater quantities of food, feed and fuel, while better protecting environmental quality. Key environmental problems in this region include water contamination by nutrients and herbicides emitted from cropland, a lack of non-agricultural habitat to support diverse communities of native plants and animals, and a high level of dependence on petrochemical energy in the dominant cropping systems. In addition, projected changes in climate for this region, which include increases in the proportion of precipitation coming from extreme events could make soil and water conservation in existing cropping systems more difficult. To address these challenges we have conducted three cropping systems projects in central Iowa: the Marsden Farm Cropping Systems experiment, the Science-based Trials of Row-crops Integrated with Prairies (STRIPs) experiment, and the Comparison of Biofuel Systems (COBS) experiment. Results from these experiments indicate that (1) diversification of the dominant corn–soybean rotation with small grains and forage legumes can permit substantial reductions in agrichemical and fossil hydrocarbon use without compromising yields or profitability; (2) conversion of small amounts of cropland to prairie buffer strips can provide disproportionately large improvements in soil and water conservation, nutrient retention, and densities of native plants and birds; and (3) native perennial species can generate large amounts of biofuel feedstocks and offer environmental benefits relative to corn- and soybean-based systems, including greater carbon inputs to soil and large reductions in nitrogen emissions to drainage water. Increasing biodiversity through the strategic integration of perennial plant species can be a viable strategy for reducing reliance on purchased inputs and for increasing agroecosystem health and resilience in the US Corn Belt.

Key words: biodiversity, multifunctionality, conservation, cropping systems

Introduction

The US Corn Belt, spanning from western Ohio to eastern Nebraska and from southern Minnesota to northern Missouri, comprises one of the largest, most productive rain-fed grain regions in the world. Its natural endowment with fertile soils and a favorable climate makes it a major focal point in discussions of how to supply an expanding world population with enough food and plant-derived renewable energy. These discussions increasingly focus not only on the technologies of farming but also on the environmental impacts of farming practices and the resilience of agricultural systems to changes in climate,

energy and nutrient availability, and other factors acting at a global scale.

Patterns of agricultural development, land use and environmental quality in Iowa exemplify the directions that conventional agriculture has followed in the Corn Belt during the past half-century. Iowa is now the leading US producer of corn, soybean, hogs, eggs and ethanol, and ranks seventh in cattle production^{1,2}. Concomitantly, the state also ranks high nationally in the number of surface waters impaired by excessive concentrations of nutrients, pathogens, pesticides and soil sediment, and its croplands are major sources of nitrogen and phosphorus delivered to the hypoxic zone in the Gulf of Mexico^{3–5}.

In addition to high levels of water pollution, the development of conventional agricultural systems in the Corn Belt has been accompanied by a major loss of biological diversity. Tall-grass prairie was the dominant land cover in Iowa and much of the central and western Corn Belt prior to Euro-American settlement⁶; smaller areas within the region were occupied by other types of perennial plant communities, including savannas, riparian forests and wetlands. Most of this native vegetation is now gone, having been converted to cropland or pasture by 1900. At present, Iowa ranks last nationally in the amount of pre-settlement vegetation still remaining⁷, with the state's prairie communities reduced to <0.1% of their former area⁸. Change has also taken place in recent years where land that had been in perennial cover through enrollment in the Conservation Reserve Program (CRP) has been converted to row crop production. Schilling and Spooner⁹ reported that in Iowa's Squaw Creek watershed, 9% of the area that had been in CRP grassland in 1990 was converted to corn and soybean production by 2005. Not surprisingly, given the loss of native plant communities and conservation grasslands, the diversity of Iowa's native flora and fauna has been greatly reduced¹⁰.

A substantial loss of crop diversity has also occurred throughout the central Corn Belt, illustrated by historical changes in land use in Iowa. Although the total amount of cropland in Iowa has remained relatively constant over the past 60 years, there has been a marked shift toward corn and soybean production; these two crops now occupy two-thirds of the state's total land area¹¹. Between 1949 and 1997 in Iowa's Raccoon River watershed, the percentage of cropland used for wheat, barley, oat, alfalfa and other hay crops fell from 42 to 3%; in contrast, the proportion in corn and soybean grew from 57 to 97%¹². Historical changes in crop diversity in the Corn Belt are linked with changes in livestock production in the region. From 1945 to 2000, cattle numbers in the Corn Belt declined by 52% as beef production shifted westward; concomitantly, land used for hay and oat production, used to feed cattle and other livestock, declined by 60 and 97%, respectively¹³. Similarly, in an analysis of landscape change during 1937–2002 in three Iowa townships, Brown and Schulte¹⁴ noted marked reductions in the area of grasslands suitable for pasturing cattle.

Reductions in crop and non-crop diversity have been linked to a variety of forms of environmental degradation and agroecosystem dysfunction; conversely, increases in diversity can be associated with improvements in agroecosystem health and function. For example, Hatfield *et al.*¹² reported that nitrate concentrations in Iowa's Raccoon River have increased since 1970 as land use in adjacent cropland has shifted from diverse cropping systems containing small grains, hay, corn and soybean to simpler corn–soybean systems, which are more prone to nutrient leaching. Reductions in diversity and

accompanying reductions in ecosystem functioning have also occurred due to removal of non-crop vegetation. In pursuit of increased field size and greater ease in the use of large agricultural machinery, trees on field borders have been removed in many areas of the Midwestern USA. Yet, fields that lack adjacent shelterbelt trees can be subject to increased wind speeds and greater wind erosion, and have reduced crop water use efficiency and lower yields^{15,16}. Non-crop vegetation can also affect pest dynamics. In a comparison of Midwestern US landscapes dominated by corn and soybean fields versus landscapes containing a mix of crop fields and non-crop habitat, Gardiner *et al.*¹⁷ found that soybean aphid (*Aphis glycines*) was subject to less biological control by natural enemies and consequently was more problematic in the less diverse landscapes, whereas greater diversity at a landscape level enhanced predator populations and biological control of the pest.

Given the loss in the Corn Belt of native biodiversity, crop diversity and the ecosystem services that diversity can provide, we ask: Are impoverished biological communities and environmental degradation unavoidable costs of high agricultural productivity or can farming systems be developed to provide both high yields and necessary ecosystem services, including water quality protection, soil conservation and pest control? Schulte *et al.*¹⁸ suggested that the strategic integration of perennial vegetation into agricultural landscapes dominated by annual crops could help reconcile agricultural productivity with environmental quality. Increasing crop and non-crop diversity in agroecosystems with perennial species may be especially important because perennial plants perform many key ecological functions more effectively than annuals, including (1) regulating the flow and storage of water, (2) reducing soil erosion, (3) storing and cycling nutrients and carbon, and (4) providing habitat for natural enemies of crop pests and for native plants and animals of conservation concern^{18,19}.

Many options exist for diversifying agroecosystems with perennial vegetation, including forage crops that are alternated with row crops in rotation sequences; riparian buffers of trees, shrubs and grasses that separate crop fields from streams; in-field strips of herbaceous species that filter runoff water; trees and shrubs on field margins that reduce wind movement; reconstructed wetlands that receive drainage water from crop fields; agroforestry plantations that supply a variety of economic products; and woody and herbaceous plants that are harvested as biofuel feedstocks¹⁸. Here, we describe some of the agronomic, economic and environmental consequences of three approaches for increasing diversity in agroecosystems in Iowa: adding perennial legumes to corn–soybean rotations; weaving strips of prairie vegetation into corn and soybean fields to serve as filter strips; and growing prairie vegetation to produce solid or liquid fuel.

Lengthening Rotations to Reduce Agrichemical Use While Maintaining Productivity: The Marsden Farm Cropping Systems Experiment

Conventional corn and soybean production is a chemically intensive activity. Together, corn and soybean receive more herbicide active ingredients and more synthetic fertilizer than any other crops grown in the USA^{20,21}. Emissions of herbicides and nitrogen from fields to water in the Corn Belt are roughly proportional to the quantities of those materials applied and the percentage of a watershed occupied by corn and soybean^{22–25}. In the case of herbicides, emissions include both runoff and leaching losses; for nitrogen, emissions result from runoff and leaching of fertilizer, as well as losses from mineralized soil organic matter and manure²⁶. Unlike deep-rooted perennial forage crops whose root systems are active for many months each year, corn and soybean have relatively shallow root systems that are active for only part of the year, rendering those crops leaky with regard to nitrogen loss during the spring and fall, when rainfall can be abundant and can transport nitrogen through the soil^{12,27}.

Concentrations of herbicides and nitrate in surface and ground water in the Corn Belt can exceed health-based standards for drinking water quality, raising concerns for impacts on human health, wildlife and fisheries^{3,28–30}. Magdoff *et al.*³¹ and Dinnes *et al.*³² noted that reductions in synthetic nitrogen fertilizer use, coupled with greater use of perennial legume crops and judicious use of manure, constitute important components of strategies to conserve nitrogen and protect water from nitrogen contamination. Similarly, reducing herbicide use by increasing reliance on physical weed control practices and diverse crop sequences that challenge weeds with a broad range of stress and mortality factors may be an effective approach for reducing herbicide concentrations in the hydrologic system^{3,33,34}.

To test the hypothesis that diversifying a simple corn–soybean cropping system could allow for substantial reductions in nitrogen fertilizer and herbicide use without compromising crop productivity and profitability, a 9-ha field experiment was established in 2001 at the Iowa State University Marsden Farm, in Boone Co., IA. The site lies within a region of intensive rain-fed corn and soybean production and is surrounded by farms with high levels of productivity. Soils at the site are fertile Mollisols. In addition to a conventionally managed 2-year rotation (corn–soybean) that receives fertilizers and herbicides at rates comparable to those used on surrounding commercial farms, the experiment includes two more diverse low-external-input (LEI) cropping systems: a 3-year rotation (corn–soybean–small grain+red clover) and a 4-year rotation (corn–soybean–small grain+alfalfa–alfalfa) managed with reduced N fertilizer and herbicide inputs.

The 2-year rotation is typical of cash grain farming systems in the region, whereas the 3-year and 4-year rotations are representative of LEI farming systems in the region that include livestock.

The experiment used a randomized complete block design with each crop phase of each rotation system present every year in four replicate blocks. Plots were 0.15 ha and managed with conventional farm machinery. After uniformly cropping the site with oat in 2001 and tuning the three management systems in 2002, intensive data collection began in 2003. Spring triticale was used as the small grain in 2003–2005, whereas oat was used in 2006–2010. During 2003–2010, solid pack manure was applied during the fall preceding corn production in the 3- and 4-year rotations at a mean dry matter rate of 9 Mg ha⁻¹, providing a mean of 120 kg total N ha⁻¹. Corn in the 2-year rotation received 112 kg N ha⁻¹ as urea at planting, whereas corn in the 3- and 4-year rotations did not. The late spring nitrate test³⁵ was used to determine rates for post-emergence side-dress N applications (as urea ammonium nitrate) for corn in all rotation systems. Weed management in the 2-year rotation was based largely on herbicides applied at conventional rates. In the 3- and 4-year systems, herbicides were applied in 38-cm-wide bands over corn and soybean rows rather than broadcast, greater reliance was placed on cultivation, and no herbicides were applied in small grain and forage legume crops. Choices of herbicides used in each system were based on the identities, densities and sizes of weed species observed in the plots. Sampling procedures and other details of farming practices used in the different cropping systems during 2003–2010 are described in Liebman *et al.*³⁶, Cruse *et al.*³⁷ and Gómez *et al.*³⁸.

Results of the experiment indicated that corn and soybean yields from the more diverse 3- and 4-year systems were higher than those from the conventional system, despite substantial reductions in the use of synthetic N fertilizer and herbicides^{36–38} (Table 1). Fossil energy use decreased with increases in crop diversity due to reductions in the use of fertilizers and pesticides, gas to dry grain, and liquid fuel to power farm machinery³⁷ (Table 1). Weed suppression was highly effective in each of the three cropping systems as reflected by the very small amounts of weed biomass found in corn and soybean phases of the rotations³⁶ (Table 1). Although labor requirements were greater in the longer, more diverse rotation systems, net returns to land and management were statistically equivalent in all three cropping systems, with a slight numerical advantage for the 3- and 4-year systems (Table 1).

Biological diversity contributed to the successful functioning of the LEI systems used at the Marsden Farm in multiple ways. For example, though triticale and oat added relatively little revenue themselves to the 3- and 4-year systems³⁶, they served as effective nurse crops for establishing red clover and alfalfa, thereby minimizing

Table 1. Inputs, yields, weed biomass and net returns for the three cropping systems in the Marsden Farm rotation experiment, Boone Co., IA, 2003–2010. Fossil energy inputs are yearly means for 2003–2008; all other data are yearly means for 2003–2010. Within rows, means followed by different letters are significantly different ($P < 0.05$); means not followed by letters are statistically equivalent. Data are from Liebman *et al.*³⁶, Cruse *et al.*³⁷ and Gómez *et al.*³⁸.

	Cropping system		
	2-year rotation: Corn–soybean	3-year rotation: Corn–soybean–small grain ¹ /red clover	4-year rotation: Corn–soybean–small grain ¹ /alfalfa–alfalfa
Whole rotation			
Fertilizer N inputs, kg N ha ⁻¹ yr ⁻¹	76 a	18 b	12 c
Herbicide inputs, kg active ingredients ha ⁻¹ yr ⁻¹	1.91 a	0.28 b	0.21 c
Fossil energy inputs, GJ ha ⁻¹ yr ⁻¹	9.1 a	4.9 b	4.3 c
Labor requirements, h ha ⁻¹ yr ⁻¹	1.8 c	2.8 b	3.5 a
Net returns to land and management ² , \$ ha ⁻¹ yr ⁻¹	690	705	701
Crop yields			
Corn, Mg ha ⁻¹	12.2 b	12.6 a	12.8 a
Soybean, Mg ha ⁻¹	3.4 b	3.7 a	3.8 a
Small grain ¹ , Mg ha ⁻¹	–	3.7	3.8
Alfalfa, Mg ha ⁻¹	–	–	9.0
Weed biomass			
In corn, kg ha ⁻¹	2.0	4.6	3.7
In soybean, kg ha ⁻¹	1.5	3.8	3.6

¹ Triticale was grown as the small grain crop in 2003–2005; oat was used in 2006–2010.

² Crop subsidy payments were not included as sources of revenue.

erosion and reducing weed growth in the absence of herbicides.

Although forage legumes were less profitable than corn³⁶, their inclusion within the diversified rotation systems allowed substantial reductions in nitrogen fertilizer use for corn production^{39,40} (Table 1). Harvest of alfalfa hay and concentrated feeds allowed integration with livestock production and fostered nutrient return to cropland through the application of manure³⁶, further reducing production costs and fossil energy consumption³⁷. In studies conducted by Randall *et al.*²⁷ and Drinkwater *et al.*⁴¹, leaching losses of nitrogen from deep-rooted perennial forage crops, and from corn–soybean systems diversified with forage legumes and small grains were lower than from corn and soybean. Although we did not measure nitrogen losses in our study, those results suggest that nitrogen losses from the diverse LEI systems in the Marsden Farm experiment are likely to have been lower than from the simpler conventional system.

As in our study, other research has shown that lengthening rotations beyond 2 years can increase soybean yields^{42,43}. In addition, both corn and soybean yields have been shown to respond positively to manure application even when the nutrient requirements of those crops are met with other fertility sources^{44–46}. In 2010, the incidence and severity of sudden death syndrome of soybean, caused by the pathogen *Fusarium virguliforme*, and subsequent losses of soybean yield were much lower in the 3- and 4-year rotation systems than in the 2-year

rotation, though the mechanisms for these effects are not yet clear (L.F. Leandro, pers. comm.).

Finally, diversifying the corn–soybean system with small grain and forage crops increased the diversity of habitats available to insects and rodents that preyed upon weed seeds^{47–49}, which is likely to have stabilized seed predator populations and increased their effectiveness in suppressing weed population growth under conditions of reduced herbicide inputs⁵⁰.

The rotation systems and management practices used in the Marsden Farm experiment are well suited to investigations of crop performance, weed dynamics, and economic and energetic costs and returns, but they should not be construed to represent optimal systems. The experiment has not addressed the need for market development for diverse crops to be viable at a larger scale, nor has the experiment provided direct data concerning the impact of the contrasting cropping systems on nutrient and pesticide emissions. Nonetheless, results from the experiment indicate diversified LEI systems for agronomic crops in the US Corn Belt can be highly productive and economically viable, and that further research into their refinement and environmental impacts is warranted. Research focused on the social and economic conditions that help to maintain the dominance of simple cropping systems and that discourage farmers from exploring and adopting more diverse LEI systems would be a useful complement to investigations targeting biophysical characteristics of contrasting cropping systems.

Using Native Perennial Vegetation to Improve Conservation in Annual Cropland: The Science-based Trials of Row-crops Integrated with Prairies (STRIPs) Experiment

Soil and water conservation are critical components of any cropping system, yet, existing rates of erosion within the US Corn Belt indicate that despite high levels of crop productivity, substantial conservation challenges need to be addressed^{51,52}. Conservation practices that have proven useful in restricting the movement of soil sediment within and out of crop fields include buffers and filter strips composed of herbaceous plants, typically perennial grasses. An extensive review by Dosskey *et al.*⁵³ found that sediment trapping efficiencies of buffers generally range between 40 and 85% for buffers that are as small as 2.5% of the source area. Modeling results predict a relatively rapid increase in sediment trapping efficiency as the percentage of buffer area to source area increases from 0 to 10%, with trapping efficiency leveling off as the percentage increases above 10%⁵³. Distributing perennial cover across several contour strips within a watershed in addition to buffers at the bases of catchments may enhance sediment and runoff trapping efficiency since most sediment deposition occurs in the first few meters of the up-slope edge of buffers⁵⁴.

In addition to the amount of vegetation cover, the composition of herbaceous vegetation used for conservation buffers can affect their function. Cool season exotic grasses, such as smooth brome (*Bromus inermis*) and Kentucky bluegrass (*Poa pratensis*), are widely used to provide ground cover in agricultural areas of the US Corn Belt, but are relatively weak-stemmed and prone to lying flat under heavy rain. In contrast, native tall-grass prairie communities are typically dominated by stiff-stemmed warm season grasses, such as Indiangrass (*Sorghastrum nutans*), big bluestem (*Andropogon gerardii*) and switchgrass (*Panicum virgatum*), and a wide range of erect forbs, including species in the genera *Solidago*, *Aster*, *Rudbeckia*, *Ratibida*, *Silphium*, *Helianthus* and *Monarda*⁵⁵, which are less prone to collapse under heavy rain, and which are more effective in providing resistance to water flow and sediment movement^{56,57}.

In addition to reducing water runoff and soil erosion from crop fields, strips and patches of perennial native plants in and around agricultural areas can provide habitat to support native animal populations, including species of conservation concern^{58–63}. Many bird species respond positively to agricultural conservation practices such as grassed waterways^{64,65}, field borders^{66,67}, and riparian buffer strips^{62,68}; the general trend among these studies is increased bird presence, abundance and richness where those small habitats are present adjacent to row crops. Compared with monocultures of cool-season

grasses, diverse prairie communities are expected to provide higher quality bird habitat and harbor more bird species^{65,69,70}.

Beginning in 2004, 12 experimental watersheds (0.5–3.2 ha each) were established within the Neal Smith National Wildlife Refuge (NSNWR), in Jasper Co., IA, to test the hydrologic and biodiversity effects of integrating prairie conservation buffers with row-crop production. The four treatments in the experiment consisted of varying proportions and different configurations of reconstructed native prairie species within corn and soybean fields. For the 100% crop treatment, each replicate watershed was used entirely for corn and soybean production, with the different crops grown in alternate years. There were two 10% prairie/90% crop treatments, one with prairie vegetation concentrated at the toe slope position and a second in which prairie vegetation is distributed in contour strips along the slope, including at the toe slope. In addition, there was a 20% prairie/80% crop treatment that also had prairie vegetation distributed in contour strips and at the toe slope. The experiment used a balanced incomplete block design with three replicates of the four treatments in four blocks. Soils in the experimental watersheds are Alfisols and Mollisols. Analysis of pretreatment data indicated that the watersheds had similar slopes, soil textures, and soil C and N concentrations.

Prior to the start of the experiment, exotic cool-season forage grasses, primarily smooth brome, dominated each of the watersheds. After tillage in the fall of 2006 to kill sod and level the soil, crop production in all watersheds was conducted using continuous no-tillage techniques; soil disturbance occurred only during corn and soybean planting and injection of anhydrous ammonia for corn production. Soybean was grown in 2007 and 2009 and corn was grown in 2008 and 2010 using conventional fertilizer and herbicide practices. Native prairie species were seeded in toe slope and strip positions in July 2007. The strips of prairie species had a minimum width of 4 m and were spaced at a minimum distance of 36 m to accommodate agricultural machinery operations.

Each watershed had a distinct surface flow outlet point where an H-flume was installed to monitor volume and rate of surface runoff. Each of these locations was also instrumented with an automated water sampler to obtain flow measurements and discrete water samples based on flow intervals. In addition, belowground water quality was monitored with a network of suction lysimeters and groundwater wells. Surface and ground water measurements were conducted in 2007–2010. Vegetation surveys were conducted between July and August in 2008, 2009 and 2010 to estimate species richness and determine plant cover using sampling quadrats along transects within the prairie and cropped areas of the watersheds. From 2007 to 2010, all watersheds were visited 9–12 times during spring and summer months and surveyed for birds using spot




Figure 1. Cumulative (a) surface runoff⁷², (b) export of sediment⁷³, (c) total phosphorus export (Helmers *et al.*, unpublished data), and (d) total nitrogen export (Helmers *et al.*, unpublished data). Measurements were made when soil was not frozen. PFS, prairie filter strips.

mapping techniques⁷¹; nest searching and monitoring were added in 2010.

Results of the experiment indicated that prairie buffers had disproportionately large effects on water movement, soil loss and nutrient loss, relative to the amount of annual cropland converted to perennial cover. During 2008, 2009 and 2010, greater water runoff occurred in the 100% crop watersheds than in the mixed perennial-annual watersheds (Fig 1a). Concomitantly, soil sediment loss from 100% crop watersheds was at least 10 times greater than from watersheds with 10 or 20% perennial cover (Fig. 1b). Losses of phosphorus (Fig. 1c) and nitrogen (Fig. 1d) in surface runoff were also markedly higher in the 100% crop watersheds. It is notable that no-tillage production techniques, generally viewed as important for soil conservation, failed to prevent large amounts of soil erosion in the 100% crop watersheds, especially in 2008, a year of intense rainfall during late May and early June.

Although differences in plant species richness and cover were not observed among the three treatments that contained prairie vegetation, plant species richness and cover by native perennial species increased with time in the prairie strips. Dominant native perennial species included Canada goldenrod (*Solidago canadensis*), pinnate prairie coneflower (*Ratibida pinnata*), wild bergamot

(*Monarda fistulosa*), smooth ox-eye (*Heliopsis helianthoides*), Indiangrass (*S. nutans*) and big bluestem (*A. gerardii*). By 2010, prairie strips contained an average of 51 plant species in a total sample area of 6m², of which 35 were native. In contrast, cropped portions of the watersheds (corn in 2010) contained only 15 plant species other than corn, of which 6 were native. Thus, by 2010, conversion of 10–20% of cropland area to reconstructed prairie gave, on average, a 240% increase in overall plant species richness and a 480% increase in native species richness. Transects within cropped areas indicated that the presence of prairie strips did not increase weed infestation in adjacent crops relative to 100% crop watersheds.

A total of 45 bird species were recorded in the watersheds, including several grassland bird species on Iowa's list of species of greatest conservation need¹⁰: common yellowthroat (*Geothlypis trichas*), dickcissel (*Spiza americana*), grasshopper sparrow (*Ammodramus savannarum*), lark sparrow (*Chondestes grammacus*), field sparrow (*Spizella pusilla*) and eastern meadowlark (*Sturnella magna*). Bird species richness was higher after prairie establishment (2008–2010) than during establishment (2007), and tended to be highest in watersheds containing 20% prairie in strips (Fig. 2). Individual bird




Figure 2. Mean bird species richness in the different experimental treatments, 2007–2010 (Schulte *et al.*, unpublished data). Lines represent one standard error. PFS, prairie filter strips.

species' responses to prairie within the watersheds varied among no response, avoidance, disproportionate use, or dependence on prairie habitat. Many bird species used both prairie and crop vegetation, but with a greater frequency of observations in prairie than expected given the extent of the habitat within a watershed (G -values 138–419; $P < 0.0001$). In 2010, 17 nests from four bird species were located and monitored, five of which fledged successfully. Although we do not have enough data to robustly assess nest success, success rates for two grassland bird species, dickcissels and vesper sparrows (*Poocetes gramineus*), are within the range of those observed in other studies^{74,75}. Placement of the STRIPs study within the NSNWR may have had some influence on these results, but they are supported by results of a recent study by Walk *et al.*⁶³, who documented that small patches (1–142 ha) of grassland habitat located within cropland matrices in Illinois could contribute to the nesting success of dickcissels and eastern meadowlarks. In this case, croplands dominated the surrounding landscape (~80% of land cover); furthermore, the investigators did not find a strong influence of patch size on nest success⁶³.

Taken together, these results provide substantial support for the hypothesis that diversifying small areas of annual cropland through the incorporation of perennial prairie vegetation can have disproportionately large beneficial effects on soil, water and biodiversity conservation. Such effects are especially important because projected climate changes in the US Corn Belt are expected to make farming more challenging. The amount of precipitation from heavy and extreme events is expected to increase in this region during the 21st century⁷⁶, with concomitant increases in crop damage⁷⁷ and erosion potential⁷⁸. Biodiversity conservation is also expected to become more challenging as rising crop prices encourage producers to convert conservation grasslands to row-crop production^{79,81}. In a review of climate change impacts

on crop production, Hatfield *et al.*⁸¹ emphasized the need for quantifying how cropping systems can be made more resilient to stress. As shown by results from the STRIPS experiment, strategic targeting to place prairie buffers in landscapes and watersheds dominated by row-crops is likely to be a valuable strategy for pursuing long-term agricultural productivity and environmental quality.

Identifying Trade-offs Among Agroecosystem Performance Indicators: The Comparison of Biofuel Systems (COBS) Experiment

Over the past decade, the USA has had a rapid expansion of interest and investment in transportation fuels derived from plant materials (biofuels), driven in part by concerns over the future availability and costs of fossil energy, and by recognition of the environmental effects of extracting and using fossil energy, including damage from petroleum spills and greenhouse gas emissions. To date, the biofuel boom in the USA has relied almost exclusively on corn grain for ethanol production^{82,83}. However, heavy reliance on corn has led to a range of environmental and conservation concerns, including increases in nitrogen and phosphorus contamination of water^{84,85}, higher rates of soil erosion⁸⁶, greater greenhouse gas emissions due to land conversion^{87,88}, reductions in wildlife habitat and wildlife populations^{89,90}, and declines in biological control of crop pests⁹¹.

Various forms of perennial vegetation have been proposed as alternatives to corn grain for use as biofuel feedstocks, including trees, certain grasses and mixtures of multiple prairie species^{92–94}. These perennials would serve as sources of lignocellulose, rather than starch substrates for fuel production, and might confer a greater degree of environmental protection compared with corn-based systems^{95,96}. Previous studies have shown that, in comparison with corn-based systems, perennial plants used as biofuel feedstocks may result in lower nitrate emissions to drainage water⁹⁷; improved soil physical characteristics that can reduce water runoff, soil erosion and phosphorus transport⁹⁸; greater root growth and below-ground carbon storage^{98,99}; lower greenhouse gas emissions^{100,101}; and better habitat for supporting pollinators, natural enemies of crop insect pests, and other wildlife populations^{17,80,87,102}.

The wide range of environmental as well as agronomic indicators that can be used to assess biofuel production systems calls attention to the multiple goods (e.g., fuel) and ecosystem services (e.g., water quality protection, carbon sequestration and wildlife conservation) that agricultural systems can provide. The multifunctionality of food and feed production systems in Iowa and Minnesota has previously been investigated by Santelmann *et al.*¹⁰³ and Boody *et al.*¹⁰⁴. It is clear from those studies that no one approach for agricultural

production is likely to score highest for all performance indicators. Similarly, the use of perennial species in lieu of corn for biofuel production may also involve trade-offs among key indicators of system performance.

The COBS experiment was initiated in 2008 at the Iowa State University South Reynoldson Farm in Boone Co., IA, to investigate the productivity characteristics and environmental impacts of corn- and prairie-based biofuel production systems. The 9-ha experimental site was on Mollisols with a relatively flat landscape (largely <1% slope, with some small areas of 2–3% slope), and was used for corn and soybean production prior to the start of the COBS experiment.

The experiment contained five cropping systems: continuous corn grown for grain and stover removal; the same corn-based system, but with rye used as a winter cover crop; a corn–soybean system from which only grain was removed; reconstructed multispecies prairie grown for whole-plant removal, without fertilizer; and the same diverse mixture of prairie species grown for whole-plant removal with fertilizer. The experimental plots were 0.17 ha each and arranged in a randomized complete block design with four replicates. All treatments were managed with no tillage. Planting, fertilization, crop protection and harvest operations were conducted with conventional farm machinery. Both prairie treatments were sown in 2008 with a mixture of 31 species native to Boone Co. or adjacent counties, including cool-season and warm-season grasses, legumes and other forbs. All corn treatments received 84 kg N ha⁻¹ at planting, with additional nitrogen applied as a side dressing during early growth, based on treatment-specific soil test results³⁵. In 2010, total nitrogen application rates for corn were 105 kg N ha⁻¹ in the corn–soybean rotation, 123 kg N ha⁻¹ in the continuous corn treatment and 169 kg N ha⁻¹ in the continuous corn plus cover crop treatment. The fertilized prairie treatment received 84 kg N ha⁻¹ in 2010; no N fertilizer was applied to unfertilized prairie or soybean plots. Weeds were suppressed in corn and soybean plots with post-emergence herbicides; no herbicides were applied to the prairie treatments.

The intent of the COBS experiment was to investigate a wide range of performance indicators. Here, we discuss measurements and calculations made in 2010 related to liquid fuel production, nitrogen loss through leaching, and root growth. Corn and soybean seeds, corn stover and prairie biomass were harvested in late September and October, and potential energy yield per unit of field area (GJ ha⁻¹) was calculated for grain-based ethanol, biodiesel and lignocellulosic ethanol, based on measured yields and conversion data given by Shapouri and Gallagher¹⁰⁵, Conley and Tao¹⁰⁶, Hill *et al.*¹⁰⁷, and Tilman *et al.*⁹². Leaching losses of nitrate–N (kg NO₃–N ha⁻¹) from the different treatments were determined using tile drains installed through the center of each plot at a depth of 1.1 m. Tile lines installed on the borders of each plot were used to prevent subsurface drainage from

entering adjacent plots laterally. Flow of water out of central tile lines was metered and flow-proportioned water samples were drawn to monitor nitrate–N concentrations in each plot. After crops were harvested, root biomass (Mg ha⁻¹) was determined using soil cores (80 or 224 cm² plot⁻¹, depending on soil stratum) taken to a depth of 1.0 m, which were subsequently washed in an elutriator to remove soil particles. Remaining materials were then subjected to flotation and inspection under a dissecting microscope, and recovered roots were dried and weighed.

To facilitate comparisons of the relative performance of the different cropping system treatments, means of each treatment were divided by the highest mean value for each of the three performance indicators: energy production, nitrate–N loss and root mass. Results for 2010, shown in Fig. 3, clearly indicated the existence of trade-offs among different facets of cropping system performance. Energy production (unadjusted for fossil energy used in producing and harvesting crop materials) was greatest from the continuous corn systems, least in the unfertilized prairie system, and intermediate in the fertilized prairie and corn–soybean rotation systems. Nitrate–N loss was greatest from the corn–soybean system, almost nil from both the fertilized and unfertilized prairie systems, and intermediate from the continuous corn systems. The unfertilized prairie system produced more roots than all other systems, including the fertilized prairie system. Root mass of the annual crop systems was ≤14% of the root mass of the unfertilized prairie.

One of the striking features of these results was that the fertilized prairie treatment performed well compared with the conventional corn–soybean system for all three indicators: its estimated fuel yield was similar, it lost less nitrate–N in drainage water and it produced more root mass. Tilman *et al.*^{108,109} have shown that above-ground productivity, N retention and root mass can increase with increasing species diversity in prairie communities, due to complementary patterns of resource use among species. High root mass in the prairie treatments of the COBS experiment may have been a consequence of including warm-season (C₄) grasses within the diverse communities, since warm-season grasses can produce more root mass than cool-season (C₃) grasses and legumes¹¹⁰. Economic performance indicators obviously need to be considered in any holistic assessment of biofuel feedstock production systems and their potential for adoption. Nonetheless, the potential for prairie-based biofuel systems as ways to address production and environmental challenges merits further investigation.

Using Biodiversity to Link Agricultural Productivity with Environmental Quality at a Landscape Scale

After a half-century of substantial increases in yield and labor efficiency, agriculture in the US Corn Belt has




Figure 3. Relative performance of different cropping systems in the COBS experiment in 2010 with respect to liquid fuel production (GJ ha^{-1}), nitrogen loss in drainage water ($\text{kg NO}_3\text{-N ha}^{-1}$) and root mass (Mg ha^{-1}). Data for the C-SB treatment represent the average of the corn and soybean phases of the rotation sequence. C-SB, corn–soybean in rotation; CC, continuous corn; CCW, continuous corn with a rye cover crop; PF, fertilized prairie; P, unfertilized prairie. Data used for the calculations are unpublished and are from M. Jarchow, M. Helmers and R. Dietzel.

entered a new and increasingly challenging era. In addition to increasing demands for food, fiber and biofuels, agriculture is now being asked to provide cleaner water, healthier soil, better wildlife habitat, and a range of other environmental goods and services¹¹¹.

In this paper, we have reviewed results from three experiments indicating that strategic timing and placement of perennial crop and non-crop species in agroecosystems dominated by annual crops can have large beneficial effects on agroecosystem functioning. In the Marsden Farm cropping systems experiment, lengthening a simple corn–soybean rotation with small grain and forage crops and recycling nutrients and carbon in the form of manure allowed for 76–84% reductions in the use of synthetic N fertilizer, 85–89% reductions in the use of herbicides and 41–56% reductions in the use of fossil energy, while increasing corn yield 3–5%, increasing soybean yield 9–12%, and maintaining equivalent returns to land and management on a whole rotation basis (Table 1). Results of the STRIPs experiment indicate that diversifying a simple corn–soybean system with small strips and patches of reconstructed prairie vegetation reduced soil and nutrient loss from watershed catchments by >90% (Fig. 1), while also increasing the diversity of native plants and birds (Fig. 2). In the COBS experiment, a fertilized prairie community was found to be capable of producing amounts of liquid fuel comparable to amounts produced by a corn–soybean systems, but with six-fold greater additions to the root C pool and with 97% less

nitrate–N released to drainage water (Fig. 3). In addition to the perennial forage crops, prairie buffer strips and prairie biofuel feedstocks we have discussed here, other vegetation types and practices exist within the ‘perennial portfolio’ for restoring agroecosystem health and function, including riparian buffers, agroforestry plantations and constructed wetlands^{18,19}. Site-specific use of the full range of such practices would lead to a patchwork mosaic of diverse crops and non-crop vegetation interwoven on landscape and watershed scales^{112,113}.

In a world in which rising population and greater affluence have created increased demands for commodity crops such as corn and soybean, should arable land be used differently to achieve adequate levels of food, feed, fuel and fiber production? We believe that it should be. Our argument for increasing biodiversity by moving beyond a narrow base of commodity crops has three parts.

The first part relates to the need to re-couple crop and livestock production. Systems in which livestock are concentrated at high densities and isolated from the land base producing the crops that feed them are often characterized by high levels of pollution¹¹⁴. In contrast, integrated crop–livestock systems, including the diversification of cereal-based rotations with forage crops, the local application of manure, and pasture-based production, are likely to play an important role in reducing water contamination, soil erosion and fossil fuel dependence in agroecosystems, while maintaining high levels of productivity and profitability^{13,37,114–116}. With appropriate forms of crop diversity and integrated management systems, livestock products might be generated in a manner that permits food security to coexist with ecosystem integrity. Results of the Marsden Farm Cropping Systems experiment provide support for the crop portion of this hypothesis, and indicate that diversification and integration can be important ways to reduce reliance on fossil energy and agrichemicals that can become environmental contaminants.

The second part of the argument relates to the need for resilience in the face of climate change. The currently dominant cropping systems in the US Corn Belt were developed in a climate regime different from the one that is beginning to express itself in the region. The types of high intensity rainfall events that are predicted to occur with increasing frequency are exemplified by the storms and flooding that took place in Iowa in May and June of 2008¹¹⁷. These events created severe erosion (defined as loss of $\geq 45 \text{ Mg}$ of soil ha^{-1}) on about 930,000 ha, representing 10% of the state’s arable cropland¹¹⁸. Existing soil conservation practices helped to prevent more damage, but were inadequate for full protection. If agricultural productivity is to be maintained or improved in the future, new conservation practices, such as the small areas of prairie vegetation used in the STRIPs project, need to be developed, refined and implemented. Patches of

perennial vegetation might be used to provide biofuel feedstocks, as in the COBS project, allowing farmers to harvest a crop with use or market value, while still achieving conservation goals^{95,119}. A key research question that needs to be addressed is: how small a proportion of arable land might be converted to perennial cover for various conservation benefits, and how does this proportion vary with site-specific conditions and specific conservation goals?

Our final point relates to the cost and value of increased biodiversity. Currently, agroecosystems are valued almost exclusively for their provisioning services, i.e., their ability to generate food and income. Their ability to conserve soil, protect water quality, provide wildlife habitat and recreational opportunities, and offer other supporting and regulating services is largely overlooked in the marketplace and, in most cases, farmers receive little or no financial compensation for providing these services¹²⁰. The development of markets for organic and grass-based crop and livestock products that command price premiums indicates one approach for compensating farmers for ecosystem services that are linked to particular sets of production practices. Nonetheless, most agricultural land in the USA is used for commodity crops, rather than organic and grass-based production. Between 1995 and 2010, US farmers received \$167 billion in federal crop subsidies for a narrow group of commodity crops, compared with \$35 billion in federal conservation payments¹²¹. Shifting commodity crop subsidies toward conservation and ecosystem services payments could provide strong financial incentives for farmers to increase crop and non-crop diversity at targeted locations within agricultural landscapes, while maintaining farm income. Moreover, a shift of support from commodity subsidies to payments for ecosystem services could generate substantial benefits for society as a whole, including a more stable food supply, cleaner water and improved outdoor recreational experiences. Moving in this direction requires an explicit recognition of the multifunctional nature of agroecosystems and the need for a balance among different functions. Politically, such a move may be difficult. Ecologically, we feel it is necessary to insure that agroecosystems are both agronomically and environmentally sustainable.

Acknowledgements. The authors gratefully acknowledge funding support from the Iowa Department of Agriculture and Land Stewardship; and the Iowa Soybean Association; the Iowa State University College of Agriculture and Life Sciences; the Leopold Center for Sustainable Agriculture; the National Science Foundation; the US Department of Agriculture's National Research Initiative, Agriculture and Food Research Initiative, and Sustainable Agriculture Research and Education programs; the US Fish and Wildlife Service; the US Forest Service; and the US National Aeronautics and Space Administration-US Department of Agriculture Carbon Cycle Science program. The authors also express their sincere thanks for being able to use data from R. Dietzel, R. Gómez, S. Hirsh, M. Jarchow, D. Larsen and A. MacDonald.

References

- 1 National Agricultural Statistics Service (NASS). 2011. 2010 State Agriculture Overview—Iowa. NASS, U.S. Department of Agriculture, Washington, DC. Available at Web site http://www.nass.usda.gov/Statistics_by_State/Ag_Overview/AgOverview_IA.pdf (verified November 15, 2011).
- 2 Nebraska Energy Office. 2011. Ethanol Facilities Capacity by State. Nebraska Energy Office, Lincoln, NE. Available at Web site <http://www.neo.ne.gov/statshtml/121.htm> (verified November 15, 2011).
- 3 Gilliom, R.J., Barbash, J.E., Crawford, C.G., Hamilton, P. A., Martin, J.D., Nakagaki, N., Nowell, L.H., Scott, J.C., Stackelberg, P.E., Thelin, G.P., and Wolock, D.M. 2006. The Quality of our Nation's Waters: Pesticides in the Nation's Streams and Ground Water, 1992–2001. Circular 1291. U.S. Geological Survey, Reston, VA.
- 4 Alexander, R.B., Smith, R.A., Schwarz, G.E., Boyer, E.W., Nolan, J.V., and Brakebill, J.W. 2008. Differences in phosphorus and nitrogen delivery to the Gulf of Mexico from the Mississippi River Basin. *Environmental Science and Technology* 42:822–830.
- 5 Iowa Department of Natural Resources (Iowa DNR). 2010. The Final 2008 Iowa List of Section 303(d) Impaired Waters. Iowa DNR, Des Moines, IA. Available at Web site <http://www.igsb.uiowa.edu/wqm/ImpairedWaters/Year2008/FactSheetForFinal-Approved2008List.pdf>. (verified November 15, 2011).
- 6 Hoekstra, J.M., Boucher, T.M., Ricketts, T.H., and Roberts, C. 2005. Confronting a biome crisis: Global disparities of habitat loss and protection. *Ecology Letters* 8:23–29.
- 7 Klopatek, J.M., Olson, R.J., Emerson, C.J., and Jones, J.L. 1979. Land-use conflicts with natural vegetation in the United States. *Environmental Conservation* 6:191–199.
- 8 Samson, F. and Knopf, F. 1994. Prairie conservation in North America. *BioScience* 44:418–421.
- 9 Schilling, K.E. and Spooner, J. 2006. Effects of watershed-scale land use change on stream nitrate concentrations. *Journal of Environmental Quality* 27:2132–2145.
- 10 Iowa Department of Natural Resources (Iowa DNR). 2007. Iowa Wildlife Action Plan. IDNR, Des Moines, IA. Available at Web site <http://www.iowadnr.gov/Environment/WildlifeStewardship/IowaWildlifeActionPlan.aspx> (verified November 15, 2011).
- 11 National Agricultural Statistics Service (NASS). 2011. Crop Summary: Acres Planted, Harvested, Yield and Production, Iowa and U.S. NASS, U.S. Department of Agriculture, Washington, DC. Available at Web site http://www.nass.usda.gov/Statistics_by_State/Iowa/Publications/Annual_Statistical_Bulletin/2009/47_09.pdf (verified November 15, 2011).
- 12 Hatfield, J.L., McMullen, L.D., and Jones, C.S. 2009. Nitrate–nitrogen patterns in the Raccoon River Basin related to agricultural practices. *Journal of Soil and Water Conservation* 64:190–199.
- 13 Sulc, R.M. and Tracy, B.F. 2007. Integrated crop–livestock systems in the U.S. Corn Belt. *Agronomy Journal* 99:335–345.
- 14 Brown, P.W. and Schulte, L.A. 2011. Agricultural landscape change (1937–2002) in three townships in Iowa, USA. *Landscape and Urban Planning* 100:202–212.

- 15 Kort, J. 1988. Benefits of windbreaks to field and forage crops. *Agriculture, Ecosystems, and Environment* 22/23:165–190.
- 16 Cruse, R.M. and Dinnes, D.L. 1995. Spatial and temporal diversity in production fields. In R. Olson, C. Francis, and S. Kaffka (eds). *Exploring the Role of Diversity in Sustainable Agriculture*. American Society of Agronomy–Crop Science Society of America–Soil Science Society of America, Madison, WI, p. 73–94.
- 17 Gardiner, M.M., Landis, D.A., Gratton, C., DiFonzo, C.D., O’Neal, M., Chacon, J., Wayo, M., Schmidt, N., Mueller, E., and Heimpel, G.E. 2009. Landscape diversity enhances biological control of an introduced crop pest in the north-central U.S. *Ecological Applications* 19:143–154.
- 18 Schulte, L.A., Liebman, M., Asbjornsen, H., and Crow, T.R. 2006. Agroecosystem restoration through strategic integration of perennials. *Journal of Soil and Water Conservation* 61:164–169.
- 19 Asbjornsen, H., Hernandez-Santana, V., Liebman, M., Bayala, J., Chen, J., Helmers, M., Ong, C.K., and Schulte, L. In review. Targeting perennial vegetation in agricultural landscapes for enhancing ecosystem services. *Agriculture, Ecosystems and Environment*.
- 20 Economic Research Service (ERS). 2011. Fertilizer Use and Price. ERS-U.S. Department of Agriculture, Washington, DC. Available at Web site <http://www.ers.usda.gov/Data/FertilizerUse/> (verified November 15, 2011).
- 21 National Agricultural Statistics Service (NASS). 2011. Agricultural Chemical use Database. NASS, U.S. Department of Agriculture, Washington, DC. Available at Web site http://www.pestmanagement.info/nass/app_stat-s3_crop.cfm (verified November 15, 2011).
- 22 Schilling, K.E. and Libra, R.D. 2000. The relationship of nitrate concentrations in streams to row crop land use in Iowa. *Journal of Environmental Quality* 29:1846–1851.
- 23 Jaynes, D.B., Colvin, T.S., Karlen, D.L., Cambardella, C. A., and Meek, D.W. 2001. Nitrate loss in subsurface drainage as affected by nitrogen fertilizer rate. *Journal of Environmental Quality* 30:1305–1314.
- 24 Broussard, W. and Turner, R.E. 2009. A century of changing land-use and water-quality relationships in the continental U.S. *Frontiers in Ecology and the Environment* 7:302–307.
- 25 Sullivan, D.J., Vecchia, A.V., Lorenz, D.L., Gilliom, R.J., and Martin, J.D. 2009. Trends in pesticide concentrations in Corn Belt streams, 1996–2006. Scientific Investigations Report 2009–5132. National Water Quality Assessment Program, U.S. Geological Survey, Reston, VA.
- 26 Burkart, M.R. and James, D.E. 1999. Agricultural-nitrogen contributions to hypoxia in the Gulf of Mexico. *Journal of Environmental Quality* 2:850–859.
- 27 Randall, G.W., Huggins, D.R., Russelle, M.P., Fuchs, D.J., Nelson, W.W. and Anderson, J.L. 1997. Nitrate losses through subsurface tile drainage in Conservation Reserve Program, alfalfa, and row crop systems. *Journal of Environmental Quality* 26:1240–1247.
- 28 Fuhrer, G.J., Gilliom, R.J., Hamilton, P.A., Morace, J.L., Nowell, L.H., Rinella, J.F., Stoner, J.D., and Wentz, D.A. 1999. The Quality of our Nation’s Waters: Nutrient and Pesticides. Circular 1225. U.S. Geological Survey, Reston, VA.
- 29 Hayes, T.B., Collins, A., Lee, M., Mendoza, M., Noriega, N., Stuart, A.A., and Vonk, A. 2002. Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine at low, ecologically relevant doses. *Proceedings of the National Academy of Sciences USA* 99:5476–5480.
- 30 Relyea, R.A. 2005. The impact of insecticides and herbicides on the biodiversity and productivity of aquatic communities. *Ecological Applications* 15:618–627.
- 31 Magdoff, F., Lanyon, L., and Liebhardt, W. 1997. Nutrient cycling, transformations, and flows: Implications for a more sustainable agriculture. *Advances in Agronomy* 60:1–73.
- 32 Dinnes, D.L., Karlen, D.L., Jaynes, D.B., Kaspar, T.C., Hatfield, J.L., Colvin, T.S. and Cambardella, C.A. 2002. Nitrogen management strategies to reduce nitrate leaching in tile-drained Midwestern soils. *Agronomy Journal* 94:153–171.
- 33 Liebman, M. and Gallandt, E.R. 1997. Many little hammers: Ecological management of crop-weed interactions. In L.E. Jackson (ed.). *Ecology in Agriculture*. Academic Press, Orlando, FL, p. 287–339.
- 34 Liebman, M. and Davis, A.S. 2009. Managing weeds in organic farming systems: An ecological approach. In C.A. Francis (ed.). *Organic Farming: The Ecological System*. American Society of Agronomy, Madison, WI, p. 173–196.
- 35 Blackmer, A.M., Voss, R.D., and Mallorino, A.P. 1997. Nitrogen Fertilizer Recommendations for Corn in Iowa. Publication PM-1714. Iowa State University Extension, Ames, IA. Available at Web site <http://www.extension.iastate.edu/Publications/PM1714.pdf> (verified November 15, 2011).
- 36 Liebman, M., Gibson, L.R., Sundberg, D.N., Heggenstaller, A.H., Westerman, P.R., Chase, C.A., Hartzler, R.G., Menalled, F.D., Davis, A.S., and Dixon, P.M. 2008. Agronomic and economic performance characteristics of conventional and low-external-input cropping systems in the central Corn Belt. *Agronomy Journal* 100:600–610.
- 37 Cruse, M.J., Liebman, M., Raman, D.R., and Wiedenhoeft, M. 2010. Fossil energy use in conventional and low-external-input cropping systems. *Agronomy Journal* 102:934–941.
- 38 Gómez, R., Liebman, M., Sundberg, D.N., and Chase, C.A. 2012. Comparison of crop management strategies involving crop genotype and weed management practices in conventional and low-external-input cropping systems. *Renewable Agriculture and Food Systems*, in press. doi:10.1017/S1742170512000142.
- 39 Fox, R.H. and Piekielek, W.P. 1988. Fertilizer N equivalence of alfalfa, birdsfoot trefoil, and red clover for succeeding corn crops. *Journal of Production Agriculture* 1:313–317.
- 40 Morris, T.F., Blackmer, A.M., and El-Hout, N.M. 1993. Optimal rates of nitrogen fertilization for first-year corn after alfalfa. *Journal of Production Agriculture* 6:344–350.
- 41 Drinkwater, L.E., Wagoner, P., and Sarrantonio, M. 1998. Legume-based cropping systems have reduced carbon and nitrogen losses. *Nature* 396:262–265.
- 42 Porter, P.M., Lauer, J.G., Lueschen, W.E., Ford, J.H., Hoverstad, T.R., Oplinger, E.S., and Crookston, R.K. 1997. Environment affects the corn and soybean rotation effect. *Agronomy Journal* 89:441–448.
- 43 Coulter, J.A., Sheaffer, C.C., Wyse, D.L., Haar, M.J., Porter, P.M., Quiring, S.R., and Klossner, L.D. 2011.

- Agronomic performance of cropping systems with contrasting crop rotations and external inputs. *Agronomy Journal* 103:182–192.
- 44 Magdoff, F.R. and Amadon, J.F. 1980. Yield trends and soil chemical change resulting from nitrogen and manure application to continuous corn. *Agronomy Journal* 72:161–164.
- 45 Eghball, B. and Power, J.F. 1999. Composted and non-composted manure application to conventional and no-tillage systems: Corn yield and nitrogen uptake. *Agronomy Journal* 91:819–825.
- 46 McAndrews, G.A., Liebman, M., Cambardella, C.A., and Richard, T.L. 2006. Residual effects of composted and fresh solid swine (*Sus scrofa* L.) manure on soybean [*Glycine max* (L.) Merr.] growth and yield. *Agronomy Journal* 98:873–882.
- 47 Heggenstaller, A.H., Menalled, F.D., Liebman, M., and Westerman, P.R. 2006. Seasonal patterns in post-dispersal seed predation of *Abutilon theophrasti* and *Setaria faberi* in three cropping systems. *Journal of Applied Ecology* 43:999–1010.
- 48 O'Rourke, M.E., Heggenstaller, A., Liebman, M., and Rice, M.E. 2006. Post-dispersal weed seed predation by invertebrates in conventional and low-external-input crop rotation systems. *Agriculture, Ecosystems and Environment* 116:280–288.
- 49 Williams, C.L., Liebman, M., Westerman, P.R., Borza, J., Sundberg, D., and Danielson, B. 2009. Over-winter predation of *Abutilon theophrasti* and *Setaria faberi* seeds in arable land. *Weed Research* 49:439–447.
- 50 Westerman, P.R., Liebman, M., Menalled, F.D., Heggenstaller, A.H., Hartzler, R.G., and Dixon, P.M. 2005. Are many little hammers effective? Velvetleaf population dynamics in two- and four-year crop rotation systems. *Weed Science* 53:382–392.
- 51 Knox, J.C. 2006. Floodplain sedimentation in the Upper Mississippi Valley: Natural versus human accelerated. *Geomorphology* 79(3–4):286–310.
- 52 Cox, C., Hug, A., and Bruzelius, N. 2011. *Losing Ground*. Environmental Working Group, Washington, DC. Available at Web site <http://www.ewg.org/losingground/> (verified November 15, 2011).
- 53 Dosskey, M.G., Helmers, M.J., Eisenhauer, D.E., Franti, T. G., and Hoagland, K.D. 2002. Assessment of concentrated flow through riparian buffers. *Journal of Soil and Water Conservation* 57:336–343.
- 54 Robinson, C.A., Ghaffarzadeh, M., and Cruse, R.M. 1996. Vegetative filter strip effects on sediment concentration in cropland runoff. *Journal of Soil and Water Conservation* 51:227–230.
- 55 Camill, P., McKone, M.J., Sturges, S.T., Severud, W.J., Ellis, E., Limmer, J., Martin, C.B., Navratil, R.T., Purdie, A. J., Sandel, B.S., Talukder, S., and Trout, A. 2004. Community and ecosystem-level changes in a species-rich tall grass prairie restoration. *Ecological Applications* 14:1680–1694.
- 56 Lee, K.H., Isenhardt, T.M., Schultz, R.C., and Mickelson, S. K. 1999. Nutrient and sediment removal by switchgrass and cool-season grass filter strips in Central Iowa, USA. *Agroforestry Systems* 44:121–132.
- 57 Dabney, S.M. 2003. Erosion control, vegetative. In R. Lal (ed.). *Encyclopedia of Soil Science*. Marcel Dekker, New York, NY, p. 209–213.
- 58 Le Couer, D., Baudry, J., Burel, F., and Thenail, C. 2002. Why and how we should study field boundary biodiversity in an agrarian landscape context. *Agriculture, Ecosystems, and Environment* 89:23–40.
- 59 Ma, M., Tarmi, S., and Helenius, J. 2002. Revisiting the species-area relationship in a semi-natural habitat: Floral richness in agricultural buffer zones in Finland. *Agriculture, Ecosystems, and Environment* 89:137–148.
- 60 Marshall, E.J.P. and Moonen, A.C. 2002. Field margins in northern Europe: Their functions and interactions with agriculture. *Agriculture, Ecosystems, and Environment* 89:5–21.
- 61 Van Buskirk, J. and Willi, Y. 2004. Enhancement of biodiversity within set-aside land. *Conservation Biology* 18:987–994.
- 62 Berges, S.A., Schulte, L.A., Isenhardt, T.M., and Schultz, R. C. 2010. Bird species diversity in riparian buffers, row crop fields, and grazed pastures within agriculturally dominated watersheds. *Agroforestry Systems* 79:97–110.
- 63 Walk, J.W., Kershner, E.L., Benson, T.J., and Warner, R.E. 2010. Nesting success of grassland birds in small patches in an agricultural landscape. *The Auk* 127:328–334.
- 64 Bryan, G.G. and Best, L.B. 1991. Bird abundance and species richness in grassed waterways in Iowa rowcrop fields. *American Midland Naturalist* 126:90–102.
- 65 Best, L.B., Freemark, K.E., Dinsmore, J.J., and Camp, M. 1995. A review and synthesis of habitat use by breeding birds in agricultural landscapes of Iowa. *American Midland Naturalist* 134:1–29.
- 66 Conover, R.R., Burger, L.W. Jr, and Linder, E.T. 2009. Breeding bird response to field border presence and width. *The Wilson Journal of Ornithology* 121:548–555.
- 67 Burger, W.L. Jr, Barbour, P.J., and Smith, M.D. 2010. Grassland Bird Population Responses to Upland Habitat Buffer Establishment. *Wildlife Insight*. U.S. Department of Agriculture, Natural Resources Conservation Service, Washington, DC.
- 68 Henningsen, J.C. and Best, L.B. 2005. Grassland bird use of riparian filter strips in southeast Iowa. *Journal of Wildlife Management* 69:198–210.
- 69 Arnold, T.W. and Higgins, K.F. 1986. Effects of shrub coverage on birds on North Dakota mixed-grass prairies. *Canadian Field-Naturalist* 100:10–14.
- 70 Hughes, J.P., Robel, R.J., Kemp, K.E., and Zimmerman, J. L. 1999. Effects of habitat on Dickcissel abundance and nest success in Conservation Reserve Program fields in Kansas. *Journal of Wildlife Management* 63:523–529.
- 71 Ralph, C.J., Geupel, G.R., Pyle, P., Martin, T.E., and DeSante, D.F. 1993. *Handbook of field methods for monitoring landbirds*. General Technical Report PSW-GTR-144. USDA Forest Service Pacific Southwest Research Station, Albany, CA.
- 72 Hernández-Santana, V., Zhou, X., Helmers, M.J., Asbjornsen, H., Kolka, R., and Tomer, M. In review. Native prairie filter strips reduce runoff from hillslopes under annual row-crop systems in Iowa, USA. *Journal of Hydrology*.
- 73 Helmers, M.J., Zhou, X., Asbjornsen, H., Kolka, R., Tomer, M.D., and Cruse, R. 2012. Sediment removal by perennial filter strips in row-cropped ephemeral watersheds. *Journal of Environmental Quality* 41 (5):1531–1539.

- 74 Jones, S.L. and Cornely, J.E. 2002. Vesper sparrow (*Poocetes gramineus*). In A. Poole (ed.). The Birds of North America Online. Cornell Lab of Ornithology, Ithaca, New York. Available at Web site <http://bna.birds.cornell.edu/bna/species/624> (verified November 15, 2011).
- 75 Temple, S.A. 2002. Dickcissel (*Spiza americana*). In: A. Poole (ed.). The Birds of North America Online. Cornell Lab of Ornithology, Ithaca, New York. Available at Web site <http://bna.birds.cornell.edu/bna/species/703> (verified November 15, 2011).
- 76 U.S. Climate Change Science Program/U.S. Global Change Research Program (USCCP/USGCRP). 2000. Climate Change Impacts on the United States—The Potential Consequences of Climate Variability and Change Overview: Midwest. USCCP/USGCRP, Washington, DC. Available at Web site <http://www.usgcrp.gov/usgcrp/Library/nationalassessment/overviewmidwest.htm> (verified November 15, 2011).
- 77 Rosenzweig, C., Tubiello, F.N., Goldberg, R., Mills, E., and Bloomfield, J. 2002. Increased crop damage in the U.S. from excess precipitation under climate change. *Global Environmental Change* 12:197–202.
- 78 Angel, J.R., Palecki, M.A., and Hollinger, S.E. 2005. Storm precipitation in the United States. Part II: Soil erosion characteristics. *Journal of Applied Meteorology* 44:947–959.
- 79 Secchi, S., Tyndall, J., Schulte, L.A., and Asbjornsen, H. 2008. Raising the stakes: High crop prices and conservation. *Journal of Soil and Water Conservation* 63:68A–73A.
- 80 Wiens, J., Fargione, J., and Hill, J. 2011. Biofuels and biodiversity. *Ecological Applications* 21:1085–1095.
- 81 Hatfield, J.L., Boote, K.J., Kimball, B.A., Ziska, L.H., Izaurralde, R.C., Ort, D.R., Thomson, A.M., and Wolfe, D. 2011. Climate impacts on agriculture: Implications for crop production. *Agronomy Journal* 103:351–370.
- 82 Economic Research Service (ERS). 2011. Bioenergy: ERS Research Findings on the Economics of Biofuels. ERS-U.S. Department of Agriculture, Washington, DC. Available at Web site <http://www.ers.usda.gov/Briefing/Bioenergy/ResearchFindings.html> (verified November 15, 2011).
- 83 Renewable Fuels Association (RFA). 2011. Ethanol Facts. RFA, Washington, DC. Available at Web site <http://www.ethanolrfa.org/pages/ethanol-facts> (verified November 15, 2011).
- 84 Donner, S.D. and Kucharik, C.J. 2008. Corn-based ethanol production compromises goal of reducing nitrogen export by the Mississippi River. *Proceedings of the National Academy of Sciences USA* 105:4513–4518.
- 85 Secchi, S., Gassman, P.W., Jha, M., Kurkalova, L., and Kling, C.L. 2011a. Potential water quality changes due to corn expansion in the Upper Mississippi River Basin. *Ecological Applications* 21:1068–1084.
- 86 Secchi, S., Kurkalova, L., Gassman, P.W., and Hart, C. 2011. Land use change in a biofuel hotspot: The case of Iowa, USA. *Biomass and Bioenergy* 35:2391–2400.
- 87 Fargione, J., Hill, J., Tilman, D., Polasky, D., and Hawthorne, P. 2008. Land clearing and the biofuel carbon debt. *Science* 319:1235–1238.
- 88 Searchinger, T., Heimlich, R., Houghton, R.A., Dong, F.X., Elobeid, A., Fabiosa, J., Tokgoz, S., Hayes, D., and Yu, T. H. 2008. Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land use change. *Science* 319:1238–1240.
- 89 Fargione, J.E., Cooper, T.R., Flaspohler, D.J., Hill, J., Lehman, C., McCoy, T., McLeod, S., Nelson, E.J., Oberhauser, K.S., and Tilman, D. 2009. Bioenergy and wildlife: Threats and opportunities for grassland conservation. *BioScience* 59:767–777.
- 90 Meehan, T.D., Hurlbert, A.H., and Gratton, C. 2010. Bird communities in future bioenergy landscapes of the Upper Midwest. *Proceedings of the National Academy of Sciences USA* 107:18533–18538.
- 91 Landis, D.A., Gardiner, M.M., van der Werf, W., and Swinton, S.M. 2008. Increasing corn for biofuel production reduces biocontrol services in agricultural landscapes. *Proceedings of the National Academy of Sciences USA* 105:20552–20557.
- 92 Tilman, D., Hill, J., and Lehman, C. 2006. Carbon-negative biofuels from low-input high-diversity grassland biomass. *Science* 314:1598–1600.
- 93 Heaton, E.A., Dohleman, F.G., and Long, S.P. 2008. Meeting US biofuel goals with less land: The potential of *Miscanthus*. *Global Change Biology* 14:2000–2014.
- 94 U.S. Department of Energy (USDOE). 2011. U.S. billion-ton update: Biomass supply for a bioenergy and bioproducts industry. R.D. Perlack and B.J. Stokes (leads). ORNL/TM-2011/224. Oak Ridge National Laboratory, Oak Ridge, TN. Available at Web site http://www1.eere.energy.gov/biomass/pdfs/billion_ton_update.pdf (verified November 15, 2011).
- 95 Jordan, N., Boody, G., Broussard, W., Glover, J.D., Keeney, D., McCown, B.H., McIsaac, G., Muller, M., Murray, H., Neal, J., Pansing, C., Turner, R.E., Warner, K., and Wyse, D. 2007. Sustainable development of the bioeconomy. *Science* 316:1570–1571.
- 96 Robertson, G.P., Hamilton, S.K., Del Grosso, S.J., and Parton, W.J. 2011. The biogeochemistry of bioenergy landscapes: Carbon, nitrogen, and water considerations. *Ecological Applications* 21:1055–1067.
- 97 McIsaac, G.F., David, M.B., and Mitchell, C.A. 2010. *Miscanthus* and switchgrass production in central Illinois: Impacts on hydrology and inorganic nitrogen leaching. *Journal of Environmental Quality* 39:1790–1799.
- 98 Blanco-Canqui, H. 2010. Energy crops and their implications on soil and environment. *Agronomy Journal* 102:403–419.
- 99 Tufekcioglu, A., Raich, J.W., Isenhardt, T.M., and Schultz, R.C. 2003. Biomass, carbon, and nitrogen dynamics of multispecies riparian buffers within an agricultural watershed in Iowa, USA. *Agroforestry Systems* 57:187–198.
- 100 Adler, P.R., Sanderson, M.A., Weimer, P.J., and Vogel, K. P. 2009. Plant species composition and biofuel yields of conservation grasslands. *Ecological Applications* 19:2202–2209.
- 101 Davis, S.C., Parton, W.J., Dohleman, F.G., Smith, C.M., Del Grosso, S., Kent, A.D., and DeLucia, E.H. 2010. Comparative biogeochemical cycles of bioenergy crops reveal nitrogen-fixation and low greenhouse gas emissions in a *Miscanthus* × *giganteus* agro-ecosystem. *Ecosystems* 13:144–156.
- 102 Fletcher, R.J. Jr, Robertson, B.A., Evans, J., Doran, P.J., Alavalapati, J.R.R., and Schemske, D.W. 2011. Biodiversity conservation in the era of biofuels: Risks and

- opportunities. *Frontiers in Ecology and the Environment* 9:161–168.
- 103 Santelmann, M.V., White, D., Freemark, K., Nassauer, J. L., Eilers, J.M., Vache, K.B., Danielson, B.J., Corry, R.C., Clark, M.E., Polasky, S., Cruse, R.M., Sifneos, J., Rustigian, H., Coiner, C., Wu, J., and Debinski, D. 2004. Assessing alternative futures for agriculture in Iowa, USA. *Landscape Ecology* 19:357–374.
- 104 Boody, G., Vondracek, B., Andow, D.A., Krinke, M., Westra, J., Zimmerman, J., and Welle, P. 2005. Multifunctional agriculture in the United States. *BioScience* 55:27–38.
- 105 Shapouri, H. and Gallagher, P. 2005. USDA's 2002 ethanol cost-of production survey. Agricultural Economic Report 841. U.S. Department of Agriculture, Washington, DC. Available at Web site http://www.usda.gov/oce/reports/energy/USDA_2002_ETHANOL.pdf (verified November 15, 2011).
- 106 Conley, S.P. and Tao, B. 2006. What is Biodiesel? ID-337. Purdue University Extension, West Lafayette, IN. Available at Web site <http://www.ces.purdue.edu/extmedia/ID/ID-337.pdf> (verified November 15, 2011).
- 107 Hill, J., Nelson, E., Tilman, D., Polasky, S., and Tiffany, D. 2006. Environmental, economic, and energetic costs and benefits of biodiesel and ethanol biofuels. *Proceedings of the National Academy of Sciences USA* 103:11206–11210.
- 108 Tilman, D., Wedin, D., and Knops, J. 1996. Productivity and sustainability influenced by biodiversity in grassland ecosystems. *Nature* 379:718–720.
- 109 Tilman, D., Reich, P.B., Knops, J., Wedin, D., Mielke, T., and Lehman, C. 2001. Diversity and productivity in a long-term grassland experiment. *Science* 294:843–845.
- 110 Jarchow, M.E. and Liebman, M. 2012. Tradeoffs in biomass and nutrient allocation in prairies and corn managed for bioenergy production. *Crop Science* 52:1330–1342.
- 111 Robertson, G.P. and Swinton, S.M. 2005. Reconciling agricultural productivity and environmental integrity: A grand challenge for agriculture. *Frontiers in Ecology and the Environment* 3:38–46.
- 112 Olson, R. 1995. Diversity in agricultural landscapes. In R. Olson, C. Francis, and S. Kaffka (eds), *Exploring the Role of Diversity in Sustainable Agriculture*. American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, Madison, WI, p. 121–160.
- 113 Larsen, G.L.D. 2011. Farming for ecosystem services: A case study of multifunctional agriculture in Iowa, USA. M. S. thesis, Iowa State University, Ames, IA.
- 114 Naylor, R., Steinfeld, H., Falcon, W., Galloway, J., Smil, V., Bradford, E., Alder, J., and Mooney, H. 2005. Losing the links between livestock and land. *Science* 310:1621–1622.
- 115 Burkart, M., James, D., Liebman, M., and Herndl, C. 2005. Impacts of integrated crop–livestock systems on nitrogen dynamics and soil erosion in western Iowa watersheds. *Journal of Geophysical Research-Biogeosciences* 110:G01009, doi:10.1029/2004JG000008 (on line).
- 116 Russelle, M.P., Entz, M.H., and Franzluebbers, A.J. 2007. Reconsidering integrated crop–livestock systems in North America. *Agronomy Journal* 99:325–334.
- 117 Mutel, C.F. (ed.). 2010. *A Watershed Year: Anatomy of the Iowa Floods of 2008*. University of Iowa Press, Iowa City, IA.
- 118 Perkins, J. 2008. Soil, Water Conservation Took a Beating in Floods. *Des Moines Register*, July 30, 2008.
- 119 Tilman, D., Socolow, R., Foley, J.A., Hill, J., Larson, E., Lynd, L., Pacala, S., Reilly, J., Searchinger, T., Somerville, C., and Williams, R. 2009. Beneficial biofuels—the food, energy, and environment trilemma. *Science* 325:270–271.
- 120 Power, A.G. 2010. Ecosystem services and agriculture: Tradeoffs and synergies. *Philosophical Transactions of the Royal Society B-Biological Sciences* 365(1554):2959–2971.
- 121 Environmental Working Group (EWG). 2011. Farm Subsidy Database. EWG, Washington, DC. Available at Web site <http://farm.ewg.org/region.php?fips=00000> (verified November 15, 2011).