

2007

Effects of Long-Term Tillage and Crop Rotation on Soil Carbon and Soil Productivity

Mahdi Al-Kaisi

Iowa State University, malkaisi@iastate.edu

Mark A. Licht

Iowa State University, lichtma@iastate.edu

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Al-Kaisi, Mahdi and Licht, Mark A., "Effects of Long-Term Tillage and Crop Rotation on Soil Carbon and Soil Productivity" (2007).
Iowa State Research Farm Progress Reports. 826.
http://lib.dr.iastate.edu/farms_reports/826

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Effects of Long-Term Tillage and Crop Rotation on Soil Carbon and Soil Productivity

Abstract

Tillage system and crop rotation have long-term effects on soil productivity and soil quality components such as soil carbon as well as on soil physical, biological, and chemical properties. In addition, both tillage and crop rotation have effects on weed and soil disease control. There is a need for well-defined, long term tillage and crop rotation studies across the different soils and climate conditions in Iowa. The objective of this study was to evaluate the long-term effects of different tillage systems and crop rotations on soil productivity.

Keywords

Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences

Effects of Long-Term Tillage and Crop Rotation on Soil Carbon and Soil Productivity

Mahdi Al-Kaisi, associate professor
Department of Agronomy
Mark Licht, crops field specialist
ISU Extension

Introduction

Tillage system and crop rotation have long-term effects on soil productivity and soil quality components such as soil carbon as well as on soil physical, biological, and chemical properties. In addition, both tillage and crop rotation have effects on weed and soil disease control. There is a need for well-defined, long-term tillage and crop rotation studies across the different soils and climate conditions in Iowa. The objective of this study was to evaluate the long-term effects of different tillage systems and crop rotations on soil productivity.

Materials and Methods

The study was originated and conducted on eight Iowa State University research and demonstration farms in 2002 and continued through 2005. Treatments include five tillage systems (no-till, strip-tillage, chisel plow, deep ripper, and moldboard plow) and two crop rotations (corn-corn-soybean and corn-soybean) across the five tillage systems and several soil associations. Initial soil samples were collected in 2002 prior to implementing the tillage treatments. Soil samples were subsequently collected in 2004. The soil samples were collected from all sites at depths of 0–6, 6–12, 12–18, and 18–24 in. and were analyzed for total carbon and total nitrogen. The

experimental design was a randomized complete block design with four replications.

The plot size was 20 rows × 65 ft. Yield was determined from the center four rows of each plot. Long-term effects of tillage and crop rotation on total soil carbon and total nitrogen were monitored on a bi-yearly or more basis. Seasonal measurements such as nitrogen use efficiency, soil bulk density, and infiltration rate were conducted on selected sites depending on availability of funding.

Results and Discussion

The average corn yields across all tillage systems for the corn-soybean and corn-corn-soybean rotation in 2006 were 205.0 and 207.9 bushels/acre, respectively (Table 1 and 2). The corn-corn-soybean rotation was in the first year of corn.

Under the corn-soybean rotation, the moldboard plow treatment more than the no-tillage treatment, but all other treatments were not different.

Under the corn-corn soybean rotation the moldboard and chisel plow treatments yielded more than the no-tillage treatment, but all other treatments were not different.

Acknowledgments

We would like to thank Bernard Havlovic and Jeff Butler for their time and labor for plot setup, planting, and harvesting.

Table 1. Corn and soybean yields under a corn-soybean rotation at the ISU Armstrong Research Farm. Yields are corrected to 15.5 and 13.0% for corn and soybean, respectively.

	Corn (C-s)			Soybean (c-S)	
	2002 ^b	2004	2006	2003 ^b	2005
	----- bushels/acre -----				
No-tillage	92.2	214.9	195.5	39.8	55.6
Strip-tillage	91.4	218.9	202.4	38.3	55.6
Deep rip	91.0	235.1	206.9	39.7	60.8
Chisel plow	88.3	232.0	206.9	35.7	56.6
Moldboard plow	107.4	226.3	213.1	33.8	55.7
LSD _(0.05) ^a	20.8	14.2	17.1	3.5	4.6
5-Tillage average	94.1	225.4	205.0	37.5	56.9

^aLeast significant differences (LSD_(0.05)) are based on a Fisher test.

Yield differences greater than the least significant difference are significantly different.

^bWeather conditions in 2002 and 2003 were 12.25 and 10.51 in. of precipitation below normal.

Table 2. Corn and soybean yields under a corn-corn-soybean rotation at the ISU Armstrong Research Farm. Yields are corrected to 15.5 and 13.0% for corn and soybean respectively.

	Corn (C-c-s)		Corn (c-C-s)	Soybean (c-c-S)	
	2003 ^b	2006	2004	2002 ^b	2005
	----- bushels/acre -----				
No-tillage	151.8	196.5	221.0	36.7	56.3
Strip-tillage	142.7	208.2	224.3	35.7	56.8
Deep rip	146.3	209.6	231.8	35.5	61.0
Chisel plow	136.8	211.9	228.7	36.7	59.1
Moldboard plow	133.8	213.1	238.2	35.7	55.4
LSD _(0.05) ^a	17.5	14.5	11.5	6.4	4.2
5-Tillage average	142.3	207.9	228.8	36.1	57.7

^aLeast significant differences (LSD_(0.05)) are based on a Fisher test.

Yield differences greater than the least significant difference are significantly different.

^bWeather conditions in 2002 and 2003 were 12.25 and 10.51 in. of precipitation below normal.