

6-4-2007

## Soybean aphids found in Minnesota and Wisconsin

Marlin E. Rice

Iowa State University, [merice@iastate.edu](mailto:merice@iastate.edu)

Matthew E. O'Neal

Iowa State University, [oneal@iastate.edu](mailto:oneal@iastate.edu)

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

---

### Recommended Citation

Rice, Marlin E. and O'Neal, Matthew E., "Soybean aphids found in Minnesota and Wisconsin" (2007). *Integrated Crop Management News*. 1048.

<http://lib.dr.iastate.edu/cropnews/1048>

**The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.**

---

# Soybean aphids found in Minnesota and Wisconsin

## **Abstract**

Our entomological colleagues to the north and northeast are reporting the first soybean aphids of the summer on seedling soybean plants. Eileen Cullen, extension entomologist, University of Wisconsin, reported that in a research plot near Madison on May 24, 8 of 60 plants were infested with 1-13 aphids per plant. Similar populations were found near Arlington on May 30. David Ragsdale, research entomologist, University of Minnesota, reported soybean aphids on V1-stage soybeans in east-central Minnesota during the week of May 28.

## **Keywords**

Entomology

## **Disciplines**

Agricultural Science | Agriculture | Entomology

# INTEGRATED CROP MANAGEMENT

Search

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2007 > IC-498(13) -- June 4, 2007

## Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

## Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

[Image Gallery](#)

## Printable Version

Printable version of this page

## Related Articles

Aphids attack!  
**August 6, 2007**

Extension agronomists update soybean aphid situation  
**July 30, 2007**

Newly revised aphids publication available  
**July 23, 2007**

Mid-July soybean aphid update

## Soybean aphids found in Minnesota and Wisconsin

by Marlin E. Rice and Matt O'Neal, Department of Entomology

Our entomological colleagues to the north and northeast are reporting the first soybean aphids of the summer on seedling soybean plants.



*Winged soybean aphids fly from buckthorn to soybean where they give birth to live aphids. (Marlin E. Rice)*

Eileen Cullen, extension entomologist, University of Wisconsin, reported that in a research plot near Madison on May 24, 8 of 60 plants were infested with 1-13 aphids per plant. Similar populations were found near Arlington on May 30. David Ragsdale, research entomologist, University of Minnesota, reported soybean aphids on V1-stage soybeans in east-central Minnesota during the week of May 28. Some fields had more than 25 percent of the plants infested with an average density of 5 aphids per plant. Furthermore, Chris DiFonzo, extension entomologist, Michigan State University, reported three research farms with infestations of soybean aphids. In Michigan, all sites had less than 1 percent of plants infested with aphids.

In Iowa in recent years, soybean aphids have been found during the last week of May in central or northeastern Iowa, but as of June 4, we have not yet received any reports of aphids in Iowa soybeans. Our aphid suction trap network with traps located near Ames, Chariton, Nashua, and Sutherland did not collect any winged soybean aphids the weeks preceding May

**July 16, 2007**

Soybean aphids exceed the economic threshold in northeast Iowa

**July 9, 2007**

Monitor soybean aphid populations on PIPE

**July 2, 2007**

Soybean aphid numbers increase...and decrease

**July 2, 2007**

Early soybean aphid outbreak--to the east

**June 11, 2007**

PIPE: Pest Information Platform for Extension and Education

**March 26, 2007**

Purdue University identifies the #1 predator of soybean aphids

**March 26, 2007**

18 and May 25. This may suggest that the Iowa population is small this year or maybe not because the traps won't necessarily represent the entire state. Also, it may take us several years to be able to interpret what the spring soybean aphid trap captures mean in relationship to early summer infestations in soybean fields.

The important point to remember is that soybean aphids will appear somewhere in Iowa in damaging numbers this summer--and field scouting will be necessary to determine the need for chemical control.

As of yet, no damaging populations occur in Iowa, but field scouting should start in 2-3 weeks so that informed decisions can be made in a timely manner.

*Marlin E. Rice is a professor of entomology with extension and research responsibilities. Matt O'Neal is an assistant professor of entomology with research and extension responsibilities in field crops.*

This article originally appeared on page 167 of the IC-498(13) -- June 4, 2007 issue.

Updated 06/06/2007 - 11:09am