

2006

Evaluation of Previous Rotation on Flax Performance in Certified Organic Production - Crawfordsville Trial, 2005

Kathleen Delate

Iowa State University, kdelate@iastate.edu

Andrea McKern

Iowa State University

Daniel Rosmann

Iowa State University

Kevin Van Dee

Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), and the [Horticulture Commons](#)

Recommended Citation

Delate, Kathleen; McKern, Andrea; Rosmann, Daniel; and Van Dee, Kevin, "Evaluation of Previous Rotation on Flax Performance in Certified Organic Production - Crawfordsville Trial, 2005" (2006). *Iowa State Research Farm Progress Reports*. 1149.

http://lib.dr.iastate.edu/farms_reports/1149

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Evaluation of Previous Rotation on Flax Performance in Certified Organic Production - Crawfordsville Trial, 2005

Abstract

Flax (*Linum usitatissimum* [Linaceae] – linen family) is an ancient crop that had been grown in Iowa for many years, but it has been displaced by the emphasis on commodity corn and soybeans. Flax has many uses, including industrial oils from oilseed flax; food-quality flaxseed oil; and linen products, fiberboard and paper products from the straw. Flaxseed oil is high in omega-3 fatty acids, which are associated with lowered risk of heart disease and lowered blood cholesterol levels. Flax has a 50-day vegetative period, a 25-day flowering period, and a 35-day period to maturity. Seeds are produced in bolls that contain 6–10 seeds. Seed color can be brown, golden, or yellow. The seed is covered with a mucilaginous coating. The flax crop responds to up to 50 lb/acre nitrogen, similar to organic small grains. Mycorrhizal association may increase the ability of flax to take up phosphorus from the soil, so growing flax after mycorrhizal wheat rather than after nonmycorrhizal canola may improve its phosphorus uptake by flax. Early-seeded flax generally produces the highest yields, when using the same planting dates as small grains. Frost seldom kills flax seedlings. Nonuniform maturity and ripening are problems in late-seeded fields.

Keywords

Horticulture, Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Horticulture

Evaluation of Previous Rotation on Flax Performance in Certified Organic Production—Crawfordsville Trial, 2005

Kathleen Delate, associate professor
Andrea McKern, research associate
Daniel Rosmann, research associate
Departments of Horticulture and Agronomy
Kevin Van Dee, farm superintendent

Introduction

Flax (*Linum usitatissimum* [Linaceae] – linen family) is an ancient crop that had been grown in Iowa for many years, but it has been displaced by the emphasis on commodity corn and soybeans. Flax has many uses, including industrial oils from oilseed flax; food-quality flaxseed oil; and linen products, fiberboard and paper products from the straw. Flaxseed oil is high in omega-3 fatty acids, which are associated with lowered risk of heart disease and lowered blood cholesterol levels. Flax has a 50-day vegetative period, a 25-day flowering period, and a 35-day period to maturity. Seeds are produced in bolls that contain 6–10 seeds. Seed color can be brown, golden, or yellow. The seed is covered with a mucilaginous coating. The flax crop responds to up to 50 lb/acre nitrogen, similar to organic small grains. Mycorrhizal association may increase the ability of flax to take up phosphorus from the soil, so growing flax after mycorrhizal wheat rather than after nonmycorrhizal canola may improve its phosphorus uptake by flax. Early-seeded flax generally produces the highest yields, when using the same planting dates as small grains. Frost seldom kills flax seedlings. Nonuniform maturity and ripening are problems in late-seeded fields.

In 2003, an organic flaxseed oil-processing facility began operations in Iowa. With the introduction of this plant and a possible worldwide market for flaxseed oil came a need for increased organic flax production in Iowa.

Materials and Methods

In the Crawfordsville organic flax trial, Norlin flax was seeded on March 29–30, 2005, at 60 lb/acre with multi-year red clover underseeded at 14 lb/acre. Flax was seeded into three blocks: 2004 crop of corn; 2004 crop of soybean; and 2004 crop of barley/red clover. Flax plant populations were counted on May 25 and flax height and weed populations were counted on May 25 and June 29, 2005. Flax height was taken by measuring three random plants in each plot. Flax population counts were taken by placing a 1-ft² quadrat in three areas of each plot and counting the number of plants inside the quadrat. Weed counts were taken by placing a 1-m² quadrat in three areas of each plot and counting the number of broadleaf and grass weeds. Biomass samples were taken by randomly clipping three 1-ft² sections from each plot on June 1 and July 12. The biomass samples were weighed and separated into flax, red clover, and weeds; then they were placed in a dryer at 155°F for 48 hours, after which time separate dry weights were taken for each group. Twelve random 1-ft² flax harvest plots per block were cut at ground level and hand-threshed on July 25. The remainder of the flax was harvested with a combine on July 25–29, 2005. Soil samples were taken on June 1 and August 11, 2005, from five random locations within each plot at a depth of 6 in.

Results and Discussion

Yields were significantly different based on the crop of the previous year. The flax field that was in barley/red clover in 2004 yielded 33.14 bushels/acre, followed by the flax that was in soybean (25.79 bu/acre), and finally the flax that followed corn (13.60 bu/acre) (Table 1). Stand counts were higher in the flax following soybean than in flax following corn and barley/red clover, which had similar stand

counts (Table 1). Red clover biomass was higher in the flax following soybeans than the other two trials (Table 1). No significant difference was observed in plant height on May 25, but on June 29, the flax following corn was found to be significantly shorter than the other two varieties (Table 1). The flax following barley/red clover had significantly higher flax biomass (Table 1) despite having higher numbers of grass weeds than the other two blocks (Table 2). Broadleaf weeds (mainly lamb's quarters) were more abundant in the flax following corn than in the other two trials, however, and contributed to the greatest weed biomass (Table 2). Significantly higher levels of NO₃-N were found in soils where barley/red

clover were grown in 2004. Recommendations from this study include the need for a nitrogen-fixing crop in the rotation prior to flax planting for maximum yield.

Acknowledgments

We would like to thank the Leopold Center for Sustainable Agriculture and Mary Wiedenhoft (Agronomy Department, ISU) for their support of this project. Thanks go to Mark and Julie Schuett of BIOWA Nutraceuticals, Cherokee, IA, and Spectrum Organic Products, Petaluma, CA, for supplying seed and assistance with this project. Thanks also go to Alzbeta Novotna for her help on production, data collection, and analytical aspects of this project.

Table 1. Organic flax plant performance, Crawfordsville, 2005.

2004 Crop	Stand plants/ft ²	Plant height (cm)	Plant height (cm)	Flax dry weight (lb/acre)	Red clover dry weight (lb/acre)	Yield (bu/acre)
		May 25	June 29			
Barley/red clover	89.75b	31.75	61.92a	3,554a	0.00b	33.14a
Corn	91.18b	30.46	52.58b	1,639c	31.6ab	13.60c
Soybeans	114.33a	30.67	61.67a	2,784b	115.0a	25.79b
LSD 0.05	15.78	NS	3.01	727	104.5	3.71

Table 2. Weed populations in organic flax, Crawfordsville, 2005.

2004 Crop	Broadleaf weeds/m ²	Grass weeds/m ²	Broadleaf weeds/m ²	Grass weeds/m ²	Weed weight (lb/acre)
	May 25, 2005		June 29, 2005		
Barley/red clover	4.83a	142.75b	5.00a	32.67b	2.9a
Corn	27.46b	10.27a	48.25c	6.00a	527.1b
Soybeans	11.50a	0.08a	19.92b	0.00a	70.0a
LSD 0.05	8.71	30.38	9.87	6.73	330.6

Table 3. Soil quality in organic flax, Crawfordsville, 2005.

2004 Crop	Bray-P (ppm)	K (ppm)	Ca (ppm)	Mg (ppm)	NO ₃ -N (ppm)	NH ₄ -N (ppm)	pH
Barley/red clover	18.50	181.50	3558.6b	640.13c	11.50a	6.13	6.54b
Corn	17.75	169.38	3878.8a	731.25b	1.38b	5.13	6.89a
Soybeans	19.13	183.63	3858.0a	783.13a	1.75b	5.63	6.78a
LSD 0.05	NS	NS	153.8	51.87	1.49	NS	0.22