

2-25-2010

The effects of age and type of carrying task on lower extremity kinematics

Jason C. Gillette
Iowa State University, gillette@iastate.edu

Catherine A. Stevermer
Iowa State University

Ross H. Miller
University of Massachusetts

Stacey A. Meardon
Iowa State University

Charles V. Schwab
Iowa State University, cvschwab@iastate.edu

Follow this and additional works at: https://lib.dr.iastate.edu/abe_eng_pubs



Part of the [Biomechanics Commons](#), [Bioresource and Agricultural Engineering Commons](#), and the [Ergonomics Commons](#)

The complete bibliographic information for this item can be found at https://lib.dr.iastate.edu/abe_eng_pubs/1115. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Agricultural and Biosystems Engineering at Iowa State University Digital Repository. It has been accepted for inclusion in Agricultural and Biosystems Engineering Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

The effects of age and type of carrying task on lower extremity kinematics

Abstract

The purpose of this study was to determine the effects of age, load amount and load symmetry on lower extremity kinematics during carrying tasks. Forty-two participants in four age groups (8–10 years, 12–14 years, 15–17 years and adults) carried loads of 0%, 10% and 20% body weight (BW) in large or small buckets unilaterally and bilaterally. Reflective markers were tracked to determine total joint range of motion and maximum joint angles during the stance phase of walking. Maximum hip extension, hip adduction and hip internal rotation angles were significantly greater for each of the child/adolescent age groups as compared with adults. In addition, maximum hip internal rotation angles significantly increased when carrying a 20% BW load. The observation that the 8–10-year-old age group carried the lightest absolute loads and still displayed the highest maximum hip internal rotation angles suggests a particular necessity in setting carrying guidelines for the youngest children.

Keywords

biomechanics, gait, load carriage, youth

Disciplines

Biomechanics | Bioresource and Agricultural Engineering | Ergonomics

Comments

This is an Accepted Manuscript of an article published by Taylor & Francis in *Ergonomics* on 25 Feb 2010, available online: DOI: [10.1080/00140130903402234](https://doi.org/10.1080/00140130903402234). Posted with permission.

Published in final edited form as:

Ergonomics. 2010 March ; 53(3): 355–364. doi:10.1080/00140130903402234.

The effects of age and type of carrying task on lower extremity kinematics

Jason C. Gillette^{a, *}, Catherine A. Stevermer^a, Ross H. Miller^b, Stacey A. Meardon^a, and Charles V. Schwab^c

Jason C. Gillette: gillette@iastate.edu; Catherine A. Stevermer: ktsteve@iastate.edu; Ross H. Miller: rhmiller@kin.umass.edu; Stacey A. Meardon: smearndon@iastate.edu; Charles V. Schwab: cvschwab@iastate.edu

^aDepartment of Kinesiology, Iowa State University, Ames, Iowa, USA

^bDepartment of Kinesiology, University of Massachusetts, Amherst, MA, USA

^cDepartment of Agricultural and Biosystems Engineering, Iowa State University, Ames, Iowa, USA

Abstract

The purpose of this study was to determine the effects of age, load amount, and load symmetry on lower extremity kinematics during carrying tasks. Forty-two participants in four age groups (8-10 years, 12-14 years, 15-17 years, and adults) carried loads of 0%, 10%, and 20% body weight (BW) in large or small buckets unilaterally and bilaterally. Reflective markers were tracked to determine total joint ROM and maximum joint angles during the stance phase of walking. Maximum hip extension, hip adduction, and hip internal rotation angles were significantly greater for each of the child/adolescent age groups as compared to adults. In addition, maximum hip internal rotation angles significantly increased when carrying a 20% BW load. The observation that the 8-10 year old age group carried the lightest absolute loads and still displayed the highest maximum hip internal rotation angles suggests a particular necessity in setting carrying guidelines for the youngest children.

Keywords

biomechanics; gait; load carriage; youth

1. Introduction

According to the United States Department of Agriculture, children suffered 23,000 agricultural-related injuries in 2001 (USDA 2004). Studies involving children participating in agricultural work have often focused on fatal and/or traumatic injuries (Rivara 1985, Bancej and Arbuckle 2000, Reed and Claunch 2000, Gerberich *et al.* 2001). To avoid hazardous

*Corresponding author. gillette@iastate.edu.

Jason C. Gillette (corresponding author), Iowa State University, Department of Kinesiology, 245 Forker Building, Ames, IA, USA 50011-1160, Phone: +1-515-294-8310, Fax: +1-515-294-8740

Catherine A. Stevermer, Iowa State University, Department of Kinesiology, 283 Forker Building, Ames, IA, USA 50011-1160, Phone: +1-515-294-2953, Fax: +1-515-294-8740

Ross H. Miller, University of Massachusetts, Department of Kinesiology, 110 Totman Building, Amherst, MA, USA 01003, Phone: +1-413-545-6075, Fax: +1-413-545-2906

Stacey A. Meardon, Iowa State University, Department of Kinesiology, 283 Forker Building, Ames, IA, USA 50011-1160, Phone: +1-515-294-2953, Fax: +1-515-294-8740

Charles V. Schwab, Iowa State University, Department of Agricultural and Biosystems Engineering, 220 Industrial Education II, Ames, IA, USA 50011-3130, Phone: +1-515-294-4131, Fax: +1-515-294-1123

Statement of relevance: Bucket carrying tasks were analyzed as a function of age group, load amount, and load symmetry. Hip joint rotations significantly increased when carrying 20% body weight loads and in children as compared to adults, which suggests particular necessity in setting carrying guidelines for the youngest age group (8-10 year olds).

activities, children commonly perform animal care tasks that are deemed to be relatively safe such as carrying feed and water, lifting bales, and spreading bedding (Marlenga *et al.* 2001, Allread *et al.* 2004). However, these tasks may require children to carry loads asymmetrically that are proportionally large and/or heavy. Children do work with adult equipment or perform tasks designed for adults who are larger and stronger (Miller and Kaufman 1998, Allread and Waters 2007). Thus, children may suffer musculoskeletal disorders due to a lack of strength or because they are still developing the movement skills to perform these challenging tasks. One study found sprains/strains to be the most common type of farm injury and the upper limbs, lower limbs, and back to be the most commonly affected sites (Pickett *et al.* 1995). Similarly, surveys of farm children have found that muscle strains of the legs, arms, shoulders, and back are considered everyday occurrences (Bartels *et al.* 2000). A field study confirmed that farm children perform lifting and carrying motions that are equivalent to those associated with high injury risk jobs in industry (Allread *et al.* 2004).

Farm children and adults have been studied to determine the effects of different bucket carrying tasks on upper extremity and low back moments (Gillette *et al.* 2009). The 8-10 year old age group displayed lower maximum normalized shoulder abduction and L5/S1 lateral bending moments than adults across carrying conditions. One explanation for these results is that the 8-10 year olds were able to compensate for smaller anthropometry by altering their carrying technique and upper body posture. What remains unknown is whether simultaneous adjustments in lower body kinematics are also required.

Prior studies suggest that whole body gait parameters do not necessarily change in adults when carrying a load. Unilateral load carrying at 15% body weight (BW) produced only minor changes in walking speed and stride frequency as compared to unloaded gait in adults (Crowe and Samson 1997). Similarly, walking speed and stride frequency were not significantly different for unilateral carrying (average 20% BW) and bilateral carrying (average 38% BW) as compared to unloaded gait in adults (Nottrodt and Manley 1989). Previous studies were inconclusive as to whether or not whole body gait parameters change in children when loads are carried. For example, decreased stride length and increased stride frequency were observed for 11-13 year olds carrying one-strap backpacks at 17% BW (Pascoe *et al.* 1997). In addition, increased stance duration was reported for 10 year old boys carrying dual-strap backpacks when walking on a treadmill at 20% BW loads (Hong and Brueggemann 2000). However, another study did not find significant differences in walking length and stride frequency for 9-10 year old boys carrying dual-strap backpacks over ground at 20% BW (Hong and Cheung 2003). When focusing on individual joint kinematics, increased hip sagittal range of motion and maximum hip adduction during stance were reported for adolescent girls carrying dual-strap backpacks at 15% BW (Chow *et al.* 2006). Thus, it appears likely that children do alter their joint coordination during carrying tasks, but few studies have examined lower extremity kinematics across a range of ages and different loading conditions.

One potentially confounding variable in the assessment of load carrying tasks is physical maturation. When adjustments were made for body size, changes in stride frequency as a function of walking velocity during unloaded gait were found to be similar when comparing children 4-10 years old to adults (Zijlstra *et al.* 1996). Similarly, a review of youth gait concluded that adult-like walking velocity and stride frequency are achieved by 4 years of age when adjusting for leg length (Sutherland 1997). This review also reported differences in joint coordination in that younger children use hip flexors/extensors more and ankle plantarflexors less than adults to achieve similar whole body gait parameters. Another study found that 7 year-old children have similar hip and knee joint angle patterns to adults during gait, but reported reduced ankle control (Ganley and Powers 2005). Taken together, these studies indicate that while children's normalized whole body gait parameters may be similar to adults, children may rely more on control of hip rotations to achieve these patterns. There also appears

to be a relative lack of joint kinematic data comparing children and adults in the frontal and transverse planes during carrying tasks. Compared to unloaded gait, the additional strength and balance challenges of bilateral and unilateral load carrying may magnify kinematic differences between children and adults.

Changes in lower extremity kinematics from unloaded gait to carrying tasks indicate that adjustments were necessary to complete the movement. Adjustments that bring joint rotations near the limits of their ranges of motion are of particular concern in regards to soft tissue strains and injury. Ultimately, changes in lower extremity kinematics may be a factor to consider when developing load-carrying guidelines for children. For load amount, studies discussed above suggested that loads carried should not exceed 10-15% BW in children (Hong and Brueggemann 2000, Hong and Cheung 2003). Similarly, the North American Guidelines for Children's Agricultural Tasks recommend limiting children's work that involves lifting and moving objects greater than 15% BW (Lee and Marlenga 1999). However, field data suggest that these standards are exceeded for children during common farm tasks (Allread *et al.* 2004). For load asymmetry, unilateral carrying may increase frontal plane motion. In addition, unilateral carrying of a proportionally large bucket positions the load further from the body and may further increase frontal plane motion. These factors are in need of more investigation for children participating in agricultural work.

The purpose of this study was to determine the effects of age, load amount, and load symmetry on lower extremity kinematics during carrying tasks in adults and in children at different ages. Consistent with increased use of hip flexors/extensors in young children (Sutherland 1997) and increased hip sagittal plane range of motion during dual-strap backpack carrying (Chow *et al.* 2006), the first set of hypotheses was that 8-10 year olds would display increased hip sagittal plane range of motion and greater maximum hip flexion angles during bucket carrying as compared to adults. Consistent with potentially reduced ankle control in children (Ganley and Powers 2005) and frontal plane loading asymmetry during bucket carrying, the second set of hypotheses was that 8-10 year olds would display increased ankle frontal plane range of motion and maximum ankle eversion angles during bucket carrying as compared to adults. Carrying greater load amounts unilaterally in larger buckets were expected to be the most challenging tasks for all participants in this study, particularly for the youngest age group.

2. Methods

2.1. Subjects

Forty-two participants in four age groups (8-10 years, 12-14 years, 15-17 years, and adults between 21-26 years) volunteered for this study. These individuals were a subset of a larger pool of 72 participants over a three year period that had their upper and/or lower body biomechanics analyzed during carrying tasks. Each individual provided informed consent as approved by the Office of Research Assurances prior to taking part in the study. Parental informed consent was also obtained for the minors participating in the data collection. Children were recruited through the county extension offices and the local 4-H chapters, which are youth organizations sponsored by the United States Department of Agriculture where participants regularly perform farming tasks. Gender distributions, average age, average height, and average mass for each age group are presented in Table 1.

2.2. Experimental protocol

Two sizes of buckets were carried: large five-gallon (18.9 litre volume, 36.8 cm high, 30 cm diameter) and small one-gallon (3.8 litre volume, 19.5 cm high, 16.7 cm diameter). The buckets were filled using sealed bags of lead shot amounting to three levels of load based on body weight: 0%, 10%, and 20% BW in addition to the weight of the bucket. Unilateral carrying

tasks were performed with both large and small buckets, while bilateral carrying tasks were only performed with small buckets. Bilateral carrying with large buckets was not performed due to the difficulty observed in some 8 year old children completing this task during initial testing and due to potential fatigue considerations limiting the number of conditions. In Figure 1, a representative 8-10 year old child and an adult are shown carrying a large bucket unilaterally and a pair of small buckets bilaterally as an example of the experimental protocol. Nine conditions were analyzed: unilateral large 0% BW, unilateral large 10% BW, unilateral large 20% BW, unilateral small 0% BW, unilateral small 10% BW, unilateral small 20% BW, bilateral small 0% BW, bilateral small 10% BW, and bilateral small 20% BW buckets. Buckets were carried unilaterally with the self-selected dominant hand, and the load was evenly split between the buckets during the bilateral conditions. Three repetitions of the nine conditions were completed, and the order of the trials was balanced to reduce potential effects of learning and fatigue.

2.3. Lower extremity kinematics

Participants carried the buckets six metres at a self-selected walking speed while reflective markers were tracked by an eight-camera Peak Motus motion analysis system (Centennial, CO). The marker set included bilateral toes, heels, lateral malleoli, medial malleoli, distal calves, lateral knee joint lines, medial knee joint lines, distal thighs, greater trochanters, and acromion processes. In addition, a single marker was placed at the L5/S1 interspace and at the supersternale. A static standing trial was captured with the full marker set and then the medial malleoli and medial knee joint line markers were removed to avoid contact while carrying buckets. The relative marker positions for the medial malleoli and medial knee joint lines were determined in reference to the lateral malleoli, distal calves, and lateral knee joint lines. During the dynamic experimental trials, the medial malleoli and medial knee joint line marker positions were recreated using these static standing trial transformations. The video data were collected at 120 Hz and noise was reduced with a fourth-order, low-pass Butterworth filter at a cut-off frequency of 6 Hz.

Lower extremity kinematics were analyzed during the stance phase of the dominant foot (side of unilaterally carried loads) near the halfway point of the six metre walking path. Initiation of the stance phase was detected when either the dominant toe or heel marker lowered from the swing phase to a point within one centimetre of the vertical static trial positions of these markers. Similarly, termination of the stance phase was detected with both the dominant toe and heel marker lifted to a point one centimetre above the static trial vertical marker positions. The ankle joint centres were estimated as the midpoint of the lateral and medial malleoli and the knee joint centres as the midpoint of the lateral and medial knee joint lines. Right and left trochanter markers were used to find the midpoint of the hip joints (midhip), and the hip joint centres were located at the midpoint between the trochanters and the midhip. Joint angles were defined as three successive rotations: flexion/extension, abduction/adduction, and internal/external rotation. The static standing trial was used to define the anatomical neutral position, which was used to adjust the joint angle values calculated for the experimental carrying trials. Maximum joint angles for ankle plantarflexion/dorsiflexion, ankle inversion/eversion, knee flexion/extension, hip flexion/extension, hip adduction/abduction, and hip internal/external rotation during the stance phase were determined. Ankle sagittal range of motion (ROM), ankle frontal ROM, knee sagittal ROM, hip sagittal ROM, hip frontal ROM, and hip transverse ROM were each determined as the difference between the maximum joint angles in opposing directions. All calculations were performed in MatLab (Natick, MA).

2.4. Data analysis

Trials with discontinuities due to marker obscurement were eliminated from the analysis, with total joint ROM and maximum joint angles averaged across remaining trials for each

participant. Multivariate ANOVA (SPSS, Chicago, IL) was used to test for main effects of age group, load amount, load symmetry and their interactions. The significance level was set to $p < 0.05$ with a Bonferroni correction of six (number of dependent variables) for the total joint ROM analysis and twelve for the maximum joint angle analysis. The majority of the variables were not normally distributed about the mean values as indicated by Shapiro-Wilk tests of normality ($p > 0.05$). Therefore, when significant main effects were found, Mann-Whitney U nonparametric comparisons were made at a significance level of $p < 0.05$. Adults were compared to the 8-10, 12-14, and 15-17 year olds to test the effects of age group on total joint ROM and maximum joint angles. The 0% BW trials were compared to the 10% BW and 20% BW trials to test the effects of load amount. Finally, the unilateral small bucket trials were compared to the unilateral large bucket trials and the bilateral small bucket trials to test the effects of load symmetry.

3. Results

Total joint ROMs were significantly dependent upon age group ($p < 0.01$) and load amount ($p < 0.01$), but were not dependent upon load symmetry ($p = 0.17$) or any interactions between comparisons ($p = 0.28$ and higher). Similarly, maximum joint angles were significantly dependent upon age group ($p < 0.01$) and carrying condition ($p < 0.01$), but were not dependent upon load symmetry ($p = 0.16$) or any interactions between comparisons ($p = 0.99$).

3.1. Effects of age group on total joint ROM

There were significant differences in ankle, knee, and hip sagittal plane ROM between the child/adolescent age groups and the adults (Figure 2). Ankle sagittal plane ROM was significantly lower for 8-10 year olds ($p = 0.013$) and significantly greater for 15-17 year olds ($p = 0.011$) as compared to adults. Knee sagittal plane ROM was significantly lower for 12-14 ($p = 0.042$) and 15-17 ($p = 0.028$) year olds as compared to adults. Hip sagittal plane ROM was significantly greater for 8-10 year olds ($p < 0.001$) as compared to adults. In addition, there were significant differences in ankle frontal and hip transverse plane ROM between children/adolescents and adults (Figure 3). Ankle frontal plane ROM was significantly greater for 8-10 ($p = 0.025$) and 15-17 ($p = 0.009$) year olds as compared to adults. Hip transverse plane ROM was significant greater for 8-10 ($p < 0.001$), 12-14 ($p = 0.047$), and 15-17 ($p < 0.001$) year olds as compared to adults. No significant differences between children/adolescents and adults were found for hip frontal plane ROM.

3.2. Effects of load amount on total joint ROM

There were significant differences in hip frontal and transverse plane ROM as a function of load amount (Figure 4). Hip frontal plane ROM was significantly lower when carrying a 20% BW load ($p = 0.003$) than when carrying a 0% BW load. In contrast, hip transverse plane ROM was significantly greater when carrying a 20% BW load ($p = 0.004$) than when carrying a 0% BW load. No significant differences in ROM were found at any joint when comparing a 10% BW load to a 0% BW load. In addition, no significant differences were found for ankle ROM, knee ROM, or hip sagittal plane ROM as a function of load amount.

3.3. Effects of age group on maximum joint angles

There were significant differences in maximum ankle plantarflexion/dorsiflexion, knee flexion/extension, and hip flexion/extension between the child/adolescent age groups and the adults (Figure 5). Maximum ankle plantarflexion was significantly lower for 8-10 year olds ($p < 0.001$) and was significantly greater for 15-17 year olds ($p = 0.010$) as compared to adults. Maximum ankle dorsiflexion was significantly greater for 8-10 year olds ($p < 0.001$) as compared to adults. Maximum knee flexion was significantly lower for 8-10 ($p = 0.008$), 12-14 ($p < 0.001$), and 15-17 ($p < 0.001$) year olds as compared to adults. Maximum knee extension

was significantly greater for 12-14 year olds ($p < 0.001$) as compared to adults. Maximum hip flexion was significantly greater for 8-10 year olds ($p = 0.007$) and significantly lower for 12-14 ($p < 0.001$) and 15-17 ($p = 0.005$) year olds as compared to adults. Maximum hip extension was significantly greater for 8-10 ($p < 0.001$), 12-14 ($p < 0.001$), and 15-17 ($p < 0.001$) year olds as compared to adults.

There were also significant differences in maximum ankle inversion/eversion, hip adduction/abduction, and hip internal/external rotation between the child/adolescent age groups and the adults (Figure 6). Maximum ankle inversion was significantly lower for 12-14 year olds ($p = 0.004$) as compared to adults. Maximum ankle eversion was significantly greater for 8-10 ($p < 0.001$) and 12-14 ($p = 0.038$) year olds as compared to adults. Maximum hip adduction was significantly greater for 8-10 ($p < 0.001$), 12-14 ($p < 0.001$), and 15-17 ($p < 0.001$) year olds as compared to adults. In contrast, maximum hip abduction was significantly lower for 8-10 ($p < 0.001$), 12-14 ($p < 0.001$), and 15-17 ($p = 0.008$) year olds as compared to adults. Maximum hip internal rotation was significantly greater for 8-10 ($p < 0.001$), 12-14 ($p < 0.001$), and 15-17 ($p < 0.001$) year olds as compared to adults. In contrast, maximum hip external rotation was significantly lower for 8-10 ($p < 0.001$), 12-14 ($p < 0.001$), and 15-17 ($p = 0.001$) year olds than for adults.

3.4. Effects of load amount on maximum joint angles

There were significant differences in maximum knee extension and hip flexion/extension as a function of load amount (Figure 7). Maximum knee extension ($p = 0.021$) was significantly greater when carrying a 20% BW load. Maximum hip flexion ($p < 0.001$) was significantly greater and extension ($p < 0.001$) was significantly lower when carrying a 20% BW load. In addition, there were significant differences in maximum hip adduction and internal/external rotation as a function of load amount (Figure 8). Maximum hip adduction ($p = 0.003$) was significantly lower when carrying a 20% BW load. Maximum hip internal rotation ($p < 0.001$) was significantly greater and external rotation ($p = 0.031$) was significantly lower when carrying a 20% BW load. No significant differences in maximum joint angles were found when comparing a 10% to a 0% BW load. In addition, no significant differences were found for maximum ankle plantarflexion/dorsiflexion, knee flexion, ankle inversion/eversion, and hip abduction as a function of load amount.

4. Discussion

The purpose of this study was to test the effects of age, load amount, and load symmetry on lower extremity kinematics. The first hypothesis was supported in that the 8-10 year old age group had significantly greater hip sagittal plane ROM and maximum hip flexion angles when compared to adults (Figures 2, 5). One explanation for this difference as a function of age may be greater use of hip flexors and extensors by children during gait (Sutherland 1997). It was also found that the 8-10 year old age group had significantly greater hip extension angles as compared to adults, which further supports the greater use of hip flexors/extensors (Figure 5). In a study of unloaded gait, Ganley and Powers (2005) did not find significant differences in maximum hip flexion or hip extension when comparing 7 year old children to adults. This discrepancy in results may be due to the fact that the weight of the empty bucket itself was carried even in the 0% BW condition in the present study, and that an impeded arm swing may have had more of an effect on the 8-10 year old group than on adults.

The second hypothesis was also supported since there were significant increases in ankle frontal plane ROM and maximum ankle eversion angles comparing 8-10 year olds to adults (Figures 3, 6). In addition, the 8-10 year old age group displayed increased maximum ankle dorsiflexion and decreased ankle plantarflexion as compared to adults (Figure 5). Reduced ankle plantarflexor moment generation and reduced ankle power absorption/generation as observed

during gait in 7 year old children seems like a plausible explanation (Ganley and Powers 2005), but it would require further kinetic analyses to confirm this. Maximum ankle eversion angles were significantly greater for both the 8-10 and 12-14 year old age groups as compared to adults (Figure 6). Increases in maximum joint angles are of particular concern when they approach the limits of joint motion. As a point of comparison, the increases in maximum hip flexion, hip extension, and ankle dorsiflexion angles mentioned thus far all remain within values commonly observed during the stance phase of running (Hamill and Knutzen 2009). While ankle pronation was not measured directly, ankle eversion can be viewed as a partial surrogate measure of pronation. Ankle pronation and internal rotation of the knee/tibia are mechanically linked due to the orientation of the subtalar joint (Inman 1976). Knee and tibial internal rotations have frequently been implicated in tissue strain and pain at the knee (James *et al.* 1978, Bellchamber and van den Bogert 2000) and have discriminated between healthy knees and ACL-deficient or reconstructed knees during walking (Ristanis *et al.* 2003, Andriacchi and Dyrby 2005), between subjects with and without patellofemoral pain (Salsich and Perman 2007), and (in a prospective study) between healthy runners and those who developed iliotibial band syndrome (Noehren *et al.* 2007). In the context of the present study population, greater ankle eversion may be a precursor to knee pain from repetitive load carrying tasks.

Perhaps the most interesting differences in movements as a function of age group occurred at the hip joint in the frontal and transverse planes of motion (Figures 3, 6). The 8-10, 12-14, and 15-17 year old age groups all displayed greater maximum hip adduction angles than adults (Figure 6). These adjustments may be used to reduce stride width in response to the frontal plane balance challenges associated with the carrying tasks. In addition, the 8-10, 12-14, and 15-17 year old age groups all displayed greater maximum hip internal rotation angles than adults (Figure 6). Hip transverse plane ROM and maximum hip internal rotation angles also increased across age groups when carrying a 20% BW load as compared to a 0% BW load (Figures 4, 8). In contrast, a study of adolescent girls carrying dual-strap backpacks did not find significant increases in hip transverse plane ROM with increased carried load (Chow *et al.* 2006). This apparent discrepancy may be attributed to asymmetrical loading for the majority of carrying conditions in the present study, where frontal and transverse plane adjustments are likely to have been more critical. The maximum hip internal rotation angles for the 8-10 year old age group across carrying tasks (Figure 6) and for the 20% BW loads across age groups (Figure 8) appear to fall outside commonly observed values for walking and running (Hamill and Knutzen 2009). These increases in maximum hip internal rotation angles are of concern, and further kinetic analysis is suggested to test if these movement changes are associated with increased torsional loading at the hip joint.

There were several limitations to this study. One limitation was potential kinematic differences between males and females, which might be further complicated by differing rates of maturation. Significant differences between males and females for hip flexion and knee extension angles during walking have been reported (Kerrigan *et al.* 1998; Oberg and Karsznia 1994). Thus, different carrying guidelines might be appropriate for males and females, although a larger research participant pool would be needed to separate out movement changes as a function of age, maturation, and gender. A second limitation was the use of total joint ROM and maximum joint angles as movement parameters since changes between age groups and carrying conditions tended to be subtle. While a kinematic-based analysis offers flexibility for potential data collection by video cameras or observation in the field, changes in joint angles need to be distinct enough to be detectable outside laboratory conditions. A third limitation also relates to the use of kinematic parameters to predict injury risk, since increases in range of motion may not result in clinically relevant increases in bone and soft tissue loading. In this study, lower extremity kinematics were analyzed during the stance phase of walking, which does ensure that the increases in range of motion are occurring during a time period in which

the joints are simultaneously exposed to loading associated with supporting body weight plus the carried load. However, further kinetic analyses of lower extremity joint moments, estimations of joint contact forces, and electromyography would provide measures that are more likely to be directly related to injury risk. A fourth limitation was the idealized experimental conditions, such as the smooth/dry floor and stability of the load being carried. Carrying water over rough/muddy ground would be a greater neuromuscular challenge and could cause greater kinematic differences between age groups and load amounts while probably representing more realistic task conditions.

There are several points of interest that can be taken from this study. Maximum hip internal rotation angles were greater for each of the child/adolescent age groups as compared to adults. In addition, maximum hip internal rotation angles significantly increased when carrying a 20% BW load. Previous studies have suggested load-carrying guidelines at the point where significant changes occur in kinematic measures. Hong and Brueggemann (2000) suggested that backpack loads should not exceed 10% BW based on the observation that trunk forward lean was similar to unloaded gait at 10% BW, but significantly increased at 15% BW. Similarly, Hong and Cheung (2003) suggested that backpack loads should not exceed 15% BW since significant increases in trunk inclination as compared to unloaded gait occurred at 20% BW loads. Applying this line of thought to the present study, suggested guidelines for unilaterally carried loads would be set at a level below 20% BW, where significant increases in maximum hip internal rotation were found to occur. As a measure of sensitivity, the 95% entire confidence interval for changes in maximum hip internal rotation was positive at 20% BW, with increases ranging from 0.8°-12.4°. This observation was also consistent across participants in this study with 100% of 8-10 year olds, 100% 12-14 of year olds, 89% of 15-17 year olds, and 86% of adults displaying increases in maximum hip internal rotation at 20% BW. By setting carried load amounts as a percentage of body weight, the experimental protocol in this study implemented carrying guidelines. Average loads for carrying feed and water measured in the field have been reported to be 83 N (Allread *et al.* 2004). In comparison, the 8-10 year old group carried less than this amount on average at 20% BW, while the other age groups carried greater than this amount. The observation that the 8-10 year old age group was tested at loads less than those measured in the field and still displayed the highest maximum hip internal rotation angles suggests a particular necessity in setting carrying guidelines for the youngest children. However, epidemiologic data combined with further field observations and kinetic measurements would be required before a causal link could be directly established between changes in lower extremity kinematics and increased risk of injury.

5. Conclusions

In this study, lower extremity kinematics were analyzed during bucket carrying tasks to determine the effects of age, load amount, and load symmetry. The child/adolescent age groups displayed significantly greater maximum hip extension, hip adduction, and hip internal rotation angles as compared to adults. In addition, 20% BW load amounts resulted in significantly greater maximum hip internal rotation angles as compared to 0% BW load amounts. These increases in maximum hip internal rotation angles are of concern since they fell outside ranges commonly seen in gait studies and may indicate increased torsional loading at the hip joint. In addition, setting carrying guidelines for the 8-10 year old group appeared to be particularly critical since this age group was tested at lower loads than commonly observed in the field, but still displayed the greatest maximum hip internal rotation angles.

References

- Allread WG, Waters TR. Interventions to reduce low-back injury risk among youth who perform feed handling and scooping tasks on farms. *Journal of Agricultural Safety and Health* 2007;13(4):375–393. [PubMed: 18075014]
- Allread WG, Wilkins JR III, Waters TR, Marras WS. Physical demands and low-back injury risk among children and adolescents working on farms. *Journal of Agricultural Safety and Health* 2004;10(4): 257–273. [PubMed: 15603225]
- Andriacchi TP, Dyrby CO. Interactions between kinematics and loading during walking for the normal and ACL deficient knee. *Journal of Biomechanics* 2005;38:293–298. [PubMed: 15598456]
- Bancej C, Arbuckle T. Injuries in Ontario farm children: A population based study. *Injury Prevention* 2000;6:135–140. [PubMed: 10875671]
- Bartels S, Niederman B, Waters TR. Job hazards for musculoskeletal disorders for youth working on farms. *Journal of Agricultural Safety and Health* 2000;6(3):191–201. [PubMed: 11202113]
- Bellchamber TL, van den Bogert AJ. Contributions of proximal and distal moments to axial tibial rotation during walking and running. *Journal of Biomechanics* 2000;33:1397–1403. [PubMed: 10940398]
- Chow DHK, Kwok MLY, Au-Yang ACK, Holmes AD, Cheng JCY, Yao FYD, Wong MS. The effect of load carriage on the gait of girls with adolescent idiopathic scoliosis and normal controls. *Medical Engineering and Physics* 2006;28:430–437. [PubMed: 16125994]
- Crowe A, Samson MM. 3-D analysis of gait: The effects upon symmetry of carrying a load in one hand. *Human Movement Science* 1997;16:357–365.
- Ganley KJ, Powers CM. Gait kinematics and kinetics of 7-year old children: a comparison to adults using age-specific anthropometric data. *Gait and Posture* 2005;21:141–145. [PubMed: 15639392]
- Gerberich SG, Gibson RW, French LR, Renier CM, Lee TY, Carr WP, Shutske J. Injuries among children and youth in farm households: Regional Rural Injury Study-I. *Injury Prevention* 2001;7:117–122. [PubMed: 11428558]
- Gillette JC, Stevermer CA, Meardon SA, Derrick TR, Schwab CV. Upper extremity and lower back moments during carrying tasks in farm children. *Journal of Applied Biomechanics* 2009;25:149–155. [PubMed: 19483259]
- Hamill, J.; Knutzen, KM. *Biomechanical Basis of Human Movement*. 3rd. Philadelphia: Lippincott, Williams & Wilkins; 2009.
- Hong Y, Brueggemann G. Changes in gait patterns in 10-year-old boys with increasing loads when walking on a treadmill. *Gait and Posture* 2000;11:254–259. [PubMed: 10802438]
- Hong Y, Cheung C. Gait and posture responses to backpack load during level walking in children. *Gait and Posture* 2003;17:28–33. [PubMed: 12535723]
- Inman, VT. *The Joints of the Ankle*. Baltimore: Williams & Wilkins; 1976.
- James SL, Bates BT, Osternig LR. Injuries to runners. *American Journal of Sports Medicine* 1978;6:40–50. [PubMed: 25589]
- Kerrigan DC, Todd MK, Croce UD. Gender differences in joint biomechanics during walking. *American Journal of Physical Medicine and Rehabilitation* 1998;77(1):2–7. [PubMed: 9482373]
- Lee, B.; Marlenga, B. *Professional Resource Manual: North American Guidelines for Children's Agricultural Tasks*. Marshfield, WI: Marshfield Clinic; 1999.
- Marlenga B, Pickett W, Berg RL. Agricultural work activities reported for children and youth on 498 North American farms. *Journal of Agricultural Safety and Health* 2001;7(4):241–252. [PubMed: 11787753]
- Miller ME, Kaufman JD. Occupational injuries among adolescents in Washington state, 1988–1991. *American Journal of Industrial Medicine* 1998;34:121–132. [PubMed: 9651621]
- Noehren B, Davis IS, Hamill J. Prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clinical Biomechanics* 2007;22:951–956. [PubMed: 17728030]
- Nottrodt JW, Manley P. Acceptable loads and locomotor patterns selected in different carriage methods. *Ergonomics* 1989;32(8):945–957. [PubMed: 2806225]
- Oberg T, Karsznia A. Joint angle parameters in gait: Reference data for normal subjects, 10–79 years of age. *Journal of Rehabilitation Research and Development* 1994;31(3):199–209. [PubMed: 7965878]

- Pascoe DD, Pascoe DE, Wang TW, Shim D, Chang CK. Influence of carrying book bags on gait cycle and posture of youths. *Ergonomics* 1997;40(6):631–641. [PubMed: 9174414]
- Pickett W, Brison RJ, Niezgoda H, Chipman ML. Nonfatal farm injuries in Ontario: A population-based survey. *Accident Analysis and Prevention* 1995;27(4):425–433. [PubMed: 7546057]
- Ristanis S, Giakas G, Papageorgiou CD, Moraiti T, Stergiou N, Georgoulis AD. The effects of anterior cruciate ligament reconstruction on tibial rotation during pivoting and after descending stairs. *Knee Surgery, Sports Traumatology, Arthroscopy* 2003;11:360–365.
- Rivara FP. Fatal and nonfatal farm injuries to children and adolescents in the United States. *Pediatrics* 1985;76(4):567–573. [PubMed: 4047800]
- Reed DB, Claunch DT. Nonfatal farm injury incidence and disability to children: a systematic review. *American Journal of Preventative Medicine* 2000;18(4S):70–79.
- Salsich GB, Perman WH. Patellofemoral joint contact area is influenced by tibiofemoral rotation alignment in individuals who have patellofemoral pain. *Journal of Orthopaedic and Sports Physical Therapy* 2007;37:521–528. [PubMed: 17939611]
- Sutherland D. The development of mature gait. *Gait and Posture* 1997;6:163–170.
- USDA (United States Department of Agriculture); National Agricultural Statistical Service. 2001 Childhood Agricultural-Related Injuries. 2004 [27 June 2008]. Available online from: www.cdc.gov/nass
- Zijlstra W, Prokop T, Berger W. Adaptability of leg movements during normal treadmill walking and split-belt walking in children. *Gait and Posture* 1996;4:212–221.



Figure 1. A representative 8-10 year old child and an adult are shown in the unilateral large bucket carrying condition and in the bilateral small bucket carrying condition.




Figure 2. Ankle, knee, and hip sagittal ROM as a function of age group. Total joint ROM are reported as averages across carrying conditions with standard deviations. Statistical differences in total joint ROM for the child/adolescent age groups as compared to the adult age group are denoted by an * ($p < 0.05$).




Figure 3. Ankle frontal and hip transverse ROM as a function of age group. Total joint ROM are reported as averages across carrying conditions with standard deviations. Statistical differences in total joint ROM for the child/adolescent age groups as compared to the adult age group are denoted by an * ($p < 0.05$).




Figure 4. Hip frontal and transverse ROM as a function of load amount. Total joint ROM are reported as averages across age groups with standard deviations. Statistical differences in total joint ROM for the 10% and 20% BW loads as compared to the 0% BW loads are denoted by an * ($p < 0.05$).




Figure 5. Maximum ankle, knee, and hip flexion/extension angles as a function of age group. Maximum joint angles are reported as averages across carrying conditions with standard deviations. Labels above the graphs indicate the positive and negative angle designations. Statistical differences in maximum joint angles for the child/adolescent age groups as compared to the adult age group are denoted by an * ($p < 0.05$).




Figure 6. Maximum ankle inversion/eversion, hip adduction/abduction, and hip internal/external rotation angles as a function of age group. Maximum joint angles are reported as averages across carrying conditions with standard deviations. Labels above the graphs indicate the positive and negative joint angle designations. Statistical differences in maximum joint angles for the child/adolescent age groups as compared to the adult age group are denoted by an * ($p < 0.05$).




Figure 7. Maximum knee and hip flexion/extension angles as a function of load amount. Maximum joint angles are reported as averages across age groups with standard deviations. Statistical differences in maximum joint angles for the 10% and 20% BW loads as compared to the 0% BW loads are denoted by an * ($p < 0.05$).




Figure 8. Maximum hip adduction/abduction and internal/external rotation angles as a function of load amount. Maximum joint angles are reported as averages across age groups with standard deviations. Statistical differences in maximum joint angles for the 10% and 20% BW loads as compared to the 0% BW loads are denoted by an * ($p < 0.05$).

Table 1

Subject characteristics. Mean values are reported for each age group with standard deviations.

Age Group	Gender (M:F)	Age (years)	Height (m)	Mass (kg)
8-10 years	6:1	9.0 ± 1.0	1.37 ± 0.09	31.2 ± 6.4
12-14 years	6:4	12.4 ± 0.8	1.55 ± 0.07	47.1 ± 7.9
15-17 years	5:4	16.0 ± 0.9	1.67 ± 0.10	61.1 ± 9.7
Adult	9:7	23.2 ± 1.6	1.76 ± 0.09	72.4 ± 12.9