

9-23-2002

Diseases are killing grasshoppers

Marlin E. Rice

Iowa State University, merice@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Rice, Marlin E., "Diseases are killing grasshoppers" (2002). *Integrated Crop Management News*. 1765.
<http://lib.dr.iastate.edu/cropnews/1765>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Diseases are killing grasshoppers

Abstract

Grasshoppers in Iowa are now dying from infection by several insect pathogens. These pathogens are causing disease that may devastate grasshopper populations in corn and soybean fields before the first frost.

Entomophaga fungus is a widespread and common pathogen of grasshoppers in North America. The advanced stage of the disease is easy to recognize. Shortly before the grasshopper dies, it climbs to the top of a fencepost, cornstalk, or other plant where it hangs on in a death grip. It typically dies with its head pointed upward and the carcass often remains in this position for several weeks until it deteriorates.

Keywords

Entomology

Disciplines

Agricultural Science | Agriculture | Entomology

INTEGRATED CROP MANAGEMENT

Diseases are killing grasshoppers

Grasshoppers in Iowa are now dying from infection by several insect pathogens. These pathogens are causing disease that may devastate grasshopper populations in corn and soybean fields before the first frost.

Entomophaga fungus is a widespread and common pathogen of grasshoppers in North America. The advanced stage of the disease is easy to recognize. Shortly before the grasshopper dies, it climbs to the top of a fencepost, cornstalk, or other plant where it hangs on in a death grip. It typically dies with its head pointed upward and the carcass often remains in this position for several weeks until it deteriorates.



Differential grasshopper killed by a fungal pathogen.

[Enlarge](#) [1]

Bacteria also kill grasshoppers but bacterial diseases are less well documented in Iowa. Recent, dead grasshoppers may have ruptured abdomens and ooze body fluids. These grasshoppers are less likely to be seen than the fungal-infected grasshoppers because they do not climb upward and clutch plants before they die.



Differential grasshopper killed, probably by a bacterial pathogen.

[Enlarge](#) [2]

As you walk field edges during the next couple of weeks, take a look for dead grass-hoppers

on fencing wire and weeds. These dead grasshoppers are a common, but not often noticed, symptom of natural mortality.

This article originally appeared on page 178 of the IC-488(21) -- September 23, 2002 issue.

Source URL:

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/2002/9-23-2002/ghopkill.html>

Links:

[1] http://www.ent.iastate.edu/imagegal/orthoptera/differential_ghop_fung.html

[2] http://www.ent.iastate.edu/imagegal/orthoptera/differential_ghop_bact.html

IOWA STATE UNIVERSITY
University Extension