

6-24-2002

Stink bugs can twist corn

Marlin E. Rice

Iowa State University, merice@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Rice, Marlin E., "Stink bugs can twist corn" (2002). *Integrated Crop Management News*. 1825.
<http://lib.dr.iastate.edu/cropnews/1825>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Stink bugs can twist corn

Abstract

Stink bug damage to corn occurs widely across Iowa but is mostly seen in fields with winter annual weeds or in corn plants adjacent to weedy fence rows. Basic symptoms of stink bug injury are stunted plants that may grow lateral shoots; wrinkled leaves; holes of various sizes, scattered randomly or in repeating patterns often with a yellow halo; and whorl leaves wrapped tight and failing to expand. Injury is most common on newly emerged plants through the 4th true-leaf stage, but it also may occur on mid-whorl-stage plants.

Keywords

Entomology

Disciplines

Agricultural Science | Agriculture | Entomology

INTEGRATED CROP MANAGEMENT

Stink bugs can twist corn

Stink bug damage to corn occurs widely across Iowa but is mostly seen in fields with winter annual weeds or in corn plants adjacent to weedy fence rows. Basic symptoms of stink bug injury are stunted plants that may grow lateral shoots; wrinkled leaves; holes of various sizes, scattered randomly or in repeating patterns often with a yellow halo; and whorl leaves wrapped tight and failing to expand. Injury is most common on newly emerged plants through the 4th true-leaf stage, but it also may occur on mid-whorl-stage plants. Stink bugs pierce the side of the stalk with their beak. Saliva injected into the leaf during feeding creates holes from pinhole diameter up to 1 inch on expanded leaves. Holes are often surrounded by dead, brown tissue and a yellow halo. Feeding patterns are often repeated across the leaf, possibly giving the appearance of severe shot hole injury from European corn borer larvae. Injured leaves are often twisted giving the plant a “buggy whip” appearance and terminal leaves may fail to fully develop because of the twisting. Heavily injured plants that are not killed by the feeding grow new lateral shoots (tillers). Insecticide control is rarely warranted because most injury is diagnosed after the problem has occurred.

Images of brown stink bugs and injury to corn may be viewed at [here](#) [1].

This article originally appeared on page 119 of the IC-488(14) -- June 24, 2002 issue.

Source URL:

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/2002/6-24-2002/stinkbugtwist.html>

Links:

[1] <http://www.ipm.iastate.edu/ipm/icm/1999/6-28-1999/midscorninj.html>

IOWA STATE UNIVERSITY

University Extension