

2001

Planting Date and Seed Treatment Effects on Soybean Yield

John Lundvall
Iowa State University

Keith Whigham
soy@iastate.edu

Mark E. Westgate
Iowa State University, westgate@iastate.edu

Dale E. Farnham
Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/farms_reports



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Lundvall, John; Whigham, Keith; Westgate, Mark E.; and Farnham, Dale E., "Planting Date and Seed Treatment Effects on Soybean Yield" (2001). *Iowa State Research Farm Progress Reports*. 1825.
http://lib.dr.iastate.edu/farms_reports/1825

This report is brought to you for free and open access by Iowa State University Digital Repository. It has been accepted for inclusion in Iowa State Research Farm Progress Reports by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Planting Date and Seed Treatment Effects on Soybean Yield

Abstract

Today's high-yielding soybean varieties respond favorably to early planting. Multi-year results from statewide university research farms suggest that mid-April to early May planting dates produce top yields in most comparisons. Research farm and on-farm strip trials also suggest that elite varieties yield similarly over a wide range of seeding rates and resulting harvest populations. Based on these results, producers are advised to plant soybeans as soon as spring field conditions allow, with a seeding rate of 150,000 to 175,000 seeds per acre.

Keywords

Agronomy

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences

Planting Date and Seed Treatment Effects on Soybean Yield

John Lundvall, ag specialist, agronomy

Keith Whigham, professor and
extension agronomist

Mark Westgate, associate professor, agronomy

Dale Farnham, assistant professor and
extension agronomist

Introduction

Today's high-yielding soybean varieties respond favorably to early planting. Multi-year results from statewide university research farms suggest that mid-April to early May planting dates produce top yields in most comparisons. Research farm and on-farm strip trials also suggest that elite varieties yield similarly over a wide range of seeding rates and resulting harvest populations. Based on these results, producers are advised to plant soybeans as soon as spring field conditions allow, with a seeding rate of 150,000 to 175,000 seeds per acre.

The yield response of modern varieties to early planting raises questions about expanding the soybean planting season (very early planting of some soybean acres in late March or early April) when weather conditions allow. In 2000 a soybean "planting date x seed treatment" study was initiated to evaluate yield response of two adapted, high-yield varieties to very early planting. Research objectives are to determine whether planting soybeans earlier than corn could be a viable management option for Iowa producers, and if fungicide seed treatments are needed to make this practice profitable.

Establishment of similar studies at four other university research farms statewide afforded yield response comparisons of adapted varieties from five unique soil associations and environments. Soils at the Southeast Research and Demonstration Farm are typical of the Taintor-Mahaska soil association.

Materials and Methods

Adapted, high yield conventional varieties from LG/Callahan (9288 variety, relative maturity 2.8) and Merschman ("Mohave V" variety, relative maturity 2.8) seed companies were tested in 2000. LG/Callahan and Merschman offer fungicide-treated soybean seed. Fungicide seed treatments fight soybean seedling diseases that can weaken or kill early-planted soybeans, causing profit-robbing stand reductions. Each company was asked to furnish their varieties with and without their respective fungicide treatments; therefore, a total of four treatments were compared on each of four planting dates. Experimental plots were planted directly into standing corn stalks using a John Deere 7100 planter with 30-inch row spacing. All plots were planted at a rate of 175,000 seeds per acre. Planting dates included March 30, April 24, May 15, and May 30. Planting dates and treatments were included in a split-plot design with four replications. Main plot treatments were planting dates, and variety/seed treatment combinations were subplot treatments. Plots were machine harvested on October 3. Grain yields (adjusted to 13% moisture) and established plant population estimates are summarized in Tables 1-3.

Results and Discussion

Averaged across varieties, fungicide-treated and untreated soybean yields were statistically similar ($P>0.05$) (Table 1). The lack of yield response to seed treatment was consistent regardless of planting date.

Averaged across treatments, soybean yield performance was statistically similar ($P>0.05$) on planting dates from March 30 to May 15 (Table 2). Yield of three individual treatments was statistically similar ($P>0.05$) on all planting dates; yield of Merschman untreated plots dropped significantly ($P<0.05$) because planting was delayed to May 30 (Table 2). Three

treatments yielded best when planted on May 15.

Producers considering very early soybean planting recognize the risk of stand losses caused by seedling diseases in cooler soils. Several plots planted March 30 and April 24 sustained additional stand losses due to severe bean leaf beetle feeding and subsequent seedling death. Poor germination of the LG/Callahan variety was another factor causing poor stands in very early-planted plots (Table 3).

Dry weather limited development of some soybean seedling diseases in 2000. Regardless

of planting date, fungicide seed treatment did not consistently improve soybean yield performance at the Southeast Research Farm; however, producers are advised to base management decisions on multiple-year data. This study will be continued at the Southeast Research Farm in 2001.

Acknowledgments

The authors wish to thank Matt Hunt and Kevin Van Dee for their assistance in conducting this research. Thanks also to LG/Callahan and Merschman seed companies for providing the seed used in this study.

Table 1. Effect of planting date and seed treatment on soybean yield in 2000 at Crawfordsville, IA.

Experimental treatment	Yield performance by planting date				All planting dates
	March 30	April 24	May 15	May 30	
	------(Bushels/acre)-----				
Untreated seed (control)	47.7 a ¹	48.0 a	49.6 a	46.4 a	47.9 a
Fungicide-treated seed	49.1 a	47.8 a	49.2 a	45.2 a	47.8 a
Mean	48.4	47.9	49.4	45.8	47.9
L.S.D. (P=0.05)	NS ²	NS	NS	NS	NS

¹ Within columns, experimental treatment mean yields followed by different letters are statistically different (P<0.05).

² "NS" indicates no statistically significant (P=0.05) seed treatment effect on soybean yield.

Table 2. Effect of planting date, variety, and seed treatment on soybean yield in 2000 at Crawfordsville, IA.

Experimental treatment (trt.)	Yield performance by planting date				Planting date yield effect
	March 30	April 24	May 15	May 30	
	------(Bushels/acre)-----				(P=0.05)
LG/Callahan with no seed trt. (control)	46.1 b ¹	46.4 c	48.9 a	46.5 a	NS ²
LG/C with fungicide seed trt.	48.5 ab	47.5 b	49.7 a	44.5 a	NS
Merschman with no seed trt. (control)	49.3 a	49.6 a	50.4 a	46.2 a	D1,2,3 > D4
Merschman with fungicide seed trt.	49.7 a	48.1 b	48.7 a	45.9 a	NS
Mean	48.4	47.9	49.4	45.8	NS
L.S.D. (P=0.05)	2.8	1.0	NS ²	NS	

¹ Within columns, experimental treatment mean yields followed by different letters are statistically different (P<0.05).

² "NS" indicates no statistically significant (P=0.05) planting date or experimental trt. effects on soybean yield.

Table 3. Estimated established plant stand levels in 2000 at Crawfordsville, IA.

Experimental treatment (trt.)	March 30	April 24	May 15	May 30
		------(Plants/acre)-----		
LG/Callahan with no seed trt. (control)	99,000	116,000	150,000	130,000
LG/C with fungicide seed trt.	129,000	130,000	152,000	129,000
Merschman with no seed trt. (control)	133,000	136,000	154,000	142,000
Merschman with fungicide seed trt.	132,000	134,000	155,000	145,000