

9-23-2002

Big caterpillar is white-lined sphinx

Marlin E. Rice

Iowa State University, merice@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Entomology Commons](#)

Recommended Citation

Rice, Marlin E., "Big caterpillar is white-lined sphinx" (2002). *Integrated Crop Management News*. 1829.
<http://lib.dr.iastate.edu/cropnews/1829>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Big caterpillar is white-lined sphinx

Abstract

During late August, large caterpillars were seen crawling out of soybean fields and across roads in Iowa. Because of the large size (approximately 2 inches in length) of this caterpillar, it was easy to assume that it was doing some damage to soybean. The caterpillar is called a white-lined sphinx and has a “horn” protruding from the rear of the body. Fortunately, it is not a pest of soybean but feeds solely on weeds.

Keywords

Entomology

Disciplines

Agricultural Science | Agriculture | Entomology

INTEGRATED CROP MANAGEMENT

Big caterpillar is white-lined sphinx

During late August, large caterpillars were seen crawling out of soybean fields and across roads in Iowa. Because of the large size (approximately 2 inches in length) of this caterpillar, it was easy to assume that it was doing some damage to soybean. The caterpillar is called a white-lined sphinx and has a “horn” protruding from the rear of the body. Fortunately, it is not a pest of soybean but feeds solely on weeds. The larva transforms into a sphinx moth, which also is called a hawk moth or hummingbird moth. During late August and September, these moths commonly probe nectar from flowers in gardens.



Caterpillar of the white-lined sphinx moth.

[Enlarge](#) [1]

This article originally appeared on page 177 of the IC-488(21) -- September 23, 2002 issue.

Source URL:

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/2002/9-23-2002/sphinx.html>

Links:

[1] http://www.ent.iastate.edu/imagegal/lepidoptera/sphinx/white_lined_sphinx.html

IOWA STATE UNIVERSITY
University Extension