

11-8-1999

Tolerance results to white mold and SDS

Michael D. Uphoff

Iowa State University, mduphoff@iastate.edu

X. B. Yang

Iowa State University, xbyang@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Uphoff, Michael D. and Yang, X. B., "Tolerance results to white mold and SDS" (1999). *Integrated Crop Management News*. 2224.
<http://lib.dr.iastate.edu/cropnews/2224>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Tolerance results to white mold and SDS

Abstract

The Department of Plant Pathology at Iowa State University conducts an annual white mold tolerance test with funding from the Iowa Soybean Promotion Board. This year, for the first time, we also have tested varieties from commercial companies for tolerance to sudden death syndrome (SDS). The data for 1999 are presented below.

Keywords

Plant Pathology

Disciplines

Agricultural Science | Agriculture | Plant Pathology



Tolerance results to white mold and SDS

The Department of Plant Pathology at Iowa State University conducts an annual white mold tolerance test with funding from the Iowa Soybean Promotion Board. This year, for the first time, we also have tested varieties from commercial companies for tolerance to sudden death syndrome (SDS). The data for 1999 are presented below.

White Mold

The white mold test was conducted at Rudd, Iowa, in a field that was severely infested in 1997. Lines were planted in late May. Each variety was replicated four times in plots four rows wide and 12 feet in length with a row spacing of 13 inches. Disease incidence data were collected in mid-September. The table below lists the average tolerance score for each variety. Tolerance of each variety was measured as the percentage of plants killed. When choosing a variety, try to select one that is consistent over years and locations or that seems to perform well for growers in your area. Keep in mind that yield losses are not significant if less than 20 percent of plants are killed. Varieties with less than 30 percent of the plants killed are considered to have an acceptable level of tolerance. In fields with mild disease pressure, varieties that did not receive a high tolerance rating in our test may do well because our fields had higher disease pressure than most production fields.

Weather in this past season was not ideal for white mold development and disease occurred very late in our plots, even though our test was done in a field that had severe white mold. Therefore, it is important to note that varieties in later maturity groups have greater disease incidence in our test than those of early maturity groups. These varieties should have higher yields and less disease when grown in southern and central Iowa, their normal area of adaptation. Comparisons should be made only among entries within maturity groups.

Sudden Death Syndrome

The SDS tolerance test was conducted in a field near Ames, Iowa. To ensure uniform disease pressure across plots, we artificially infested test plots at planting with a method developed in our laboratory. Lines were planted in plots consisting of two 10-foot-long rows at eight seeds per foot with a row spacing of 30 inches. Lines were replicated three times. Lines were evaluated in early September and each line was given a numerical tolerance score from 1 to 10 based on leaf symptoms with 10 indicating 100 percent defoliation. Low scores indicate a high level of tolerance. Most entries in our tests are tolerant to this disease.

If you observed severe SDS in your field in the past few years, you may want to select varieties that are tolerant to this disease based on the information in the table below. Yield

loss may not be a concern if the tolerance score is less than 3.

| ISU SDS tolerance test results (1999). | | | |
|---|----------------|--------------------------|----------------------|
| Company | Variety | Relative Maturity | Average Score |
| AgriPro Seeds | AP 2002 RR | 2.0 | 1.3 |
| AgriPro Seeds | AP 2415 RR | 2.4 | 2.7 |
| Asgrow Seeds | AG2903 | 2.9 | 4.3 |
| Asgrow Seeds | AG3302 | 3.3 | 1.3 |
| Dairyland Seeds | DSR-275 | 2.7 | 3.7 |
| Dekalb Seeds | CX339C | 2.8 | 0.7 |
| Dekalb Seeds | CX284C | 3.0 | 1.0 |
| Dekalb Seeds | CX302C | 3.3 | 2.7 |
| Growmark | HS 2861 | | 2.7 |
| Growmark | RT 3885 | | 2.7 |
| Latham Seed | 950 Brand | 2.9 | 1.0 |
| Latham Seeds | 640 Brand | 2.3 | 1.0 |
| LG Seeds | LG6222CRR | 2.2 | 1.7 |
| LG Seeds | LG6278CRR | 2.7 | 2.7 |
| Mark Seed | 9824CTA | 2.4 | 2.3 |
| Mark Seed | 97CN22CTB | 2.2 | 1.3 |
| Merchman | Shawnee VIIRR | 2.9 | 3.0 |
| Merchman | Fillmore VRR | 3.4 | 2.0 |
| Naylor Seed | NS 2770 | 2.5 | 1.7 |
| Naylor Seed | NS 2450 | 2.5 | 2.3 |
| Novartis | S33-N1 | 3.3 | 1.3 |
| Novartis | X9925 | 2.5 | 2.3 |
| Ottlie Seed | 8350 | 3.5 | 1.7 |
| Ottlie Seed | 8240 RR | 2.4 | 2.0 |

| | | | |
|---------------------|------------|-----|-----|
| Sands Seed | Exp 2424RR | 2.4 | 1.3 |
| Sands Seed | Exp 3434RR | 3.4 | 2.3 |
| Sieben | SS 3001RR | | 2.7 |
| Sieben | SS 289 | | 4.0 |
| Pioneer Hi-Bred | 93B82 | | 1.7 |
| Pioneer Hi-Bred | 92B91 | | 1.0 |
| Tolerant control | Pharaoh | | 0.3 |
| Tolerant control | Ripley | | 0.3 |
| Susceptible control | Spencer | | 5.0 |
| Susceptible control | P9344 | | 4.0 |

ISU white mold tolerance test results (1999).

| Company | Variety | Relative Maturity | Average Score | Average Yield (bu/acre) |
|-------------------------|-------------|-------------------|---------------|-------------------------|
| Agri-Pro Seeds | AP2002RR | 2.0 | 15.3 | 50.0 |
| Agri-Pro Seeds | AP2415RR | 2.4 | 31.3 | 39.0 |
| Agri-Pro Seeds | AP2220 | 2.2 | 13.8 | 52.2 |
| Agri-Pro Seeds | AP2889 | 2.8 | 21.3 | 52.8 |
| Albert Lea Seed House | NS1903 | 1.9 | 6.5 | 53.5 |
| Albert Lea Seed House | NS 1904 RR | 1.9 | 10.3 | 46.2 |
| Asgrow Seeds | AG1901 | 1.9 | 2.3 | 52.2 |
| Asgrow Seeds | AG2501 | 2.5 | 12.8 | 37.3 |
| Asgrow Seeds | AG2001 | 2.0 | 1.8 | 50.4 |
| Dairyland Seed Co. Inc. | DSR-218 | 2.1 | 4.0 | 50.8 |
| Dairyland Seed Co. Inc. | DSR-215/RR | 2.1 | 2.3 | 47.6 |
| Dairyland Seed Co. Inc. | DSR-220/STS | 2.4 | 5.3 | 46.9 |
| Dairyland Seed Co. Inc. | DSR-293/RR | 2.7 | 8.8 | 35.5 |

| | | | | |
|-------------------------|---------------------|-----|------|------|
| Dekalb Seeds | CX195 | 1.9 | 5.3 | 44.4 |
| Fontanelle Hybrids | 8890 RR | 3.0 | 38.8 | 56.2 |
| Fontanelle Hybrids | 8933 RR | 2.3 | 28.8 | 39.8 |
| Great Lakes Hybrids Inc | GL1715 | 1.7 | 1.3 | 35.5 |
| Great Lakes Hybrids Inc | GL1902RR | 1.9 | 10.3 | 39.9 |
| Growmark | HT 261 STS | | 17.5 | 47.9 |
| Growmark | RT 2175 | | 2.5 | 54.3 |
| Growmark | RT 2587 | | 18.3 | 50.4 |
| Latham Seed Co. | Latham 656RR Brand | 2.3 | 15.0 | 48.7 |
| Latham Seed Co. | Latham 1056RR Brand | 3.0 | 15.3 | 54.6 |
| LG Seed Inc. | LG6200 | 2.0 | 6.3 | 40.7 |
| LG Seed Inc. | LG6222CRR | 2.2 | 13.8 | 46.0 |
| LG Seed Inc. | LG6284RR | 2.8 | 9.0 | 45.8 |
| LG Seed Inc. | LG6283STS | 2.8 | 26.3 | 56.8 |
| Mark Seed Co. | 9519 | 1.9 | 4.0 | 47.3 |
| Mark Seed Co. | 9921 | 2.1 | 14.0 | 49.2 |
| Merschman Seeds | Munsee IIRR | 2.1 | 28.8 | 49.8 |
| Merschman Seeds | Comanche V | 2.1 | 22.5 | 45.4 |
| Merschman Seeds | Mars VRR | 1.9 | 26.7 | 49.8 |
| Merschman Seeds | Apache VIIRR | 2.4 | 15.0 | 43.5 |
| Naylor Seed | NS 2770 | 2.5 | 33.8 | 24.4 |
| Naylor Seed | NS 2450 | 2.5 | 21.3 | 49.4 |
| Naylor Seed | Excel 8261 RR | 2.5 | 18.8 | 43.9 |
| Novartis Seeds Inc. | X9818 | 1.9 | 5.3 | 42.0 |
| Novartis Seeds Inc. | X9923R | 2.3 | 3.8 | 47.5 |
| Novartis Seeds Inc. | X9919R | 1.9 | 12.8 | 44.4 |
| Ottlie RO Seed | Ottlie 8299 | 2.9 | 32.5 | 49.6 |
| Ottlie RO Seed | Ottlie 8240 RR | 2.4 | 30.0 | 43.9 |
| | | | | |

| | | | | |
|-----------------------|--------------------|-----|------|------|
| Pioneer Hi-Bred | 93B11 | | 46.3 | 51.8 |
| Pioneer Hi-Bred | 9306 | | 31.3 | 50.8 |
| Profiseed Inc. | PS 4241 RR | 2.4 | 13.8 | 51.4 |
| Profiseed Inc. | PS 2509 | 2.4 | 21.3 | 51.3 |
| Sand Seed Service | SOI 260 | 2.0 | 3.0 | 49.2 |
| Sand Seed Service | SOI 275RR | 2.7 | 13.8 | 43.9 |
| Sieben | SS289 | | 27.5 | 38.4 |
| Sieben | SS298 | | 46.3 | 47.1 |
| Sieben | SS2601RR | | 11.5 | 41.7 |
| Terra Seeds | TS194 | | 3.8 | 41.1 |
| Wilson Genetics L.L.C | Wilson 2832 RR | 2.8 | 17.5 | 44.2 |
| Wilson Genetics L.L.C | Wilson 2844 RR | 2.8 | 38.8 | 35.3 |
| Wilson Genetics L.L.C | Wilson 3111 RR/SCN | 3.1 | 26.3 | 49.3 |
| | BSR101 | 1.7 | 2.5 | 50.6 |
| | Corsoy79 | 1.5 | 2.5 | 47.4 |
| | A2242 | 2.2 | 17.5 | 48.3 |
| | Kenwood 94 | 2.5 | 6.3 | 49.3 |
| | Williams 82 | 3.8 | 50.0 | 17.4 |

This article originally appeared on pages 191-192 of the IC-482(24) -- November 8, 1999 issue.

Source URL:

<http://www.ipm.iastate.edu/ipm/icm//ipm/icm/1999/11-8-1999/wmsdstolres.html>

IOWA STATE UNIVERSITY
University Extension