

2-10-2016

Spray Volume is Critical for Postemergence Herbicides

Robert G. Hartzler

Iowa State University, hartzler@iastate.edu

H. Mark Hanna

Iowa State University, hmhanna@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>

 Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Bioresource and Agricultural Engineering Commons](#), and the [Weed Science Commons](#)

Recommended Citation

Hartzler, Robert G. and Hanna, H. Mark, "Spray Volume is Critical for Postemergence Herbicides" (2016). *Integrated Crop Management News*. 2328.

<http://lib.dr.iastate.edu/cropnews/2328>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Spray Volume is Critical for Postemergence Herbicides

Abstract

Proper herbicide application is critical to ensure products achieve their full potential for weed control. For postemergence herbicides, this includes application to appropriately sized weeds and equipping the sprayer to achieve uniform coverage of target weeds. Weed scientists at Purdue University recently reported on the influence of nozzle type and spray volume on target coverage within the soybean canopy. A traditional flat fan (XR) nozzle and three drift reduction nozzles were included in the study. The XR nozzle produced more driftable droplets than the other nozzles, and the two types of air induction nozzles had much less volume in small droplets than the XR and TTJ nozzles

Keywords

Agronomy, Agricultural and Biosystems Engineering

Disciplines

Agricultural Science | Agriculture | Agronomy and Crop Sciences | Bioresource and Agricultural Engineering
| Weed Science

IOWA STATE UNIVERSITY
Extension and Outreach
Integrated Crop Management

[Home](#)

Mailing Lists

Subscribe to ICM News updates and receive email alerts when new information is posted.

Your Email address *

Spray Volume is Critical for Postemergence Herbicides

ICM News

February 10, 2016

Proper herbicide application is critical to ensure products achieve their full potential for weed control. For postemergence herbicides, this includes application to appropriately sized weeds and equipping the sprayer to achieve uniform coverage of target weeds.

Weed scientists at Purdue University recently reported on the influence of nozzle type and spray volume on target coverage within the soybean canopy. A traditional flat fan (XR) nozzle and three drift reduction nozzles were included in the study. The XR nozzle produced more driftable droplets than the other nozzles, and the two types of air induction nozzles had much less volume in small droplets than the XR and TTJ nozzles (Figure 1).




Figure 1. Effect of nozzle type and size on production of driftable droplets. Legleiter and Johnson. 2016.

Coverage of targets within the canopy was evaluated by placing water sensitive paper at different heights within 12" tall soybean planted in 15" rows. Averaged over nozzle types and spray volume, coverage declined by approximately 50% from the top of the canopy to targets placed 4" above the soil surface.

The TTI nozzle was less consistent than the other nozzles on coverage at the top and middle of the canopy. The TTI nozzle produces the largest droplet size of the nozzles evaluated, and thus would be expected to provide less uniform coverage than the others. However, nozzle type did not have a significant affect on coverage of targets at the bottom of the canopy. The authors speculated that large droplets were more efficient at penetrating the canopy than small droplets, therefore resulting in similar coverage by the different nozzles of targets placed at the base of the canopy.

Regardless of nozzle type or target placement, better coverage was achieved with the larger nozzle size that provided greater gallons per acre (GPA) spray volume. Averaged over nozzle types and the two years of the study, 15 GPA provided 26% coverage compared to 13% at 10 GPA.

Thorough coverage of weeds is required to achieve consistent weed control, especially when relying on herbicides other than glyphosate. The concern over off-target movement of pesticides has led to the movement towards nozzle types that produce fewer small droplets. This research demonstrates that spray volume has a greater affect on spray coverage than the type of nozzle. Using spray volumes at the high end of the recommended range should improve the consistency of weed control. Nozzle selection

should be based on label specifications.

Reference: Legleiter, T.R. and W.G. Johnson. 2016. Herbicide coverage in narrow row soybean as influenced by spray nozzle design and carrier volume. *Crop Protec.* 83:1-8.ods

Category: Weeds

Tags:

herbicide application

spray nozzles

herbicide

postemergence herbicides

postemergence application

Authors:



Bob Hartzler *Professor of Agronomy*

Dr. Bob Hartzler is a professor of agronomy and an extension weed specialist. He conducts research on weed biology and how it impacts the efficacy of weed management programs in corn and soybean. Dr. Hartzler also teaches undergraduate classes in weed science and weed identificatio...



H Hanna *Scientist II*

Dr. H. Mark Hanna is an extension agricultural engineer with Iowa State University. Hanna's main focus is sustainable agricultural systems, including chemical application, energy consumption, tillage/planting, and harvest. His research focus has been on developing ways for field equipment to en...

[Site Index](#) [Site map](#) [Policies](#)

