

9-10-2019

Nutrient Removal When Harvesting Corn Stover

John E. Sawyer

Iowa State University, jsawyer@iastate.edu

Antonio Mallarino

Iowa State University, apmallar@iastate.edu

Follow this and additional works at: <https://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), and the [Agriculture Commons](#)

Recommended Citation

Sawyer, John E. and Mallarino, Antonio, "Nutrient Removal When Harvesting Corn Stover" (2019).

Integrated Crop Management News. 2589.

<https://lib.dr.iastate.edu/cropnews/2589>

The Iowa State University Digital Repository provides access to Integrated Crop Management News for historical purposes only. Users are hereby notified that the content may be inaccurate, out of date, incomplete and/or may not meet the needs and requirements of the user. Users should make their own assessment of the information and whether it is suitable for their intended purpose. For current information on integrated crop management from Iowa State University Extension and Outreach, please visit <https://crops.extension.iastate.edu/>.

Nutrient Removal When Harvesting Corn Stover

Abstract

Increasing demand to use corn plant biomass for producing energy and various products has spurred interest in harvesting corn stover and specific plant components in addition to grain. Harvesting more biomass means increased carbon (C) and other nutrient removal from fields. What is the nutrient removal when different corn plant components are harvested?

Disciplines

Agricultural Science | Agriculture

IOWA STATE UNIVERSITY

Extension and Outreach

Integrated Crop Management

Nutrient Removal When Harvesting Corn Stover

September 10, 2019

Increasing demand to use corn plant biomass for producing energy and various products has spurred interest in harvesting corn stover and specific plant components in addition to grain. Harvesting more biomass means increased carbon (C) and other nutrient removal from fields. What is the nutrient removal when different corn plant components are harvested?

Corn Plant Nutrient Content

Most producers in Iowa are familiar with harvesting corn for grain, but less familiar with other plant biomass removal. Those harvesting corn silage are aware of the increase in phosphorus (P) and especially potassium (K) removal, and fertilization guidelines consider this increase. For example, with corn silage based on a bushel grain equivalent, the pounds of P as P_2O_5 goes from 0.32 to 0.44 and the pounds of K as K_2O goes from 0.22 to 1.10, respectively, for corn grain and corn silage (15% grain moisture content, ISU Extension publication [PM 1688](#)).

For corn stover the pounds per ton are 4.8 lb P_2O_5 and 18.0 lb K_2O (15% stover moisture content, publication PM 1688). That data indicate there is considerable K in the non-grain part of the corn plant. Calculating the P and K removal in harvested stover is somewhat complicated as nutrients can be leached (especially K) from leaves and corn stalks with rainfall after grain harvest. The effect of rainfall on K is much more important than for P because all plant K is in a soluble inorganic form (K^+) while most P is in organic forms of low solubility. Therefore, the concentration of nutrients at plant maturity will typically be higher than found for baled stover because it often rains after grain harvest. Also, the

frequency and amount of rainfall will affect the concentration remaining. Costs for replacing removed nutrients will vary depending upon prevailing prices and stover removal amounts.

Corn biomass also contains other plant nutrients. Examples for nitrogen (N), P, K, calcium (Ca), magnesium (Mg), and sulfur (S) are shown in Table 1. There is variation in concentration due to differing data sources, hybrids, fertilization levels, etc. While there may typically be adequate supply of nutrients like K, Ca, and Mg from Iowa soils, stover harvest can lead to removal of large amounts of “basic cations” (K^+ , Ca^{+2} , and Mg^{+2}) and result in accelerated decrease in soil pH. Of course fertilization, liming, or manuring soils aids in replacement of these nutrients.

Dry Matter and Nutrient Composition in Corn Plant Components

Silage harvest results in almost complete removal of aboveground plant biomass. Baling corn stover typically does not remove as much plant biomass and amounts removed vary greatly across fields, years, and desired removal level. Also, there may be interest in specific plant component removal, such as targeting corn cobs. Table 1 lists the various corn plant components, the associated dry matter, and the nutrient composition from N rate trials at 14 site-years in Iowa. While not measured in those trials, cob composition for P would be approximately 1.1 lb P_2O_5 /ton cobs and for K 14 lb K_2O /ton cobs (both dry matter based). The corn grain harvest index averaged 51% (percent of the total aboveground biomass as grain on a dry matter basis at maturity). The grain harvest index can be used to estimate non-grain biomass from grain yield. For example, at a grain yield of 200 bu/acre (at 15% moisture standard, the per bu grain dry matter is 47.6 lb) the grain dry matter is 9,520 lb/acre and the non-grain (vegetation plus cob) dry matter is 9,150 lb/acre.

Summary

Harvesting corn plant components in addition to grain does result in greater removal of plant nutrients. Effects of increased P and K removal on nutrient application needs are immediate and should to be accounted for in fertilization plans. Effects on needs of other nutrients such as N and S, liming requirements to maintain desirable soil pH levels, soil organic C, and several physical, chemical, and biological soil properties are less apparent in the short-term but have consequence in the long term. Therefore, consideration of the impact on nutrient cycling, nutrient removal, and soil resources should be a part of the decision process regarding harvesting corn biomass. Additional information on nutrient management related to stover harvest can be found in publication [PM 3052C](#), available from the ISU Extension and Outreach [Extension Store](#).

Table 1. Corn nutrient composition at plant maturity by plant part.

C	
Grain	795
Vegetation	840
Cob	787
N	
Grain	24
Vegetation	12
Cob	10
P (P ₂ O ₅)	
Grain	12
Vegetation	3
Cob	--
K (K ₂ O)	
Grain	8

Vegetation	22
Cob	--
Ca	
Grain	2
Vegetation	9
Cob	--
Mg	
Grain	2
Vegetation	6
Cob	--
S	
Grain	2
Vegetation	1
Cob	--
Zn	

Grain	0.030
Vegetation	0.031
Cob	--
<hr/>	
Mn	
Grain	0.006
Vegetation	0.069
Cob	--
<hr/>	
Cu	
Grain	0.004
Vegetation	0.015
Cob	--
<hr/>	
B	
Grain	0.009
Vegetation	0.015
Cob	--

Fe	
Grain	0.047
Vegetation	0.281
Cob	--

Values are lb/ton (DM)

From 14 site years in 2006-2007.

J.E. Sawyer and D.W. Barker.

Category: Crop Production

Links to this article are strongly encouraged, and this article may be republished without further permission if published as written and if credit is given to the author, Integrated Crop Management News, and Iowa State University Extension and Outreach. If this article is to be used in any other manner, permission from the author is required. This article was originally published on September 10, 2019. The information contained within may not be the most current and accurate depending on when it is accessed.

Crop:

Corn

Tags: corn stover harvest crop nutrients

Authors:



John Sawyer Professor

Dr. John Sawyer is a professor of agronomy and extension specialist in soil fertility and nutrient management at Iowa State University. His extension program involves soil fertility management, efficient crop nutrient utilization, and environmentally sound fertilizer and manure systems. Dr. Sawye...



Antonio Mallarino Professor of Soil Fertility and Nutrient Management, Extension Specialist

Dr. Antonio Mallarino is a professor of agronomy and nutrient management research and an extension specialist at Iowa State University. His programs focus on agronomic and environmental issues of nutrient management with emphasis on phosphorus, potassium, lime, and micronutrients. Issues addresse...